



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 04:09 AM GMT

PDB ID : 1A08
Title : C-SRC (SH2 DOMAIN) COMPLEXED WITH ACE-DIFLUORO PHOSPH
OTYR-GLU-(N,N-DIPENTYLAMINE)
Authors : Shewchuk, L.; Jordan, S.
Deposited on : 1997-12-09
Resolution : 2.20 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3751 (2.20-2.20)
Ramachandran outliers	78287	3681 (2.20-2.20)
Sidechain outliers	78261	3682 (2.20-2.20)
RSRZ outliers	66119	2939 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	107	
1	B	107	
2	C	4	
2	D	4	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2199 atoms, of which 448 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C-SRC TYROSINE KINASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	105	Total	C	H	N	O	S	0	0	0
			1020	520	196	145	156	3			
1	B	104	Total	C	H	N	O	S	0	0	0
			1012	517	194	144	154	3			

- Molecule 2 is a protein called ACE-DIFLUORO PHOSPHOTYR-GLU-(N,N-DIPENTYLAMINE).

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
2	C	4	Total	C	F	H	N	O	P	0	0	0
			43	27	2	2	3	8	1			
2	D	4	Total	C	F	H	N	O	P	0	0	0
			43	27	2	2	3	8	1			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	16	Total	H	O	0	0
			48	32	16		
3	B	7	Total	H	O	0	0
			21	14	7		
3	C	3	Total	H	O	0	0
			9	6	3		
3	D	1	Total	H	O	0	0
			3	2	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: C-SRC TYROSINE KINASE

Chain A: 



- Molecule 1: C-SRC TYROSINE KINASE

Chain B: 



- Molecule 2: ACE-DIFLUORO PHOSPHOTYR-GLU-(N,N-DIPENTYLAMINE)

Chain C: 

There are no outlier residues recorded for this chain.

- Molecule 2: ACE-DIFLUORO PHOSPHOTYR-GLU-(N,N-DIPENTYLAMINE)

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.90Å 67.40Å 75.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.20 36.08 – 2.20	Depositor EDS
% Data completeness (in resolution range)	83.0 (6.00-2.20) 83.1 (36.08-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.15 (at 2.20Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.192 , (Not available) 0.183 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	0 of 11573 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2199	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FTY, DIP, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	0/841	1.77	17/1136 (1.5%)
1	B	1.04	1/835 (0.1%)	1.79	19/1128 (1.7%)
2	C	1.23	0/8	0.93	0/9
2	D	1.43	0/8	0.87	0/9
All	All	1.05	1/1692 (0.1%)	1.77	36/2282 (1.6%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	240	LEU	C-N	-7.29	1.17	1.34

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	220	ARG	NE-CZ-NH2	-12.71	113.95	120.30
1	A	158	ARG	NE-CZ-NH2	-12.62	113.99	120.30
1	B	220	ARG	NE-CZ-NH2	-11.37	114.62	120.30
1	A	158	ARG	NE-CZ-NH1	9.86	125.23	120.30
1	A	216	TYR	CB-CG-CD1	-8.96	115.63	121.00
1	A	230	VAL	CG1-CB-CG2	-8.67	97.03	110.90
1	B	208	ARG	NE-CZ-NH2	-8.34	116.13	120.30
1	B	151	TRP	CE2-CD2-CG	-8.25	100.70	107.30
1	B	151	TRP	CG-CD2-CE3	8.09	141.18	133.90
1	A	216	TYR	CB-CG-CD2	8.04	125.82	121.00
1	A	232	TYR	CB-CG-CD2	-7.91	116.25	121.00
1	B	166	LEU	CA-C-N	-7.86	99.92	117.20
1	B	163	ARG	NE-CZ-NH2	-7.79	116.41	120.30
1	B	151	TRP	CB-CG-CD1	-7.54	117.20	127.00
1	B	208	ARG	NE-CZ-NH1	7.39	123.99	120.30
1	A	182	THR	N-CA-CB	-7.27	96.48	110.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	151	TRP	CD1-CG-CD2	7.00	111.90	106.30
1	B	222	GLN	CA-CB-CG	6.74	128.24	113.40
1	A	149	GLU	C-N-CA	6.46	137.84	121.70
1	B	209	LYS	CA-CB-CG	-6.32	99.49	113.40
1	A	163	ARG	NE-CZ-NH1	6.29	123.44	120.30
1	B	158	ARG	NE-CZ-NH2	-6.25	117.17	120.30
1	B	158	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	A	159	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	B	167	ASN	CA-CB-CG	-5.96	100.29	113.40
1	A	243	ARG	CA-CB-CG	5.96	126.50	113.40
1	A	178	ARG	NE-CZ-NH1	5.73	123.17	120.30
1	A	220	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	A	205	TYR	CB-CG-CD1	-5.37	117.78	121.00
1	B	243	ARG	NE-CZ-NH2	5.36	122.98	120.30
1	B	200	LEU	CB-CG-CD1	-5.29	102.00	111.00
1	B	208	ARG	CB-CG-CD	5.25	125.26	111.60
1	A	158	ARG	CG-CD-NE	-5.24	100.79	111.80
1	B	240	LEU	CB-CG-CD1	-5.13	102.28	111.00
1	B	220	ARG	NE-CZ-NH1	5.09	122.85	120.30
1	A	177	VAL	CA-CB-CG2	-5.05	103.32	110.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	824	196	591	15	0
1	B	818	194	588	11	0
2	C	41	2	35	0	0
2	D	41	2	36	1	0
3	A	16	32	0	4	0
3	B	7	14	0	3	0
3	C	3	6	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	1	2	0	0	0
All	All	1751	448	1250	25	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (25) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:217:ILE:H	3:A:506:HOH:H2	1.12	0.96
1:B:217:ILE:H	3:B:510:HOH:H2	1.27	0.79
1:B:220:ARG:HD3	3:B:523:HOH:H1	1.62	0.63
1:B:242:HIS:HD2	1:B:243:ARG:O	1.84	0.61
1:A:243:ARG:NH2	1:A:245:THR:HG22	2.15	0.60
1:A:204:HIS:H	3:A:500:HOH:H2	1.48	0.60
1:B:188:CYS:SG	2:D:101:FTY:HE2	2.41	0.59
1:A:187:TYR:HB2	1:A:207:ILE:HB	1.83	0.59
1:B:220:ARG:NH1	1:B:238:ASP:OD2	2.36	0.59
1:A:146:ILE:HD11	1:A:151:TRP:CH2	2.38	0.58
1:A:146:ILE:HD12	1:A:149:GLU:HG3	1.86	0.58
1:B:174:THR:HA	1:B:246:THR:O	2.08	0.53
1:A:204:HIS:H	3:A:500:HOH:H1	1.55	0.53
1:A:163:ARG:HG2	1:A:163:ARG:HH11	1.73	0.53
1:A:242:HIS:HD2	1:A:243:ARG:O	1.94	0.51
1:B:201:ASN:HA	3:B:504:HOH:H2	1.79	0.47
1:B:175:PHE:CD1	1:B:244:LEU:HB3	2.50	0.46
1:B:158:ARG:O	1:B:162:GLU:HG3	2.16	0.46
1:A:159:ARG:HH11	1:B:249:PRO:HA	1.81	0.45
1:A:174:THR:HA	1:A:246:THR:O	2.17	0.44
1:A:219:SER:HA	3:A:525:HOH:H2	1.83	0.43
1:A:247:VAL:HG12	1:A:248:CYS:O	2.19	0.43
1:A:170:ASN:CG	1:A:174:THR:HG21	2.39	0.42
1:A:233:TYR:HA	1:A:236:HIS:O	2.19	0.42
1:B:151:TRP:HA	1:B:247:VAL:HG13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/107 (96%)	100 (97%)	3 (3%)	0	100	100
1	B	102/107 (95%)	97 (95%)	5 (5%)	0	100	100
2	C	1/4 (25%)	1 (100%)	0	0	100	100
2	D	1/4 (25%)	1 (100%)	0	0	100	100
All	All	207/222 (93%)	199 (96%)	8 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/94 (92%)	85 (99%)	1 (1%)	82	90
1	B	85/94 (90%)	85 (100%)	0	100	100
2	C	1/1 (100%)	1 (100%)	0	100	100
2	D	1/1 (100%)	1 (100%)	0	100	100
All	All	173/190 (91%)	172 (99%)	1 (1%)	92	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	171	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	242	HIS
1	B	242	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FTY	C	101	2	18,18,19	4.63	8 (44%)	25,27,29	3.09	12 (48%)
2	FTY	D	101	2	18,18,19	4.98	5 (27%)	25,27,29	3.35	10 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FTY	C	101	2	-	1/15/21/23	0/1/1/1
2	FTY	D	101	2	-	1/15/21/23	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	101	FTY	O-C	20.11	1.25	1.11
2	C	101	FTY	O-C	18.11	1.23	1.11
2	C	101	FTY	C1-CZ	3.75	1.55	1.50
2	C	101	FTY	P-O2P	-3.57	1.47	1.54
2	D	101	FTY	P-O2P	-3.27	1.48	1.54
2	D	101	FTY	C1-CZ	2.71	1.53	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	101	FTY	F1-C1	2.69	1.41	1.37
2	C	101	FTY	F2-C1	2.65	1.41	1.37
2	C	101	FTY	CA-C	2.23	1.52	1.48
2	C	101	FTY	P-O1P	-2.18	1.46	1.50
2	C	101	FTY	P-O3P	-2.18	1.50	1.54
2	C	101	FTY	F1-C1	2.12	1.40	1.37
2	D	101	FTY	F2-C1	2.09	1.40	1.37

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	101	FTY	C-CA-N	-9.83	104.02	113.83
2	C	101	FTY	F2-C1-F1	-7.21	97.47	106.69
2	D	101	FTY	O1P-P-C1	-6.91	101.70	112.41
2	C	101	FTY	O1P-P-C1	-6.52	102.30	112.41
2	D	101	FTY	O2P-P-C1	-6.11	99.27	106.07
2	C	101	FTY	P-C1-CZ	5.84	121.11	108.69
2	D	101	FTY	F2-C1-F1	-4.70	100.68	106.69
2	D	101	FTY	P-C1-CZ	4.41	118.08	108.69
2	C	101	FTY	CE1-CZ-C1	3.86	123.45	119.94
2	C	101	FTY	F2-C1-CZ	-3.61	104.75	110.75
2	C	101	FTY	O2P-P-C1	-3.54	102.13	106.07
2	C	101	FTY	CE2-CZ-C1	-3.35	116.89	119.94
2	C	101	FTY	O3P-P-O2P	3.22	117.09	107.93
2	D	101	FTY	O3P-P-O2P	3.15	116.91	107.93
2	D	101	FTY	F2-C1-P	3.12	113.51	107.79
2	D	101	FTY	O3P-P-C1	-2.95	102.78	106.07
2	C	101	FTY	CD1-CE1-CZ	-2.95	117.30	121.28
2	C	101	FTY	F2-C1-P	2.84	112.99	107.79
2	D	101	FTY	F1-C1-CZ	-2.70	106.27	110.75
2	D	101	FTY	F2-C1-CZ	-2.38	106.80	110.75
2	C	101	FTY	O3P-P-C1	-2.36	103.44	106.07
2	C	101	FTY	C-CA-N	-2.36	111.47	113.83

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	101	FTY	O1P-P-C1-CZ
2	D	101	FTY	O1P-P-C1-CZ

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	105/107 (98%)	-0.41	1 (0%) 79 80	7, 23, 46, 68	0
1	B	104/107 (97%)	-0.27	4 (3%) 38 39	9, 22, 57, 69	0
2	C	4/4 (100%)	0.30	0 100 100	14, 15, 17, 18	0
2	D	4/4 (100%)	-1.15	0 100 100	21, 23, 24, 28	0
All	All	217/222 (97%)	-0.35	5 (2%) 57 58	7, 22, 56, 69	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	194	PHE	3.8
1	B	197	ALA	3.1
1	B	183	THR	3.1
1	B	182	THR	2.1
1	B	195	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	FTY	C	101	18/19	0.12	0.84	0,13,20,23	0
2	FTY	D	101	18/19	0.09	-0.62	0,26,37,37	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.