



# Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 04:08 AM GMT

PDB ID : 1A09  
Title : C-src (SH2 domain) complexed with ace-formyl phosphotyrl-glu-(n,n-dipenty  
lamine)  
Authors : Shewchuk, L.; Jordan, S.  
Deposited on : 1997-12-10  
Resolution : 2.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	107	
1	B	107	
2	C	4	
2	D	4	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2733 atoms, of which 799 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C-SRC TYROSINE KINASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	106	Total	C	H	N	O	S	0	0	0
			1047	530	204	150	160	3			
1	B	102	Total	C	H	N	O	S	0	0	0
			1016	514	201	146	152	3			

- Molecule 2 is a protein called ACE-FORMYL PHOSPHOTYR-GLU-(N,N-DIPENTYLAMINE).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	4	Total	C	H	N	O	P	0	0	0
			44	27	3	3	10	1			
2	D	4	Total	C	H	N	O	P	0	0	0
			44	27	3	3	10	1			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	108	Total	H	O	0	0
			324	216	108		
3	B	77	Total	H	O	0	0
			231	154	77		
3	C	4	Total	H	O	0	0
			12	8	4		
3	D	5	Total	H	O	0	0
			15	10	5		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: C-SRC TYROSINE KINASE



- Molecule 1: C-SRC TYROSINE KINASE



- Molecule 2: ACE-FORMYL PHOSPHOTYR-GLU-(N,N-DIPENTYLAMINE)



- Molecule 2: ACE-FORMYL PHOSPHOTYR-GLU-(N,N-DIPENTYLAMINE)



There are no outlier residues recorded for this chain.

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.90Å 66.00Å 74.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.00 42.51 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.0 (6.00-2.00) 96.5 (42.51-2.00)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.49 (at 2.00Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.198 , (Not available) 0.208 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	25.8	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 47.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 17233 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2733	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DIP, ACE, PTH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.79	0/860	1.47	12/1160 (1.0%)
1	B	0.78	0/832	1.37	10/1121 (0.9%)
2	C	1.29	0/8	0.95	0/9
2	D	1.00	0/8	0.75	0/9
All	All	0.79	0/1708	1.42	22/2299 (1.0%)

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	158	ARG	NE-CZ-NH2	-14.98	112.81	120.30
1	A	158	ARG	NE-CZ-NH1	12.60	126.60	120.30
1	B	151	TRP	CD1-CG-CD2	9.58	113.96	106.30
1	B	151	TRP	CE2-CD2-CG	-8.23	100.72	107.30
1	A	243	ARG	NE-CZ-NH1	7.86	124.23	120.30
1	A	151	TRP	CE2-CD2-CG	-7.61	101.21	107.30
1	A	151	TRP	CD1-CG-CD2	7.47	112.28	106.30
1	A	151	TRP	CG-CD2-CE3	7.36	140.53	133.90
1	A	151	TRP	CB-CG-CD1	-6.79	118.17	127.00
1	B	172	ARG	NE-CZ-NH2	-6.44	117.08	120.30
1	B	151	TRP	CB-CG-CD1	-6.43	118.64	127.00
1	B	151	TRP	CG-CD1-NE1	-6.28	103.82	110.10
1	B	151	TRP	CG-CD2-CE3	6.05	139.35	133.90
1	A	188	CYS	N-CA-CB	-5.67	100.39	110.60
1	B	158	ARG	NE-CZ-NH1	5.67	123.13	120.30
1	B	152	TYR	CB-CG-CD1	-5.61	117.64	121.00
1	A	163	ARG	NE-CZ-NH1	5.60	123.10	120.30
1	A	216	TYR	CB-CG-CD1	-5.48	117.71	121.00
1	A	151	TRP	CG-CD1-NE1	-5.42	104.68	110.10
1	A	172	ARG	NE-CZ-NH1	5.41	123.01	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	158	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	B	172	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	843	204	606	25	0
1	B	815	201	591	16	0
2	C	41	3	37	1	0
2	D	41	3	37	0	0
3	A	108	216	0	18	0
3	B	77	154	0	13	0
3	C	4	8	0	2	0
3	D	5	10	0	0	0
All	All	1934	799	1271	41	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

All (41) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:217:ILE:H	3:A:518:HOH:H2	1.12	0.95
1:A:144:ASP:N	3:A:625:HOH:H1	1.67	0.91
1:B:155:LYS:HG3	3:B:690:HOH:H2	1.44	0.83
1:B:243:ARG:H	3:B:539:HOH:H1	1.29	0.78
1:B:217:ILE:H	3:B:537:HOH:H1	1.29	0.77
1:B:170:ASN:H	3:B:678:HOH:H2	1.36	0.74
1:A:243:ARG:H	3:A:528:HOH:H1	1.32	0.74
1:A:239:GLY:H	3:A:610:HOH:H2	1.37	0.71
1:B:236:HIS:HA	3:B:634:HOH:H1	1.60	0.66
1:B:238:ASP:H	3:B:594:HOH:H1	1.43	0.65

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:157:THR:H	3:A:543:HOH:H2	1.43	0.64
1:A:206:LYS:H	3:C:501:HOH:H1	1.46	0.64
1:B:227:GLN:HE22	3:B:648:HOH:H1	1.45	0.63
1:B:234:SER:O	1:B:243:ARG:HD3	2.01	0.61
1:B:155:LYS:NZ	3:B:670:HOH:H1	1.99	0.59
1:A:242:HIS:HD2	1:A:243:ARG:O	1.87	0.58
1:A:146:ILE:HG22	3:A:554:HOH:H1	1.70	0.57
1:A:174:THR:HA	1:A:246:THR:O	2.04	0.57
1:A:224:ASN:H	3:A:639:HOH:H1	1.53	0.54
1:A:172:ARG:NH2	1:A:194:PHE:HD2	2.06	0.54
1:A:209:LYS:NZ	3:A:660:HOH:H2	2.05	0.54
1:B:202:VAL:H	3:B:533:HOH:H1	1.55	0.53
1:B:155:LYS:HZ1	3:B:670:HOH:H1	1.55	0.52
1:B:214:GLY:H	3:B:622:HOH:H2	1.58	0.52
1:A:226:LEU:H	3:A:526:HOH:H1	1.58	0.51
1:B:160:GLU:HG3	3:B:611:HOH:H2	1.75	0.51
1:A:242:HIS:HB2	3:A:685:HOH:H1	1.75	0.50
1:A:206:LYS:HA	2:C:101:PTH:OF	2.13	0.48
1:A:144:ASP:N	3:A:625:HOH:O	2.41	0.47
1:A:238:ASP:H	3:A:642:HOH:H2	1.62	0.46
1:B:166:LEU:HD22	1:B:200:LEU:HD11	2.00	0.44
1:A:204:HIS:H	3:C:500:HOH:H1	1.64	0.43
1:A:206:LYS:HZ3	3:A:516:HOH:H1	1.66	0.43
1:B:212:SER:HB2	3:B:622:HOH:O	2.19	0.42
1:B:174:THR:HA	1:B:246:THR:O	2.20	0.42
1:A:211:ASP:H	3:A:520:HOH:H1	1.67	0.41
1:A:246:THR:HA	3:A:565:HOH:O	2.19	0.41
1:A:206:LYS:NZ	3:A:516:HOH:H1	2.19	0.41
1:A:163:ARG:HH12	3:A:601:HOH:H2	1.68	0.41
1:A:175:PHE:CD1	1:A:244:LEU:HB3	2.55	0.41
1:A:236:HIS:HA	3:A:629:HOH:H1	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	104/107 (97%)	100 (96%)	3 (3%)	1 (1%)	22	12
1	B	100/107 (94%)	98 (98%)	2 (2%)	0	100	100
2	C	1/4 (25%)	1 (100%)	0	0	100	100
2	D	1/4 (25%)	1 (100%)	0	0	100	100
All	All	206/222 (93%)	200 (97%)	5 (2%)	1 (0%)	38	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	198	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/94 (95%)	86 (97%)	3 (3%)	49	45
1	B	86/94 (92%)	84 (98%)	2 (2%)	63	63
2	C	1/1 (100%)	1 (100%)	0	100	100
2	D	1/1 (100%)	1 (100%)	0	100	100
All	All	177/190 (93%)	172 (97%)	5 (3%)	56	54

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	167	ASN
1	A	188	CYS
1	A	211	ASP
1	B	200	LEU
1	B	212	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	147	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	242	HIS
1	B	227	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PTH	C	101	1,2	18,18,19	5.11	3 (16%)	23,25,27	1.86	7 (30%)
2	PTH	D	101	1,2	18,18,19	5.29	5 (27%)	23,25,27	1.55	4 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTH	C	101	1,2	-	0/11/13/15	0/1/1/1
2	PTH	D	101	1,2	-	0/11/13/15	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	101	PTH	O-C	19.34	1.24	1.11
2	C	101	PTH	O-C	17.97	1.23	1.11
2	C	101	PTH	P-OH	11.40	1.77	1.60
2	D	101	PTH	P-OH	10.31	1.76	1.60
2	D	101	PTH	CA-C	2.46	1.52	1.48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	101	PTH	CA-C	2.16	1.52	1.48
2	D	101	PTH	P-O2P	-2.15	1.46	1.54
2	D	101	PTH	P-O3P	-2.09	1.47	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	101	PTH	C-CA-N	-4.59	109.24	113.83
2	D	101	PTH	C-CA-N	-3.27	110.56	113.83
2	D	101	PTH	O3P-P-O2P	3.22	120.16	107.61
2	C	101	PTH	OH-P-O1P	-3.16	93.44	108.29
2	C	101	PTH	P-OH-CZ	-3.13	114.54	122.86
2	C	101	PTH	CF-CE1-CD1	2.98	127.45	120.49
2	C	101	PTH	O3P-P-O2P	2.91	118.94	107.61
2	D	101	PTH	CB-CG-CD1	2.77	125.33	120.41
2	C	101	PTH	O2P-P-O1P	2.51	118.66	110.44
2	C	101	PTH	CF-CE1-CZ	-2.41	114.30	119.20
2	D	101	PTH	O3P-P-OH	-2.21	96.58	105.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	106/107 (99%)	0.55	7 (6%) 18 17	15, 25, 52, 70	0
1	B	102/107 (95%)	0.44	4 (3%) 37 37	15, 25, 45, 64	0
2	C	4/4 (100%)	-0.50	0 100 100	20, 22, 23, 24	0
2	D	4/4 (100%)	0.37	0 100 100	24, 28, 31, 33	0
All	All	216/222 (97%)	0.47	11 (5%) 27 26	15, 25, 51, 70	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	194	PHE	5.4
1	A	197	ALA	4.5
1	B	148	ALA	4.3
1	B	197	ALA	4.1
1	B	198	LYS	3.1
1	A	196	ASN	2.7
1	A	145	SER	2.5
1	B	196	ASN	2.5
1	A	144	ASP	2.4
1	A	213	GLY	2.2
1	A	172	ARG	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PTH	D	101	18/19	0.17	1.14	0,29,41,54	0
2	PTH	C	101	18/19	0.16	0.74	0,22,32,39	0

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

There are no ligands in this entry.

### 6.5 Other polymers

There are no such residues in this entry.