



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 04:07 AM GMT

PDB ID : 1A0Q
Title : 29G11 COMPLEXED WITH PHENYL [1-(1-N-SUCCINYLAMINO)PENTYL]PHOSPHONATE
Authors : Buchbinder, J.L.; Stephenson, R.C.; Scanlan, T.S.; Fletterick, R.J.
Deposited on : 1997-12-05
Resolution : 2.30 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	212	
2	H	217	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3301 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 29G11 FAB (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	211	Total	C	N	O	S	0	0	0
			1643	1032	278	328	5			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	4	LEU	MET	CONFLICT	GB 12002896
L	22	THR	ALA	CONFLICT	GB 12002896
L	26	SER	ASN	CONFLICT	GB 12002896
L	30	LYS	ASN	CONFLICT	GB 12002896
L	34	GLY	ALA	CONFLICT	GB 12002896
L	43	GLN	GLY	CONFLICT	GB 12002896
L	55	LEU	GLN	CONFLICT	GB 12002896
L	63	ARG	SER	CONFLICT	GB 12002896
L	92	TYR	ASP	CONFLICT	GB 12002896
L	141	SER	PRO	CONFLICT	GB 12002896
L	187	GLU	GLY	CONFLICT	GB 12002896

- Molecule 2 is a protein called 29G11 FAB (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	205	Total	C	N	O	S	0	0	0
			1540	977	249	308	6			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	8	ASP	GLY	CONFLICT	GB 3399661
H	9	ALA	PRO	CONFLICT	GB 3399661
H	20	ILE	MET	CONFLICT	GB 3399661
H	31	ASP	SER	CONFLICT	GB 3399661
H	32	HIS	TYR	CONFLICT	GB 3399661

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	34	ILE	MET	CONFLICT	GB 3399661
H	42	GLU	GLY	CONFLICT	GB 3399661
H	52	SER	ASN	CONFLICT	GB 3399661
H	53	GLY	TYR	CONFLICT	GB 3399661
H	55	GLY	ASP	CONFLICT	GB 3399661
H	56	ASP	GLY	CONFLICT	GB 3399661
H	57	ILE	THR	CONFLICT	GB 3399661
H	71	ALA	SER	CONFLICT	GB 3399661
H	81	GLN	GLU	CONFLICT	GB 3399661
H	82A	ASN	SER	CONFLICT	GB 3399661
H	91	LEU	TYR	CONFLICT	GB 3399661
H	93	LYS	VAL	CONFLICT	GB 3399661
H	?	-	GLY	DELETION	GB 3399661
H	97	TYR	ARG	CONFLICT	GB 3399661
H	98	GLY	PRO	CONFLICT	GB 3399661
H	99	ARG	TYR	CONFLICT	GB 3399661
H	100	SER	TYR	CONFLICT	GB 3399661
H	100A	ASN	ALA	CONFLICT	GB 3399661
H	100B	VAL	MET	CONFLICT	GB 3399661
H	108	THR	SER	CONFLICT	GB 3399661
H	109	LEU	VAL	CONFLICT	GB 3399661

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	H	2	Total Zn 2 2	0	0
3	L	1	Total Zn 1 1	0	0

- Molecule 4 is PHENYL[1-(N-SUCCINYLAMINO)PENTYL]PHOSPHONATE (three-letter code: HEP) (formula: C₁₅H₂₂NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	H	1	Total	C	N	O	P	0	0
			23	15	1	6	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	47	Total	O	0	0
			47	47		
5	L	45	Total	O	0	0
			45	45		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: 29G11 FAB (LIGHT CHAIN)



• Molecule 2: 29G11 FAB (HEAVY CHAIN)



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	36.78Å 82.61Å 132.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30	Depositor
% Data completeness (in resolution range)	93.0 (50.00-2.30)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	0.09	Depositor
Refinement program	X-PLOR 3.843	Depositor
R, R_{free}	0.203 , 0.275	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3301	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, HEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.67	0/1681	1.36	19/2277 (0.8%)
2	H	0.68	0/1577	1.35	14/2150 (0.7%)
All	All	0.67	0/3258	1.36	33/4427 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	1

There are no bond length outliers.

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	163	TRP	CD1-CG-CD2	8.17	112.84	106.30
1	L	140	TYR	CA-C-N	-8.10	99.39	117.20
2	H	47	TRP	CD1-CG-CD2	8.06	112.75	106.30
2	H	188	TRP	CD1-CG-CD2	8.03	112.72	106.30
2	H	103	TRP	CD1-CG-CD2	7.87	112.59	106.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	L	43	GLN	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1643	0	1590	18	0
2	H	1540	0	1503	12	0
3	H	2	0	0	0	0
3	L	1	0	0	0	0
4	H	23	0	20	1	0
5	H	47	0	0	0	0
5	L	45	0	0	0	0
All	All	3301	0	3113	30	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

The worst 5 of 30 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:L:149:LYS:HB2	1:L:193:THR:HB	1.77	0.66
2:H:35:HIS:HD2	2:H:47:TRP:HE1	1.49	0.59
1:L:43:GLN:HG3	1:L:45:ARG:HB2	1.85	0.58
2:H:4:LEU:HG	2:H:22:CYS:SG	2.48	0.53
1:L:145:ASN:HB3	1:L:197:THR:HB	1.92	0.51

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	L	209/212 (99%)	202 (97%)	6 (3%)	1 (0%)	38 45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	199/217 (92%)	192 (96%)	6 (3%)	1 (0%)	38	45
All	All	408/429 (95%)	394 (97%)	12 (3%)	2 (0%)	38	45

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	141	SER
2	H	40	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	185/189 (98%)	179 (97%)	6 (3%)	51	67
2	H	175/187 (94%)	169 (97%)	6 (3%)	49	64
All	All	360/376 (96%)	348 (97%)	12 (3%)	50	66

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	143	ASP
2	H	8	ASP
2	H	147	PRO
1	L	141	SER
2	H	85	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	5	GLN
2	H	82(A)	ASN
2	H	32	HIS
1	L	138	ASN
2	H	35	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	HEP	H	214	-	23,23,23	1.76	2 (8%)	30,30,30	1.34	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEP	H	214	-	-	0/24/24/24	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	214	HEP	P-C1	-5.81	1.74	1.83
4	H	214	HEP	P-O2P	-4.90	1.48	1.56

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	H	214	HEP	O2P-P-O1P	3.75	119.73	110.45
4	H	214	HEP	C13-C14-C12	2.68	116.56	112.65
4	H	214	HEP	C2-C1-N1	-2.60	109.64	112.02
4	H	214	HEP	O3P-P-O1P	-2.08	110.20	114.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.