



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 11:10 PM GMT

PDB ID : 4A1X
Title : Co-Complex structure of NS3-4A protease with the inhibitory peptide CP5-46-A (Synchrotron data)
Authors : Schmelz, S.; Kuegler, J.; Collins, J.; Heinz, D.
Deposited on : 2011-09-20
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	203	
1	B	203	
2	C	21	
2	D	21	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	CL	A	1182	-	X
4	CL	A	1188	-	X
4	CL	B	1184	-	X
4	CL	B	1187	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3397 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NONSTRUCTURAL PROTEIN 4A, SERINE PROTEASE NS3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	191	Total	C	N	O	S	0	10	0
			1461	908	263	277	13			
1	B	194	Total	C	N	O	S	0	10	0
			1479	917	269	281	12			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	LYS	-	LINKER	UNP P26662
A	0	GLY	-	LINKER	UNP P26662
A	1	GLY	-	LINKER	UNP P26662
A	181	GLY	-	EXPRESSION TAG	UNP P26662
A	182	SER	-	EXPRESSION TAG	UNP P26662
A	183	HIS	-	EXPRESSION TAG	UNP P26662
A	184	HIS	-	EXPRESSION TAG	UNP P26662
A	185	HIS	-	EXPRESSION TAG	UNP P26662
A	186	HIS	-	EXPRESSION TAG	UNP P26662
A	187	HIS	-	EXPRESSION TAG	UNP P26662
A	188	HIS	-	EXPRESSION TAG	UNP P26662
A	26	ARG	LYS	SEE REMARK 999	UNP P26662
A	30	GLU	ASP	SEE REMARK 999	UNP P26662
A	36	VAL	LEU	SEE REMARK 999	UNP P26662
A	86	GLN	PRO	SEE REMARK 999	UNP P26662
A	94	LEU	MET	SEE REMARK 999	UNP P26662
A	114	ILE	VAL	VARIANT	UNP P26662
A	132	VAL	ILE	VARIANT	UNP P26662
A	150	ALA	VAL	SEE REMARK 999	UNP P26662
A	170	VAL	ILE	SEE REMARK 999	UNP P26662
B	-1	LYS	-	LINKER	UNP P26662
B	0	GLY	-	LINKER	UNP P26662
B	1	GLY	-	LINKER	UNP P26662
B	181	GLY	-	EXPRESSION TAG	UNP P26662

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	182	SER	-	EXPRESSION TAG	UNP P26662
B	183	HIS	-	EXPRESSION TAG	UNP P26662
B	184	HIS	-	EXPRESSION TAG	UNP P26662
B	185	HIS	-	EXPRESSION TAG	UNP P26662
B	186	HIS	-	EXPRESSION TAG	UNP P26662
B	187	HIS	-	EXPRESSION TAG	UNP P26662
B	188	HIS	-	EXPRESSION TAG	UNP P26662
B	26	ARG	LYS	SEE REMARK 999	UNP P26662
B	30	GLU	ASP	SEE REMARK 999	UNP P26662
B	36	VAL	LEU	SEE REMARK 999	UNP P26662
B	86	GLN	PRO	SEE REMARK 999	UNP P26662
B	94	LEU	MET	SEE REMARK 999	UNP P26662
B	114	ILE	VAL	SEE REMARK 999	UNP P26662
B	132	VAL	ILE	VARIANT	UNP P26662
B	150	ALA	VAL	SEE REMARK 999	UNP P26662
B	170	VAL	ILE	SEE REMARK 999	UNP P26662

- Molecule 2 is a protein called CP5-46-A PEPTIDE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	17	Total	C	N	O	S	0	0	0
			132	86	22	23	1			
2	D	15	Total	C	N	O	S	0	0	0
			117	78	17	21	1			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	6	Total	Cl	0	0
			6	6		
4	A	7	Total	Cl	0	0
			7	7		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	87	Total 87	O 87	0	0
5	B	95	Total 95	O 95	0	0
5	C	8	Total 8	O 8	0	0
5	D	3	Total 3	O 3	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	93.32Å 93.32Å 82.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	80.82 – 1.90 19.96 – 1.90	Depositor EDS
% Data completeness (in resolution range)	80.8 (80.82-1.90) 99.7 (19.96-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.182 , 0.182 0.182 , 0.216	Depositor DCC
R_{free} test set	1619 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	21.8	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 42.1	EDS
Estimated twinning fraction	0.064 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 32044 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3397	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1487	0.66	0/2023
1	B	0.51	1/1505 (0.1%)	0.67	0/2046
2	C	0.58	0/136	0.62	0/185
2	D	0.56	0/121	0.63	0/166
All	All	0.51	1/3249 (0.0%)	0.66	0/4420

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	85	TRP	CD2-CE2	5.13	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1461	0	1466	26	0
1	B	1479	0	1490	20	0
2	C	132	0	126	7	0
2	D	117	0	110	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	7	0	0	3	0
4	B	6	0	0	4	0
5	A	87	0	0	2	0
5	B	95	0	0	2	0
5	C	8	0	0	0	0
5	D	3	0	0	0	0
All	All	3397	0	3192	46	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (46) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:89[B]:PRO:O	5:B:2040:HOH:O	1.66	1.12
1:A:89[B]:PRO:O	5:A:2035:HOH:O	2.04	0.74
1:B:154:PHE:CZ	2:D:9:LEU:HD13	2.22	0.74
1:B:154:PHE:CZ	2:D:9:LEU:CD1	2.71	0.73
1:A:132:VAL:CG1	1:A:159[A]:CYS:SG	2.82	0.67
1:A:132:VAL:HG11	1:A:159[A]:CYS:SG	2.37	0.65
1:A:132:VAL:HG22	4:A:1188:CL:CL	2.35	0.64
1:B:132:VAL:HG11	1:B:159:CYS:SG	2.38	0.64
1:B:154:PHE:CE2	2:D:9:LEU:HD13	2.36	0.61
1:A:25:ASP:OD2	5:A:2007:HOH:O	2.16	0.60
1:A:132:VAL:HG12	1:A:159[A]:CYS:SG	2.43	0.58
1:A:154:PHE:CZ	2:C:9:LEU:CD1	2.87	0.58
1:B:-4:LEU:O	1:B:-3:SER:HB2	2.04	0.57
1:B:47[B]:CYS:HG	1:B:93:SER:HG	1.50	0.56
1:A:130:ARG:HD2	1:A:134:TYR:CE2	2.41	0.56
1:B:139[B]:SER:OG	5:B:2053:HOH:O	2.17	0.55
1:A:157:ALA:HB2	2:C:9:LEU:HD11	1.89	0.53
1:A:159[B]:CYS:SG	2:C:7:VAL:CG2	2.98	0.52
1:B:132:VAL:HG22	4:B:1184:CL:CL	2.46	0.52
1:A:104:LEU:HD22	1:A:151[A]:VAL:HG21	1.92	0.51
1:A:104:LEU:HD22	1:A:151[B]:VAL:HG11	1.92	0.51
1:B:154:PHE:HZ	2:D:9:LEU:CD1	2.24	0.49
1:B:-14:GLY:HA2	4:B:1186:CL:CL	2.49	0.49
1:B:104:LEU:HD22	1:B:151[A]:VAL:HG21	1.95	0.49
1:A:28:GLN:NE2	1:A:30:GLU:OE2	2.46	0.48
1:B:13:LEU:O	1:B:17:ILE:HG12	2.12	0.48
1:A:159[B]:CYS:SG	2:C:7:VAL:HG21	2.54	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:154:PHE:CZ	2:C:9:LEU:HD12	2.49	0.47
1:A:40:THR:OG1	4:A:1182:CL:CL	2.69	0.47
1:B:143:LEU:HB3	1:B:151[B]:VAL:HG12	1.97	0.47
1:B:132:VAL:CG1	1:B:159:CYS:SG	3.04	0.46
1:A:154:PHE:HZ	2:C:9:LEU:CD1	2.30	0.45
1:A:-3:SER:HB3	1:A:4:THR:OG1	2.16	0.45
1:A:75:TYR:CZ	1:A:179:MET:HG2	2.52	0.44
1:A:151[B]:VAL:HG22	1:A:151[B]:VAL:O	2.18	0.43
1:A:130:ARG:HD2	1:A:134:TYR:CD2	2.53	0.43
1:A:-6:ILE:HB	1:A:33:VAL:HB	2.00	0.43
1:B:158:VAL:HB	1:B:166:ALA:HB3	2.00	0.43
1:A:116:VAL:HG22	1:A:126:LEU:HD23	2.01	0.42
1:A:159[B]:CYS:SG	2:C:7:VAL:HG23	2.59	0.42
1:B:15:GLY:HA3	4:B:1188:CL:CL	2.57	0.41
1:A:15:GLY:HA3	4:A:1183:CL:CL	2.58	0.41
1:B:-4:LEU:HD23	1:B:5:ALA:HB2	2.02	0.41
1:B:38:THR:HB	4:B:1185:CL:CL	2.57	0.40
1:B:-4:LEU:O	1:B:-3:SER:CB	2.70	0.40
1:A:47[B]:CYS:SG	1:A:93:SER:OG	2.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/203 (97%)	195 (99%)	2 (1%)	0	100	100
1	B	200/203 (98%)	197 (98%)	3 (2%)	0	100	100
2	C	15/21 (71%)	14 (93%)	1 (7%)	0	100	100
2	D	13/21 (62%)	12 (92%)	1 (8%)	0	100	100
All	All	425/448 (95%)	418 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	164/164 (100%)	163 (99%)	1 (1%)	92	92
1	B	165/164 (101%)	164 (99%)	1 (1%)	92	92
2	C	14/17 (82%)	14 (100%)	0	100	100
2	D	13/17 (76%)	13 (100%)	0	100	100
All	All	356/362 (98%)	354 (99%)	2 (1%)	92	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-7	ARG
1	B	-7	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 15 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/203 (94%)	0.23	6 (3%) 47 48	11, 24, 42, 60	0
1	B	194/203 (95%)	0.33	10 (5%) 26 26	12, 26, 49, 67	0
2	C	17/21 (80%)	1.15	5 (29%) 1 1	16, 32, 66, 72	0
2	D	15/21 (71%)	0.85	3 (20%) 2 1	17, 30, 64, 76	0
All	All	417/448 (93%)	0.34	24 (5%) 22 22	11, 26, 49, 76	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	180	ARG	4.2
1	B	0	GLY	3.9
2	C	19	HIS	3.8
2	D	20	CYS	3.8
1	B	99	CYS	3.7
1	B	1	GLY	3.7
1	B	181	GLY	3.7
2	C	18	ILE	3.6
1	A	79	ASP	3.5
2	C	20	CYS	3.3
1	B	100	GLY	3.2
1	A	177	THR	3.2
1	B	121	ASP	3.2
1	B	98	THR	3.0
1	A	99	CYS	2.7
1	A	100	GLY	2.6
1	A	2	PRO	2.5
2	D	19	HIS	2.5
2	C	5	ARG	2.5
1	A	180	ARG	2.4
2	C	8	TYR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	8	TYR	2.3
1	B	177	THR	2.3
1	B	-3	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	CL	A	1188	1/1	0.30	6.19	57,57,57,57	0
4	CL	B	1184	1/1	0.21	4.08	52,52,52,52	0
4	CL	B	1187	1/1	0.26	3.56	55,55,55,55	0
4	CL	A	1182	1/1	0.19	2.66	55,55,55,55	0
4	CL	B	1185	1/1	0.17	1.32	51,51,51,51	0
4	CL	A	1187	1/1	0.11	-0.34	45,45,45,45	0
3	ZN	B	1182	1/1	0.13	-0.66	62,62,62,62	0
4	CL	A	1186	1/1	0.06	-1.43	32,32,32,32	0
4	CL	B	1183	1/1	0.04	-1.55	33,33,33,33	0
3	ZN	A	1181	1/1	0.04	-1.56	42,42,42,42	0
4	CL	A	1183	1/1	0.04	-2.14	24,24,24,24	0
4	CL	B	1188	1/1	0.04	-3.05	26,26,26,26	0
4	CL	A	1184	1/1	0.06	-3.18	31,31,31,31	0
4	CL	B	1186	1/1	0.05	-4.14	33,33,33,33	0
4	CL	A	1185	1/1	0.08	-6.14	52,52,52,52	0

6.5 Other polymers ⓘ

There are no such residues in this entry.