



# Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 01:19 AM GMT

PDB ID : 2A2A  
Title : High-resolution crystallographic analysis of the autoinhibited conformation of a human death-associated protein kinase  
Authors : Kursula, P.; Wilmanns, M.  
Deposited on : 2005-06-22  
Resolution : 1.47 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2222 (1.50-1.46)
Clashscore	79885	2555 (1.50-1.46)
Ramachandran outliers	78287	2496 (1.50-1.46)
Sidechain outliers	78261	2494 (1.50-1.46)
RSRZ outliers	66119	2223 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	321	
1	B	321	
1	C	321	
1	D	321	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	NA	A	3178	-	X
2	NA	C	3158	-	X
2	NA	C	3291	-	X
4	DTT	A	2002	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Geometry	Electron density
4	DTT	B	2003	-	X
4	DTT	C	2004	-	X
4	DTT	D	2005	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11556 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Death-associated protein kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	304	Total	C	N	O	S	0	10	0
			2557	1628	436	488	5			
1	B	303	Total	C	N	O	S	0	7	0
			2523	1609	434	476	4			
1	C	303	Total	C	N	O	S	0	8	0
			2532	1619	431	478	4			
1	D	303	Total	C	N	O	S	0	5	0
			2512	1605	433	470	4			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	CLONING ARTIFACT	UNP Q9UIK4
A	231	SER	ALA	SEE REMARK 999	UNP Q9UIK4
A	243	HIS	GLN	SEE REMARK 999	UNP Q9UIK4
B	0	GLY	-	CLONING ARTIFACT	UNP Q9UIK4
B	231	SER	ALA	SEE REMARK 999	UNP Q9UIK4
B	243	HIS	GLN	SEE REMARK 999	UNP Q9UIK4
C	0	GLY	-	CLONING ARTIFACT	UNP Q9UIK4
C	231	SER	ALA	SEE REMARK 999	UNP Q9UIK4
C	243	HIS	GLN	SEE REMARK 999	UNP Q9UIK4
D	0	GLY	-	CLONING ARTIFACT	UNP Q9UIK4
D	231	SER	ALA	SEE REMARK 999	UNP Q9UIK4
D	243	HIS	GLN	SEE REMARK 999	UNP Q9UIK4

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Na	0	0
			2	2		
2	A	2	Total	Na	0	0
			2	2		

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total	Na	0	0
			1	1		
2	C	3	Total	Na	0	0
			3	3		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		
3	C	1	Total	Cl	0	0
			1	1		

- Molecule 4 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (three-letter code: DTT) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>2</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			8	4	2	2		
4	B	1	Total	C	O	S	0	0
			8	4	2	2		
4	C	1	Total	C	O	S	0	0
			8	4	2	2		
4	D	1	Total	C	O	S	0	0
			8	4	2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		
5	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	361	Total	O	0	0
			361	361		
6	B	315	Total	O	0	0
			315	315		
6	C	346	Total	O	0	0
			346	346		
6	D	349	Total	O	0	0
			349	349		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Death-associated protein kinase 2




- Molecule 1: Death-associated protein kinase 2



- Molecule 1: Death-associated protein kinase 2



- Molecule 1: Death-associated protein kinase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.30Å 60.65Å 98.71Å 92.16° 103.45° 94.25°	Depositor
Resolution (Å)	20.00 – 1.47 38.62 – 1.47	Depositor EDS
% Data completeness (in resolution range)	94.8 (20.00-1.47) 94.8 (38.62-1.47)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 1.47Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.150 , 0.207 0.165 , 0.218	Depositor DCC
$R_{free}$ test set	2004 reflections (1.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.4	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 55.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 200381 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	11556	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, NA, DTT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.82	1/2605 (0.0%)	0.82	1/3513 (0.0%)
1	B	0.77	0/2571	0.82	2/3466 (0.1%)
1	C	0.81	0/2580	0.86	2/3480 (0.1%)
1	D	0.82	0/2560	0.83	0/3452
All	All	0.81	1/10316 (0.0%)	0.83	5/13911 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	208	TYR	CD1-CE1	5.22	1.47	1.39

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	287	ARG	NE-CZ-NH1	5.75	123.18	120.30
1	C	170	ASP	CB-CG-OD2	5.58	123.32	118.30
1	B	237	ASP	CB-CG-OD1	5.32	123.09	118.30
1	C	63	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	B	237	ASP	CB-CG-OD2	-5.25	113.57	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the

chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2557	0	0	12	0
1	B	2523	0	0	7	0
1	C	2532	0	0	11	0
1	D	2512	0	0	8	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	3	0	0	0	0
2	D	1	0	0	0	0
3	A	2	0	0	1	0
3	C	1	0	0	0	0
4	A	8	0	0	2	0
4	B	8	0	0	0	0
4	C	8	0	0	0	0
4	D	8	0	0	0	0
5	A	6	0	0	0	0
5	C	6	0	0	0	0
5	D	6	0	0	0	0
6	A	361	0	0	9	0
6	B	315	0	0	4	0
6	C	346	0	0	5	0
6	D	349	0	0	3	0
All	All	11556	0	0	40	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (40) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:A:2002:DTT:S1	4:A:2002:DTT:S4	2.29	1.30
1:A:118[B]:SER:OG	6:A:3437:HOH:O	1.53	1.24
1:A:63[B]:ARG:NH1	6:A:3392:HOH:O	1.68	1.21
1:A:63[B]:ARG:CZ	6:A:3392:HOH:O	2.12	0.85
1:A:23:GLN:N	6:A:3645:HOH:O	2.08	0.85
1:A:261:ARG:NE	6:A:3371:HOH:O	2.24	0.69
1:D:59:GLU:OE2	6:D:3512:HOH:O	2.15	0.65
1:B:110[A]:SER:OG	6:B:3920:HOH:O	2.16	0.62
1:D:250:ASP:OD2	6:D:3476:HOH:O	2.16	0.61
1:C:125:ASP:OD2	6:C:3605:HOH:O	2.17	0.59
1:A:182:GLU:OE1	1:A:303:ARG:NH2	2.37	0.57
1:B:118[B]:SER:O	6:B:3930:HOH:O	2.18	0.56

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:58:ARG:NE	1:C:62:GLU:OE1	2.40	0.55
1:B:63[A]:ARG:NH1	6:B:4051:HOH:O	2.39	0.54
1:A:108:LYS:CE	6:A:3627:HOH:O	2.56	0.53
1:C:106:ALA:O	1:C:297:ARG:NE	2.43	0.52
1:B:103:ASP:OD2	1:B:300:TYR:CE1	2.66	0.49
1:C:49:SER:N	1:C:52:SER:OG	2.47	0.48
1:A:50:ARG:N	6:A:3620:HOH:O	2.48	0.47
1:D:292:ASN:CB	1:D:295:ASN:ND2	2.77	0.47
4:A:2002:DTT:S1	4:A:2002:DTT:C4	3.01	0.47
1:C:25:ALA:N	6:C:3610:HOH:O	2.47	0.47
1:D:134:LYS:NZ	1:D:170:ASP:OD2	2.49	0.46
1:A:134:LYS:NZ	6:A:3640:HOH:O	2.48	0.46
1:C:246[A]:GLU:OE2	1:C:249:LYS:NZ	2.50	0.45
1:A:146:MET:CE	6:A:3495:HOH:O	2.64	0.45
1:A:167:GLU:N	3:A:3065:CL:CL	2.87	0.44
1:B:49:SER:N	1:B:52:SER:OG	2.50	0.44
1:D:249:LYS:NZ	6:D:3275:HOH:O	2.50	0.44
1:C:78[A]:THR:CG2	6:C:3307:HOH:O	2.66	0.43
1:C:24:PHE:C	6:C:3610:HOH:O	2.55	0.43
1:B:109:GLU:O	6:B:4041:HOH:O	2.21	0.43
1:D:292:ASN:CG	1:D:295:ASN:ND2	2.73	0.42
1:B:49:SER:O	1:B:52:SER:N	2.52	0.42
1:A:46:LYS:NZ	1:A:84:GLU:OE2	2.53	0.42
1:C:109:GLU:CD	1:C:109:GLU:N	2.73	0.41
1:D:295:ASN:N	1:D:295:ASN:ND2	2.69	0.41
1:C:21:SER:O	1:C:22:GLY:O	2.38	0.41
1:C:258:LYS:NZ	6:C:3446:HOH:O	2.54	0.40
1:D:71[A]:VAL:CG1	1:D:76:VAL:CG1	2.99	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	312/321 (97%)	303 (97%)	9 (3%)	0	100 100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	308/321 (96%)	298 (97%)	9 (3%)	1 (0%)	50 19
1	C	309/321 (96%)	295 (96%)	12 (4%)	2 (1%)	33 9
1	D	306/321 (95%)	298 (97%)	7 (2%)	1 (0%)	50 19
All	All	1235/1284 (96%)	1194 (97%)	37 (3%)	4 (0%)	50 19

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	23	GLN
1	C	22	GLY
1	C	50	ARG
1	D	21	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	283/290 (98%)	276 (98%)	7 (2%)	60 23
1	B	279/290 (96%)	273 (98%)	6 (2%)	64 28
1	C	280/290 (97%)	273 (98%)	7 (2%)	60 23
1	D	277/290 (96%)	273 (99%)	4 (1%)	78 50
All	All	1119/1160 (96%)	1095 (98%)	24 (2%)	66 30

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	LEU
1	A	50	ARG
1	A	86	ARG
1	A	261	ARG
1	A	293	LEU
1	A	294[A]	GLU
1	A	294[B]	GLU
1	B	37	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	54	ARG
1	B	86	ARG
1	B	169	GLU
1	B	299	GLN
1	B	300	TYR
1	C	2	GLU
1	C	19	LEU
1	C	49	SER
1	C	50	ARG
1	C	109	GLU
1	C	267	GLN
1	C	293	LEU
1	D	19	LEU
1	D	294	GLU
1	D	295	ASN
1	D	300	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 18 ligands modelled in this entry, 11 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	DTT	A	2002	-	7,7,7	0.95	0	8,8,8	2.11	4 (50%)
5	GOL	A	3001	-	5,5,5	0.20	0	5,5,5	0.57	0
4	DTT	B	2003	-	7,7,7	0.78	0	8,8,8	1.84	2 (25%)
4	DTT	C	2004	-	7,7,7	0.79	0	8,8,8	2.08	4 (50%)
5	GOL	C	3002	-	5,5,5	0.44	0	5,5,5	0.88	0
4	DTT	D	2005	-	7,7,7	0.68	0	8,8,8	1.49	1 (12%)
5	GOL	D	3003	-	5,5,5	0.26	0	5,5,5	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DTT	A	2002	-	1/1/2/2	0/8/8/8	0/0/0/0
5	GOL	A	3001	-	-	0/4/4/4	0/0/0/0
4	DTT	B	2003	-	1/1/2/2	0/8/8/8	0/0/0/0
4	DTT	C	2004	-	1/1/2/2	0/8/8/8	0/0/0/0
5	GOL	C	3002	-	-	0/4/4/4	0/0/0/0
4	DTT	D	2005	-	1/1/2/2	0/8/8/8	0/0/0/0
5	GOL	D	3003	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	2003	DTT	C2-C1-S1	-4.12	108.17	114.66
4	A	2002	DTT	O3-C3-C4	-3.65	103.01	109.87
4	C	2004	DTT	O2-C2-C1	-3.21	103.84	109.87
4	C	2004	DTT	C3-C4-S4	-2.90	110.09	114.66
4	D	2005	DTT	C2-C1-S1	-2.84	110.18	114.66
4	A	2002	DTT	O2-C2-C1	-2.84	104.54	109.87
4	A	2002	DTT	C2-C1-S1	-2.56	110.62	114.66
4	A	2002	DTT	C4-C3-C2	2.21	118.11	113.07
4	C	2004	DTT	O2-C2-C3	2.19	114.11	109.83
4	C	2004	DTT	O3-C3-C2	2.04	113.82	109.83
4	B	2003	DTT	O3-C3-C2	-2.04	105.86	109.83

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	2003	DTT	C3
4	A	2002	DTT	C3
4	D	2005	DTT	C3
4	C	2004	DTT	C3

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	304/321 (94%)	0.06	16 (5%) 25 29	15, 23, 52, 76	0
1	B	303/321 (94%)	0.15	17 (5%) 24 26	14, 23, 54, 83	0
1	C	303/321 (94%)	0.06	14 (4%) 31 35	15, 23, 48, 73	0
1	D	303/321 (94%)	0.12	17 (5%) 24 26	15, 23, 47, 93	0
All	All	1213/1284 (94%)	0.10	64 (5%) 26 29	14, 23, 52, 93	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	300	TYR	9.0
1	B	304	ARG	8.1
1	D	301	VAL	6.9
1	B	48	GLN	6.8
1	C	301	VAL	6.8
1	D	304	ARG	6.7
1	A	49	SER	6.6
1	A	51	ALA	6.5
1	B	302	ARG	6.3
1	B	24	PHE	6.1
1	A	302	ARG	6.1
1	A	53	ARG	6.0
1	A	1	MET	5.6
1	D	302	ARG	5.6
1	A	54	ARG	5.4
1	B	51	ALA	5.3
1	A	48	GLN	5.3
1	D	303	ARG	5.2
1	D	21	SER	5.0
1	B	53	ARG	5.0
1	C	53	ARG	5.0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	24	PHE	5.0
1	C	48	GLN	4.9
1	C	49	SER	4.6
1	B	52	SER	4.6
1	B	300	TYR	4.6
1	D	48	GLN	4.5
1	D	49	SER	4.5
1	B	303	ARG	4.3
1	B	49	SER	4.3
1	D	51	ALA	4.2
1	A	304	ARG	4.1
1	C	302	ARG	4.0
1	C	54	ARG	3.9
1	B	23	GLN	3.7
1	A	24	PHE	3.7
1	C	24	PHE	3.5
1	C	22	GLY	3.5
1	D	278	VAL	3.5
1	B	54	ARG	3.4
1	C	300	TYR	3.3
1	C	304	ARG	3.2
1	B	22	GLY	3.0
1	A	23	GLN	2.9
1	D	86	ARG	2.9
1	A	303	ARG	2.8
1	A	301	VAL	2.8
1	A	52	SER	2.7
1	D	171	GLY	2.6
1	A	300	TYR	2.6
1	B	170	ASP	2.6
1	C	50	ARG	2.5
1	B	278	VAL	2.5
1	B	21	SER	2.5
1	D	52	SER	2.5
1	A	22	GLY	2.4
1	C	303	ARG	2.3
1	C	51	ALA	2.3
1	D	170	ASP	2.3
1	D	23	GLN	2.2
1	B	86	ARG	2.2
1	A	86	ARG	2.1
1	D	261	ARG	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	145[A]	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NA	A	3178	1/1	0.17	12.82	43,43,43,43	0
2	NA	C	3158	1/1	0.13	5.85	31,31,31,31	0
4	DTT	C	2004	8/8	0.10	3.77	39,42,42,46	0
4	DTT	A	2002	8/8	0.12	3.76	39,43,49,61	0
4	DTT	B	2003	8/8	0.13	2.90	35,41,48,50	0
2	NA	C	3291	1/1	0.17	2.34	46,46,46,46	0
4	DTT	D	2005	8/8	0.11	2.16	42,47,51,54	0
2	NA	D	3175	1/1	0.10	1.20	46,46,46,46	0
5	GOL	D	3003	6/6	0.09	0.76	25,32,38,44	0
2	NA	A	3152	1/1	0.07	0.46	32,32,32,32	0
2	NA	C	3281	1/1	0.07	0.13	66,66,66,66	0
2	NA	B	3221	1/1	0.09	-0.42	40,40,40,40	0
2	NA	B	3763	1/1	0.09	-0.56	52,52,52,52	0
5	GOL	C	3002	6/6	0.08	-0.68	27,36,38,41	0
3	CL	A	3318	1/1	0.05	-0.96	53,53,53,53	0
5	GOL	A	3001	6/6	0.07	-1.07	26,33,36,37	0
3	CL	C	3039	1/1	0.04	-1.74	38,38,38,38	0
3	CL	A	3065	1/1	0.03	-3.29	37,37,37,37	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.