



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 09:05 PM GMT

PDB ID : 3A25
Title : Crystal structure of P. horikoshii TYW2 in complex with AdoMet
Authors : Umitsu, M.; Nishimasu, H.; Ishitani, R.; Nureki, O.
Deposited on : 2009-04-28
Resolution : 2.30 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	301	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2337 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein PH0793.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	264	2163	1409	371	376	7	0	1	0

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MET	-	EXPRESSION TAG	UNP O58523
A	-21	GLY	-	EXPRESSION TAG	UNP O58523
A	-20	SER	-	EXPRESSION TAG	UNP O58523
A	-19	SER	-	EXPRESSION TAG	UNP O58523
A	-18	HIS	-	EXPRESSION TAG	UNP O58523
A	-17	HIS	-	EXPRESSION TAG	UNP O58523
A	-16	HIS	-	EXPRESSION TAG	UNP O58523
A	-15	HIS	-	EXPRESSION TAG	UNP O58523
A	-14	HIS	-	EXPRESSION TAG	UNP O58523
A	-13	HIS	-	EXPRESSION TAG	UNP O58523
A	-12	SER	-	EXPRESSION TAG	UNP O58523
A	-11	SER	-	EXPRESSION TAG	UNP O58523
A	-10	GLY	-	EXPRESSION TAG	UNP O58523
A	-9	LEU	-	EXPRESSION TAG	UNP O58523
A	-8	GLU	-	EXPRESSION TAG	UNP O58523
A	-7	VAL	-	EXPRESSION TAG	UNP O58523
A	-6	LEU	-	EXPRESSION TAG	UNP O58523
A	-5	PHE	-	EXPRESSION TAG	UNP O58523
A	-4	GLN	-	EXPRESSION TAG	UNP O58523
A	-3	GLY	-	EXPRESSION TAG	UNP O58523
A	-2	PRO	-	EXPRESSION TAG	UNP O58523
A	-1	LEU	-	EXPRESSION TAG	UNP O58523
A	0	HIS	-	EXPRESSION TAG	UNP O58523

- Molecule 2 is S-ADENOSYLMETHIONINE (three-letter code: SAM) (formula: C₁₅H₂₂N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	27	15	6	5	1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	147	Total	O	0	0
			147	147		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein PH0793

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	131.60Å 131.60Å 99.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.54 – 2.30 45.54 – 2.30	Depositor EDS
% Data completeness (in resolution range)	94.4 (45.54-2.30) 94.4 (45.54-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.29Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.205 , 0.240 0.203 , 0.238	Depositor DCC
R_{free} test set	1344 reflections (4.99%)	DCC
Wilson B-factor (Å ²)	51.5	Xtriage
Anisotropy	0.260	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 65.9	EDS
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 26936 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2337	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2211	0.63	2/2986 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	7[A]	LYS	N-CA-C	5.03	124.58	111.00
1	A	7[B]	LYS	N-CA-C	5.03	124.58	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2163	0	2259	99	0
2	A	27	0	22	1	0
3	A	147	0	0	9	0
All	All	2337	0	2281	99	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 22.

All (99) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:263:ALA:HB1	1:A:264:PRO:HA	1.28	1.15
1:A:6:ILE:N	1:A:8:PRO:HD2	1.68	1.08
1:A:258:LYS:HB3	1:A:267:TRP:CZ3	1.91	1.05
1:A:259:ILE:HG22	1:A:260:LYS:H	1.31	0.94
1:A:201:VAL:HB	1:A:226:THR:HG23	1.50	0.89
1:A:261:ARG:HG3	1:A:267:TRP:CE2	2.08	0.89
1:A:276:PHE:HB2	3:A:296:HOH:O	1.73	0.87
1:A:255:ASN:HD22	1:A:256:GLU:N	1.74	0.83
1:A:263:ALA:HB1	1:A:264:PRO:CA	2.09	0.80
1:A:259:ILE:HG22	1:A:260:LYS:N	1.96	0.79
1:A:6:ILE:O	1:A:6:ILE:HD12	1.83	0.79
1:A:6:ILE:N	1:A:8:PRO:CD	2.46	0.78
1:A:17:GLU:OE2	1:A:51:ARG:HD3	1.84	0.77
1:A:113:VAL:O	1:A:117:VAL:HG23	1.86	0.76
1:A:9:ARG:HG2	1:A:9:ARG:HH11	1.51	0.74
1:A:6:ILE:HG22	1:A:9:ARG:HB2	1.70	0.73
1:A:7[A]:LYS:H	1:A:7[A]:LYS:HE3	1.56	0.71
1:A:258:LYS:HB3	1:A:267:TRP:CE3	2.26	0.70
1:A:183:ASP:OD1	1:A:185:ARG:HD2	1.92	0.69
1:A:9:ARG:NH1	1:A:9:ARG:HG2	2.07	0.68
1:A:135:GLY:HA3	2:A:279:SAM:HE1	1.76	0.68
1:A:261:ARG:HB2	1:A:267:TRP:CZ3	2.29	0.67
1:A:118:ARG:HH11	1:A:257:LEU:HD12	1.58	0.67
1:A:25:LEU:CD1	1:A:44:GLU:HG3	2.25	0.67
1:A:263:ALA:CB	1:A:264:PRO:HA	2.18	0.65
1:A:274:ARG:HD2	1:A:276:PHE:CZ	2.33	0.64
1:A:217:ASP:OD1	1:A:277:LYS:N	2.31	0.64
1:A:118:ARG:HH11	1:A:257:LEU:CD1	2.12	0.63
1:A:235:GLU:HB3	1:A:236:PRO:HA	1.80	0.63
1:A:261:ARG:HG3	1:A:267:TRP:CZ2	2.33	0.63
1:A:245:LYS:HA	1:A:245:LYS:HE3	1.79	0.62
1:A:212:LEU:O	1:A:277:LYS:HE2	2.00	0.62
1:A:25:LEU:HD13	1:A:44:GLU:HG3	1.81	0.61
1:A:230:LYS:HG3	3:A:337:HOH:O	2.03	0.59
1:A:261:ARG:HB2	1:A:267:TRP:CE3	2.37	0.59
1:A:229:GLU:HA	1:A:232:MET:HG3	1.87	0.57
1:A:262:TYR:CG	1:A:262:TYR:O	2.57	0.57
1:A:255:ASN:HD22	1:A:256:GLU:H	1.49	0.56
1:A:145:VAL:HG12	1:A:146:TYR:CD2	2.42	0.55
1:A:110:PRO:O	1:A:113:VAL:HG12	2.07	0.54
1:A:25:LEU:HB3	1:A:42:ARG:HD3	1.89	0.54
1:A:6:ILE:HA	1:A:7[A]:LYS:C	2.28	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:253:LYS:HA	1:A:273:LEU:HD23	1.90	0.54
1:A:258:LYS:CB	1:A:267:TRP:CZ3	2.80	0.54
1:A:58:GLU:HG3	1:A:59:VAL:N	2.22	0.53
1:A:260:LYS:HG3	1:A:262:TYR:HB3	1.90	0.53
1:A:229:GLU:HA	1:A:232:MET:SD	2.50	0.52
1:A:6:ILE:HA	1:A:7[B]:LYS:C	2.30	0.52
1:A:128:LEU:HD23	1:A:192:ILE:HB	1.92	0.52
1:A:175:ASP:OD1	1:A:176:ARG:HB3	2.10	0.51
1:A:186:ASP:O	1:A:188:PRO:HD3	2.10	0.51
1:A:187:PHE:HE2	1:A:192:ILE:HD11	1.74	0.51
1:A:98:LYS:NZ	3:A:380:HOH:O	2.43	0.51
1:A:259:ILE:O	1:A:261:ARG:N	2.41	0.51
1:A:259:ILE:CG2	1:A:260:LYS:N	2.66	0.51
1:A:254:LEU:O	1:A:255:ASN:HB2	2.10	0.51
1:A:9:ARG:CG	1:A:9:ARG:HH11	2.18	0.50
1:A:110:PRO:C	1:A:112:ASN:H	2.15	0.50
1:A:261:ARG:HH12	1:A:263:ALA:N	2.10	0.49
1:A:7[A]:LYS:O	1:A:8:PRO:C	2.50	0.49
1:A:19:PRO:HG2	1:A:22:LEU:HD12	1.95	0.49
1:A:7[A]:LYS:O	1:A:10:ILE:N	2.46	0.49
1:A:274:ARG:HD2	1:A:276:PHE:CE2	2.47	0.49
1:A:258:LYS:HB3	1:A:267:TRP:HZ3	1.70	0.49
1:A:261:ARG:NH1	1:A:263:ALA:N	2.61	0.48
1:A:260:LYS:NZ	3:A:373:HOH:O	2.44	0.48
1:A:7[A]:LYS:N	1:A:7[A]:LYS:HE3	2.26	0.48
1:A:129:VAL:HG13	1:A:195:ARG:HB3	1.96	0.47
1:A:224:HIS:HE1	3:A:293:HOH:O	1.97	0.47
1:A:150:LYS:NZ	3:A:393:HOH:O	2.47	0.47
1:A:16:LYS:NZ	1:A:16:LYS:HB2	2.30	0.47
1:A:185:ARG:NH2	3:A:339:HOH:O	2.48	0.47
1:A:54:GLU:HG3	1:A:84:TYR:CE2	2.50	0.46
1:A:187:PHE:CE2	1:A:192:ILE:HD11	2.51	0.46
1:A:39:LEU:HD12	1:A:40:PRO:HD2	1.99	0.45
1:A:7[A]:LYS:HB3	1:A:8:PRO:HD3	1.99	0.45
1:A:55:VAL:O	1:A:59:VAL:HG13	2.17	0.45
1:A:157:ASP:HA	1:A:158:PRO:HD3	1.83	0.45
1:A:18:LEU:HA	1:A:19:PRO:HD2	1.85	0.44
1:A:277:LYS:O	1:A:278:SER:HB2	2.17	0.44
1:A:7[B]:LYS:O	1:A:10:ILE:N	2.50	0.43
1:A:262:TYR:O	1:A:263:ALA:O	2.37	0.43
1:A:7[B]:LYS:O	1:A:8:PRO:C	2.55	0.43
1:A:168:ILE:HD11	1:A:177:MET:HG2	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:118:ARG:HD2	1:A:257:LEU:HD11	2.01	0.43
1:A:261:ARG:CG	1:A:267:TRP:CE2	2.92	0.43
1:A:249:TYR:CD2	1:A:277:LYS:HA	2.55	0.42
1:A:229:GLU:HA	1:A:232:MET:CG	2.48	0.42
1:A:261:ARG:HH12	1:A:263:ALA:CA	2.31	0.42
1:A:7[A]:LYS:HB3	1:A:8:PRO:CD	2.49	0.42
1:A:46:GLU:OE2	1:A:67:ARG:NH2	2.46	0.42
1:A:37:LEU:HB2	1:A:62:VAL:HG21	2.01	0.41
1:A:250:ASP:N	3:A:296:HOH:O	2.46	0.41
1:A:251:VAL:HA	1:A:274:ARG:O	2.20	0.41
1:A:250:ASP:CB	3:A:296:HOH:O	2.68	0.41
1:A:21:GLU:CD	1:A:21:GLU:H	2.24	0.40
1:A:263:ALA:CB	1:A:264:PRO:CA	2.84	0.40
1:A:236:PRO:HD2	1:A:271:LEU:HD21	2.02	0.40
1:A:195:ARG:HA	1:A:220:ILE:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	261/301 (87%)	236 (90%)	16 (6%)	9 (3%)	6 3

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	7[A]	LYS
1	A	7[B]	LYS
1	A	263	ALA
1	A	260	LYS
1	A	261	ARG
1	A	49	LYS
1	A	259	ILE
1	A	255	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	264	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/267 (88%)	213 (90%)	23 (10%)	12	13

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7[A]	LYS
1	A	7[B]	LYS
1	A	9	ARG
1	A	16	LYS
1	A	35	ASP
1	A	37	LEU
1	A	38	LEU
1	A	51	ARG
1	A	55	VAL
1	A	58	GLU
1	A	59	VAL
1	A	60	LEU
1	A	79	ASP
1	A	172	LYS
1	A	192	ILE
1	A	241	LYS
1	A	250	ASP
1	A	255	ASN
1	A	260	LYS
1	A	261	ARG
1	A	266	VAL
1	A	268	HIS
1	A	278	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	191	ASN
1	A	224	HIS
1	A	255	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	SAM	A	279	-	26,29,29	1.13	2 (7%)	38,42,42	2.31	11 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAM	A	279	-	-	0/13/33/33	0/1/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	279	SAM	C2-N3	3.37	1.38	1.32
2	A	279	SAM	C2-N1	2.49	1.38	1.33

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	279	SAM	N3-C2-N1	-9.98	120.37	128.71
2	A	279	SAM	CG-CB-CA	3.78	117.78	112.22
2	A	279	SAM	N3-C4-N9	3.72	132.16	125.43
2	A	279	SAM	CB-CA-C	3.42	116.59	111.44
2	A	279	SAM	O4'-C1'-N9	3.29	111.50	108.44
2	A	279	SAM	O4'-C4'-C5'	2.68	115.96	108.95
2	A	279	SAM	C5-C4-N3	-2.46	120.34	125.70
2	A	279	SAM	C4-C5-N7	-2.23	107.61	109.52
2	A	279	SAM	C2-N3-C4	2.23	120.36	114.01
2	A	279	SAM	N7-C8-N9	-2.18	108.21	114.36
2	A	279	SAM	C5'-SD-CG	2.16	109.48	102.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/301 (87%)	0.13	5 (1%) 64 73	39, 64, 94, 105	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	267	TRP	2.9
1	A	262	TYR	2.5
1	A	148	LYS	2.3
1	A	168	ILE	2.2
1	A	278	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SAM	A	279	27/27	0.13	0.10	51,57,90,96	0

6.5 Other polymers ⓘ

There are no such residues in this entry.