



# wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 04:23 PM GMT

PDB ID : 4A21  
Title : STRUCTURE OF MYCOBACTERIUM TUBERCULOSIS FRUCTOSE 1,6-BISPHOSPHATE ALDOLASE BOUND TO SULFATE  
Authors : Coincon, M.; De La Paz Santangelo, M.; Gest, P.M.; Guerin, M.E.; Pham, H.; Ryan, G.; Puckett, S.E.; Spencer, J.S.; Gonzalez-Juarrero, M.; Daher, R.; Lenaerts, A.J.; Schnappinger, D.; Therisod, M.; Ehrt, S.; Jackson, M.; Sygusch, J.  
Deposited on : 2011-09-21  
Resolution : 2.35 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3327 (2.40-2.32)
Clashscore	79885	1064 (2.38-2.34)
Ramachandran outliers	78287	1048 (2.38-2.34)
Sidechain outliers	78261	1049 (2.38-2.34)
RSRZ outliers	66119	3330 (2.40-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	344	
1	B	344	
1	C	344	
1	D	344	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	NA	A	1347	-	X

## 2 Entry composition (i)


There are 4 unique types of molecules in this entry. The entry contains 19522 atoms, of which 9296 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FRUCTOSE-BISPHOSPHATEALDOLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	314	4671	1488	2325	395	453	10	0	0	0
1	B	314	4671	1488	2325	395	453	10	0	0	0
1	C	314	4669	1488	2323	395	453	10	0	0	0
1	D	314	4669	1488	2323	395	453	10	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Na	0	0
			2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	215	Total	O	0	0
			215	215		
4	B	205	Total	O	0	0
			205	205		
4	C	185	Total	O	0	0
			185	185		
4	D	195	Total	O	0	0
			195	195		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FRUCTOSE-BISPHOSPHATEALDOLASE

Chain A:




- Molecule 1: FRUCTOSE-BISPHOSPHATEALDOLASE


Chain B:



- Molecule 1: FRUCTOSE-BISPHOSPHATEALDOLASE


Chain C:





● Molecule 1: FRUCTOSE-BISPHOSPHATEALDOLASE

Chain D:



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	336.94Å 43.23Å 103.02Å 90.00° 99.57° 90.00°	Depositor
Resolution (Å)	42.87 – 2.35 42.87 – 2.35	Depositor EDS
% Data completeness (in resolution range)	91.2 (42.87-2.35) 99.1 (42.87-2.35)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.07 (at 2.34Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.211 , 0.261 0.215 , 0.257	Depositor DCC
$R_{free}$ test set	4631 reflections (7.52%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.4	Xtrriage
Anisotropy	0.214	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 31.8	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Outliers	0 of 61593 reflections	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	19522	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/2391	0.51	0/3236
1	B	0.36	0/2391	0.52	0/3236
1	C	0.33	0/2391	0.48	0/3236
1	D	0.35	0/2391	0.50	0/3236
All	All	0.35	0/9564	0.50	0/12944

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2346	2325	0	52	0
1	B	2346	2325	0	65	0
1	C	2346	2323	0	66	0
1	D	2346	2323	0	60	0
2	A	10	0	0	2	0
2	B	10	0	0	0	0
2	C	10	0	0	0	0
2	D	10	0	0	2	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	2	0	0	0	0
4	A	215	0	0	21	1
4	B	205	0	0	22	1
4	C	185	0	0	22	1
4	D	195	0	0	14	0
All	All	10226	9296	0	237	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 237 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:258:LEU:HB3	4:A:2170:HOH:O	1.57	1.03
1:B:257:SER:HA	4:B:2161:HOH:O	1.58	1.03
1:A:181:TYR:CZ	4:A:2138:HOH:O	2.22	0.92
1:C:267:ARG:HA	4:C:2033:HOH:O	1.71	0.89
1:C:163:GLY:HA3	1:C:207:THR:OG1	1.85	0.77

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:A:2126:HOH:O	4:A:2211:HOH:O[1_545]	2.17	0.03
4:B:2180:HOH:O	4:C:2159:HOH:O[1_565]	2.17	0.03

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	308/344 (90%)	297 (96%)	10 (3%)	1 (0%)	50 62
1	B	308/344 (90%)	293 (95%)	12 (4%)	3 (1%)	22 24
1	C	308/344 (90%)	290 (94%)	14 (4%)	4 (1%)	18 17
1	D	308/344 (90%)	296 (96%)	10 (3%)	2 (1%)	33 39

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1232/1376 (90%)	1176 (96%)	46 (4%)	10 (1%)	27 31

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	255	SER
1	A	96	HIS
1	C	96	HIS
1	D	96	HIS
1	B	303	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	242/266 (91%)	237 (98%)	5 (2%)	66 82
1	B	242/266 (91%)	236 (98%)	6 (2%)	60 76
1	C	242/266 (91%)	234 (97%)	8 (3%)	50 65
1	D	242/266 (91%)	236 (98%)	6 (2%)	60 76
All	All	968/1064 (91%)	943 (97%)	25 (3%)	59 75

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	95	ASP
1	C	146	GLU
1	D	258	LEU
1	C	127	SER
1	C	207	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	B	122	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	252	HIS
1	D	326	GLN

### 5.3.3 RNA [i](#)

There are no RNA chains in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	1344	-	4,4,4	0.19	0	6,6,6	0.12	0
2	SO4	A	1345	-	4,4,4	0.19	0	6,6,6	0.13	0
2	SO4	B	1344	-	4,4,4	0.11	0	6,6,6	0.24	0
2	SO4	B	1345	-	4,4,4	0.16	0	6,6,6	0.19	0
2	SO4	C	1344	-	4,4,4	0.12	0	6,6,6	0.13	0
2	SO4	C	1345	-	4,4,4	0.08	0	6,6,6	0.16	0
2	SO4	D	1344	-	4,4,4	0.11	0	6,6,6	0.33	0
2	SO4	D	1345	-	4,4,4	0.21	0	6,6,6	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1344	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1345	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1344	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1345	-	-	0/0/0/0	0/0/0/0
2	SO4	C	1344	-	-	0/0/0/0	0/0/0/0
2	SO4	C	1345	-	-	0/0/0/0	0/0/0/0
2	SO4	D	1344	-	-	0/0/0/0	0/0/0/0
2	SO4	D	1345	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	314/344 (91%)	0.86	37 (11%) 5 6	14, 27, 52, 81	0
1	B	314/344 (91%)	0.82	36 (11%) 5 7	15, 27, 49, 91	0
1	C	314/344 (91%)	1.07	50 (15%) 3 3	18, 35, 63, 99	0
1	D	314/344 (91%)	0.78	28 (8%) 10 11	15, 27, 49, 77	0
All	All	1256/1376 (91%)	0.88	151 (12%) 5 6	14, 29, 55, 99	0

The worst 5 of 151 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	254	GLY	7.6
1	C	256	GLY	7.0
1	C	163	GLY	6.9
1	B	257	SER	6.4
1	D	305	VAL	5.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NA	A	1347	1/1	0.30	3.97	43,43,43,43	0
2	SO4	C	1345	5/5	0.19	1.66	50,50,68,80	0
3	NA	A	1346	1/1	0.18	-0.03	47,47,47,47	0
2	SO4	C	1344	5/5	0.26	-0.15	51,67,74,83	0
2	SO4	D	1344	5/5	0.16	-0.26	38,43,63,87	0
2	SO4	B	1345	5/5	0.19	-0.52	50,54,67,83	0
2	SO4	D	1345	5/5	0.20	-0.58	57,59,68,88	0
2	SO4	A	1345	5/5	0.14	-1.27	38,46,52,60	0
2	SO4	A	1344	5/5	0.10	-1.35	52,55,64,65	0
2	SO4	B	1344	5/5	0.14	-1.55	41,44,65,75	0

## 6.5 Other polymers

There are no such residues in this entry.