



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 04:22 PM GMT

PDB ID : 4A2E
Title : CRYSTAL STRUCTURE OF A CORIOLOPSIS GALLICA LACCASE AT
1.7 Å RESOLUTION PH 5.5
Authors : De La Mora, E.; Valderrama, B.; Horjales, E.; Rudino-Pinera, E.
Deposited on : 2011-09-26
Resolution : 1.80 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance


The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3513 (1.80-1.80)
Clashscore	79885	4461 (1.80-1.80)
Ramachandran outliers	78287	4404 (1.80-1.80)
Sidechain outliers	78261	4403 (1.80-1.80)
RSRZ outliers	66119	3515 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	496	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4466 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LACCASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	496	Total	C	N	O	S	0	10	0
			3800	2411	640	740	9			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASP	TYR	CONFLICT	UNP Q1W6B1
A	151	ASN	GLN	CONFLICT	UNP Q1W6B1
A	178	LYS	ARG	CONFLICT	UNP Q1W6B1
A	182	PRO	ALA	CONFLICT	UNP Q1W6B1
A	183	VAL	ILE	CONFLICT	UNP Q1W6B1
A	229	TYR	HIS	CONFLICT	UNP Q1W6B1
A	256	LEU	ILE	CONFLICT	UNP Q1W6B1
A	287	THR	ASN	CONFLICT	UNP Q1W6B1
A	288	GLN	THR	CONFLICT	UNP Q1W6B1
A	294	THR	VAL	CONFLICT	UNP Q1W6B1
A	314	THR	ALA	CONFLICT	UNP Q1W6B1
A	329	LYS	GLU	CONFLICT	UNP Q1W6B1
A	356	ASN	ARG	CONFLICT	UNP Q1W6B1
A	358	THR	SER	CONFLICT	UNP Q1W6B1
A	423	VAL	ALA	CONFLICT	UNP Q1W6B1

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Cu	0	0
			4	4		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	A	2	Total	C	N	O	0	0
			28	16	2	10		

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASP	TYR	CONFLICT	UNP Q1W6B1
A	151	ASN	GLN	CONFLICT	UNP Q1W6B1
A	178	LYS	ARG	CONFLICT	UNP Q1W6B1
A	182	PRO	ALA	CONFLICT	UNP Q1W6B1
A	183	VAL	ILE	CONFLICT	UNP Q1W6B1
A	229	TYR	HIS	CONFLICT	UNP Q1W6B1
A	256	LEU	ILE	CONFLICT	UNP Q1W6B1
A	287	THR	ASN	CONFLICT	UNP Q1W6B1
A	288	GLN	THR	CONFLICT	UNP Q1W6B1
A	294	THR	VAL	CONFLICT	UNP Q1W6B1
A	314	THR	ALA	CONFLICT	UNP Q1W6B1
A	329	LYS	GLU	CONFLICT	UNP Q1W6B1
A	356	ASN	ARG	CONFLICT	UNP Q1W6B1
A	358	THR	SER	CONFLICT	UNP Q1W6B1
A	423	VAL	ALA	CONFLICT	UNP Q1W6B1
A	39	ASP	TYR	CONFLICT	UNP Q1W6B1
A	151	ASN	GLN	CONFLICT	UNP Q1W6B1
A	178	LYS	ARG	CONFLICT	UNP Q1W6B1
A	182	PRO	ALA	CONFLICT	UNP Q1W6B1
A	183	VAL	ILE	CONFLICT	UNP Q1W6B1
A	229	TYR	HIS	CONFLICT	UNP Q1W6B1
A	256	LEU	ILE	CONFLICT	UNP Q1W6B1
A	287	THR	ASN	CONFLICT	UNP Q1W6B1
A	288	GLN	THR	CONFLICT	UNP Q1W6B1
A	294	THR	VAL	CONFLICT	UNP Q1W6B1
A	314	THR	ALA	CONFLICT	UNP Q1W6B1
A	329	LYS	GLU	CONFLICT	UNP Q1W6B1
A	356	ASN	ARG	CONFLICT	UNP Q1W6B1
A	358	THR	SER	CONFLICT	UNP Q1W6B1
A	423	VAL	ALA	CONFLICT	UNP Q1W6B1

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	606	Total 606	O 606	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LACCASE

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.38Å 86.13Å 152.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.63 – 1.80 25.63 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.6 (25.63-1.80) 96.6 (25.63-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.88 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.154 , 0.187 0.154 , 0.188	Depositor DCC
R_{free} test set	3389 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	16.4	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 35.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 67354 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4466	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.48	18/3925 (0.5%)	1.13	16/5398 (0.3%)

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	221	VAL	CB-CG1	-8.43	1.35	1.52
1	A	517	GLN	C-OXT	7.86	1.38	1.23
1	A	464	PRO	CA-CB	7.59	1.68	1.53
1	A	301	TYR	CD1-CE1	7.24	1.50	1.39
1	A	423	VAL	CB-CG2	-7.09	1.38	1.52

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	282	LEU	CB-CG-CD1	7.20	123.23	111.00
1	A	485	MET	CG-SD-CE	-6.92	89.14	100.20
1	A	271	ASP	CB-CG-OD1	6.69	124.32	118.30
1	A	142	ARG	NE-CZ-NH2	-6.40	117.10	120.30
1	A	371	VAL	CG1-CB-CG2	-6.33	100.78	110.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the

chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3800	0	3601	33	0
2	A	4	0	0	0	0
3	A	56	0	50	0	0
4	A	606	0	0	6	0
All	All	4466	0	3651	33	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

The worst 5 of 33 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:418[B]:HIS:HE1	1:A:470:HIS:CE1	1.36	1.41
1:A:418[B]:HIS:CE1	1:A:470:HIS:CE1	2.09	1.40
1:A:329:LYS:HE2	1:A:329:LYS:H	1.13	1.08
1:A:418[B]:HIS:CE1	1:A:470:HIS:NE2	2.32	0.97
1:A:418[B]:HIS:HE1	1:A:470:HIS:HE1	1.09	0.89

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	504/496 (102%)	487 (97%)	15 (3%)	2 (0%)	43 25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	226	ASP
1	A	79	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	411/401 (102%)	402 (98%)	9 (2%)	64 48

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	329	LYS
1	A	481	PHE
1	A	439	ASN
1	A	282	LEU
1	A	407	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	269	ASN
1	A	272	GLN
1	A	380	GLN
1	A	251	HIS
1	A	422	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	1522	1,3	12,14,15	1.23	2 (16%)	15,19,21	1.92	2 (13%)
3	NAG	A	1523	3	12,14,15	1.66	3 (25%)	15,19,21	1.60	4 (26%)
3	NAG	A	1524	1,3	12,14,15	1.31	1 (8%)	15,19,21	2.13	6 (40%)
3	NAG	A	1525	3	12,14,15	1.03	0	15,19,21	1.43	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1522	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1523	3	-	0/6/23/26	0/1/1/1
3	NAG	A	1524	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1525	3	-	0/6/23/26	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1524	NAG	C2-N2	3.32	1.50	1.46
3	A	1523	NAG	C3-C2	2.92	1.58	1.52
3	A	1522	NAG	C4-C5	2.89	1.59	1.53
3	A	1523	NAG	O4-C4	2.54	1.49	1.43
3	A	1523	NAG	C2-N2	2.49	1.49	1.46

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1522	NAG	O5-C5-C6	6.06	113.33	106.98
3	A	1524	NAG	O5-C5-C4	-3.79	105.84	110.65
3	A	1524	NAG	C2-N2-C7	-3.41	117.36	123.09
3	A	1524	NAG	O7-C7-N2	3.30	128.80	121.90
3	A	1525	NAG	C3-C2-N2	-3.10	107.04	111.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	496/496 (100%)	-0.68	0 100 100	8, 13, 22, 34	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	A	1525	14/15	0.18	14.80	16,27,31,37	0
3	NAG	A	1523	14/15	0.13	2.30	10,17,26,27	0
3	NAG	A	1524	14/15	0.07	0.04	12,17,26,28	0
3	NAG	A	1522	14/15	0.05	-0.57	12,15,19,20	0

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy

less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CU	A	1520	1/1	0.02	-1.82	17,17,17,17	1
2	CU	A	1521	1/1	0.03	-1.99	12,12,12,12	1
2	CU	A	1518	1/1	0.02	-2.37	14,14,14,14	1
2	CU	A	1519	1/1	0.02	-4.15	12,12,12,12	1

6.5 Other polymers ⓘ

There are no such residues in this entry.