



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 02:35 PM GMT

PDB ID : 4A2G
Title : Coriolopsis gallica laccase collected at 8.98 keV
Authors : De La Mora, E.; Rudino-Pinera, E.
Deposited on : 2011-09-26
Resolution : 1.80 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance


The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3513 (1.80-1.80)
Clashscore	79885	4461 (1.80-1.80)
Ramachandran outliers	78287	4404 (1.80-1.80)
Sidechain outliers	78261	4403 (1.80-1.80)
RSRZ outliers	66119	3515 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	496	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4374 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LACCASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	496	Total	C	N	O	S	0	9	0
			3792	2406	638	739	9			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASP	TYR	CONFLICT	UNP Q1W6B1
A	151	ASN	GLN	CONFLICT	UNP Q1W6B1
A	155	ALA	LYS	CONFLICT	UNP Q1W6B1
A	178	LYS	ARG	CONFLICT	UNP Q1W6B1
A	181	ALA	PRO	CONFLICT	UNP Q1W6B1
A	182	PRO	ALA	CONFLICT	UNP Q1W6B1
A	183	VAL	ILE	CONFLICT	UNP Q1W6B1
A	198	ALA	ILE	CONFLICT	UNP Q1W6B1
A	199	ALA	ASN	CONFLICT	UNP Q1W6B1
A	202	ALA	ASN	CONFLICT	UNP Q1W6B1
A	229	TYR	HIS	CONFLICT	UNP Q1W6B1
A	256	LEU	ILE	CONFLICT	UNP Q1W6B1
A	287	THR	ASN	CONFLICT	UNP Q1W6B1
A	288	GLN	THR	CONFLICT	UNP Q1W6B1
A	291	ALA	ASP	CONFLICT	UNP Q1W6B1
A	294	THR	VAL	CONFLICT	UNP Q1W6B1
A	314	THR	ALA	CONFLICT	UNP Q1W6B1
A	329	LYS	GLU	CONFLICT	UNP Q1W6B1
A	356	ASN	ARG	CONFLICT	UNP Q1W6B1
A	358	THR	SER	CONFLICT	UNP Q1W6B1
A	383	ALA	GLN	CONFLICT	UNP Q1W6B1
A	388	ALA	THR	CONFLICT	UNP Q1W6B1
A	408	ALA	THR	CONFLICT	UNP Q1W6B1
A	423	VAL	ALA	CONFLICT	UNP Q1W6B1
A	438	ALA	GLU	CONFLICT	UNP Q1W6B1
A	450	ALA	SER	CONFLICT	UNP Q1W6B1

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Cu 2 2	0	0

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total C N O 28 16 2 10	0	0
3	A	2	Total C N O 28 16 2 10	0	0

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASP	TYR	CONFLICT	UNP Q1W6B1
A	151	ASN	GLN	CONFLICT	UNP Q1W6B1
A	155	ALA	LYS	CONFLICT	UNP Q1W6B1
A	178	LYS	ARG	CONFLICT	UNP Q1W6B1
A	181	ALA	PRO	CONFLICT	UNP Q1W6B1
A	182	PRO	ALA	CONFLICT	UNP Q1W6B1
A	183	VAL	ILE	CONFLICT	UNP Q1W6B1
A	198	ALA	ILE	CONFLICT	UNP Q1W6B1
A	199	ALA	ASN	CONFLICT	UNP Q1W6B1
A	202	ALA	ASN	CONFLICT	UNP Q1W6B1
A	229	TYR	HIS	CONFLICT	UNP Q1W6B1
A	256	LEU	ILE	CONFLICT	UNP Q1W6B1
A	287	THR	ASN	CONFLICT	UNP Q1W6B1
A	288	GLN	THR	CONFLICT	UNP Q1W6B1
A	291	ALA	ASP	CONFLICT	UNP Q1W6B1
A	294	THR	VAL	CONFLICT	UNP Q1W6B1
A	314	THR	ALA	CONFLICT	UNP Q1W6B1
A	329	LYS	GLU	CONFLICT	UNP Q1W6B1
A	356	ASN	ARG	CONFLICT	UNP Q1W6B1
A	358	THR	SER	CONFLICT	UNP Q1W6B1
A	383	ALA	GLN	CONFLICT	UNP Q1W6B1
A	388	ALA	THR	CONFLICT	UNP Q1W6B1
A	408	ALA	THR	CONFLICT	UNP Q1W6B1
A	423	VAL	ALA	CONFLICT	UNP Q1W6B1
A	438	ALA	GLU	CONFLICT	UNP Q1W6B1
A	450	ALA	SER	CONFLICT	UNP Q1W6B1
A	39	ASP	TYR	CONFLICT	UNP Q1W6B1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	151	ASN	GLN	CONFLICT	UNP Q1W6B1
A	155	ALA	LYS	CONFLICT	UNP Q1W6B1
A	178	LYS	ARG	CONFLICT	UNP Q1W6B1
A	181	ALA	PRO	CONFLICT	UNP Q1W6B1
A	182	PRO	ALA	CONFLICT	UNP Q1W6B1
A	183	VAL	ILE	CONFLICT	UNP Q1W6B1
A	198	ALA	ILE	CONFLICT	UNP Q1W6B1
A	199	ALA	ASN	CONFLICT	UNP Q1W6B1
A	202	ALA	ASN	CONFLICT	UNP Q1W6B1
A	229	TYR	HIS	CONFLICT	UNP Q1W6B1
A	256	LEU	ILE	CONFLICT	UNP Q1W6B1
A	287	THR	ASN	CONFLICT	UNP Q1W6B1
A	288	GLN	THR	CONFLICT	UNP Q1W6B1
A	291	ALA	ASP	CONFLICT	UNP Q1W6B1
A	294	THR	VAL	CONFLICT	UNP Q1W6B1
A	314	THR	ALA	CONFLICT	UNP Q1W6B1
A	329	LYS	GLU	CONFLICT	UNP Q1W6B1
A	356	ASN	ARG	CONFLICT	UNP Q1W6B1
A	358	THR	SER	CONFLICT	UNP Q1W6B1
A	383	ALA	GLN	CONFLICT	UNP Q1W6B1
A	388	ALA	THR	CONFLICT	UNP Q1W6B1
A	408	ALA	THR	CONFLICT	UNP Q1W6B1
A	423	VAL	ALA	CONFLICT	UNP Q1W6B1
A	438	ALA	GLU	CONFLICT	UNP Q1W6B1
A	450	ALA	SER	CONFLICT	UNP Q1W6B1

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	524	Total O 524 524	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LACCASE

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.01Å 85.23Å 151.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	75.76 – 1.80 34.33 – 1.80	Depositor EDS
% Data completeness (in resolution range)	94.0 (75.76-1.80) 94.0 (34.33-1.80)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.02 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.157 , 0.188 0.160 , 0.192	Depositor DCC
R_{free} test set	3240 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	17.9	Xtriage
Anisotropy	0.098	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 38.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 64012 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4374	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.26	13/3913 (0.3%)	1.05	10/5383 (0.2%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	221	VAL	CB-CG1	-8.42	1.35	1.52
1	A	242	GLU	CG-CD	6.73	1.62	1.51
1	A	471	CYS	CB-SG	-5.94	1.72	1.81
1	A	26	VAL	CB-CG1	5.72	1.64	1.52
1	A	137	TYR	CD2-CE2	5.66	1.47	1.39
1	A	101	ALA	CA-CB	5.47	1.64	1.52
1	A	447	SER	CB-OG	5.37	1.49	1.42
1	A	137	TYR	CD1-CE1	5.27	1.47	1.39
1	A	245	SER	CB-OG	5.23	1.49	1.42
1	A	148	TYR	CD1-CE1	5.11	1.47	1.39
1	A	231	PHE	CE1-CZ	5.10	1.47	1.37
1	A	120	VAL	CB-CG1	5.07	1.63	1.52
1	A	401	GLU	CD-OE1	-5.04	1.20	1.25

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	142	ARG	NE-CZ-NH2	-8.61	116.00	120.30
1	A	271	ASP	CB-CG-OD1	7.31	124.88	118.30
1	A	462	ASP	CB-CG-OD1	7.27	124.84	118.30
1	A	343[A]	ASP	CB-CG-OD2	6.47	124.12	118.30
1	A	343[B]	ASP	CB-CG-OD2	6.47	124.12	118.30
1	A	254	ASP	CB-CG-OD1	5.76	123.48	118.30
1	A	444	ASP	CB-CG-OD2	5.53	123.27	118.30
1	A	244	ASP	CB-CG-OD1	5.40	123.16	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	219	ARG	NE-CZ-NH2	-5.16	117.72	120.30
1	A	214	LYS	CD-CE-NZ	-5.01	100.17	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3792	0	3595	34	0
2	A	2	0	0	0	0
3	A	56	0	50	3	0
4	A	524	0	0	10	0
All	All	4374	0	3645	36	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

All (36) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:329:LYS:HE2	1:A:329:LYS:H	1.26	0.98
1:A:475:PHE:HD1	4:A:2248:HOH:O	1.54	0.89
1:A:22:ALA:HB2	1:A:163:ASP:OD2	1.83	0.79
1:A:422:HIS:HE1	1:A:487:GLU:OE1	1.67	0.77
1:A:276:ASN:HD21	1:A:300:ARG:HE	1.35	0.75
1:A:475:PHE:CD1	4:A:2248:HOH:O	2.36	0.72
3:A:1523:NAG:O4	4:A:2524:HOH:O	2.11	0.68
1:A:276:ASN:ND2	1:A:300:ARG:HE	1.91	0.67
1:A:284:ASN:HD22	1:A:284:ASN:H	1.46	0.63
1:A:33:ASN:ND2	1:A:74:THR:H	2.01	0.59
1:A:242:GLU:HB3	1:A:265:SER:HB2	1.86	0.57
1:A:436:ASN:HD21	1:A:439:ASN:ND2	2.02	0.57
1:A:422:HIS:CE1	1:A:487:GLU:OE1	2.52	0.57
3:A:1521:NAG:O4	4:A:2514:HOH:O	2.18	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:475:PHE:HA	4:A:2248:HOH:O	2.06	0.56
1:A:329:LYS:CE	1:A:329:LYS:H	2.11	0.54
1:A:33:ASN:HD22	1:A:74:THR:H	1.55	0.52
1:A:251:HIS:HE1	4:A:2318:HOH:O	1.92	0.52
1:A:22:ALA:HB3	4:A:2003:HOH:O	2.08	0.52
1:A:329:LYS:HE2	1:A:329:LYS:N	2.10	0.51
1:A:92:HIS:HD2	4:A:2124:HOH:O	1.94	0.50
1:A:211:THR:OG1	1:A:302[B]:ASP:OD2	2.24	0.47
1:A:135:THR:HA	1:A:474:ASP:OD2	2.13	0.47
1:A:269:ASN:C	1:A:269:ASN:HD22	2.18	0.47
1:A:251:HIS:HD2	4:A:2316:HOH:O	1.98	0.46
1:A:87:HIS:CE1	1:A:261:ALA:HB1	2.50	0.46
1:A:227:PRO:HA	1:A:284:ASN:HD21	1.82	0.45
1:A:269:ASN:HD22	1:A:270:ALA:N	2.15	0.44
1:A:84[A]:ILE:HG21	1:A:86:TRP:CH2	2.54	0.43
1:A:43:ARG:HB2	3:A:1521:NAG:H83	2.01	0.42
1:A:228:ASN:H	1:A:284:ASN:ND2	2.18	0.42
1:A:269:ASN:HB2	4:A:2331:HOH:O	2.20	0.41
1:A:120:VAL:O	1:A:120:VAL:HG23	2.20	0.41
1:A:85:HIS:HA	1:A:98:ASP:O	2.21	0.40
1:A:239:THR:HA	1:A:252:THR:HA	2.02	0.40
1:A:226:ASP:HB3	1:A:227:PRO:CD	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	503/496 (101%)	486 (97%)	15 (3%)	2 (0%)	43 25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	226	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	79	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/401 (102%)	400 (98%)	10 (2%)	61	44

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	263	ARG
1	A	269	ASN
1	A	282	LEU
1	A	284	ASN
1	A	329	LYS
1	A	407	THR
1	A	439	ASN
1	A	468	PHE
1	A	472	HIS
1	A	481	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	A	44	GLN
1	A	91	GLN
1	A	92	HIS
1	A	104	ASN
1	A	105	GLN
1	A	251	HIS
1	A	269	ASN
1	A	276	ASN
1	A	284	ASN
1	A	318	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	356	ASN
1	A	380	GLN
1	A	422	HIS
1	A	439	ASN
1	A	476	HIS
1	A	517	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1520	1,3	12,14,15	1.12	1 (8%)	15,19,21	1.97	3 (20%)
3	NAG	A	1521	3	12,14,15	1.64	3 (25%)	15,19,21	1.54	2 (13%)
3	NAG	A	1522	1,3	12,14,15	0.90	1 (8%)	15,19,21	1.54	3 (20%)
3	NAG	A	1523	3	12,14,15	0.67	0	15,19,21	1.17	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1520	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1521	3	-	0/6/23/26	0/1/1/1
3	NAG	A	1522	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1523	3	-	0/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1521	NAG	C2-N2	3.38	1.50	1.46
3	A	1520	NAG	O5-C5	-3.12	1.39	1.45
3	A	1521	NAG	O4-C4	2.28	1.48	1.43
3	A	1521	NAG	C3-C2	2.22	1.57	1.52
3	A	1522	NAG	C2-N2	2.11	1.48	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1520	NAG	O5-C5-C4	-4.56	104.87	110.65
3	A	1520	NAG	O5-C5-C6	4.54	111.75	106.98
3	A	1522	NAG	O5-C5-C4	-3.20	106.60	110.65
3	A	1521	NAG	O6-C6-C5	-3.04	100.88	111.36
3	A	1523	NAG	O6-C6-C5	-2.93	101.26	111.36
3	A	1522	NAG	C2-N2-C7	-2.92	118.19	123.09
3	A	1520	NAG	O4-C4-C5	-2.75	102.05	109.28
3	A	1522	NAG	O7-C7-N2	2.38	126.88	121.90
3	A	1521	NAG	O7-C7-N2	2.13	126.36	121.90
3	A	1523	NAG	C3-C2-N2	-2.11	108.55	111.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	496/496 (100%)	-0.53	4 (0%) 83 81	10, 15, 29, 39	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	353	ALA	3.4
1	A	182	PRO	2.4
1	A	477	LEU	2.2
1	A	352	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	A	1523	14/15	0.16	6.64	24,31,37,39	0
3	NAG	A	1521	14/15	0.12	1.92	13,21,25,28	0
3	NAG	A	1522	14/15	0.07	0.82	14,20,26,26	0
3	NAG	A	1520	14/15	0.06	-0.04	16,18,25,29	0

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CU	A	1519	1/1	0.05	-2.77	20,20,20,20	1
2	CU	A	1518	1/1	0.07	-5.79	20,20,20,20	1

6.5 Other polymers

There are no such residues in this entry.