



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 10:52 PM GMT

PDB ID : 3A3E
Title : Crystal structure of penicillin binding protein 4 (dacB) from Haemophilus influenzae, complexed with novel beta-lactam (CMV)
Authors : Kawai, F.; Roper, D.I.; Park, S.-Y.; Tame, J.R.H.
Deposited on : 2009-06-12
Resolution : 2.40 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2207 (2.40-2.40)
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)
RSRZ outliers	66119	2210 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	453	
1	B	453	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6865 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Penicillin-binding protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	438	Total	C	N	O	S	0	0	0
			3378	2154	582	630	12			
1	B	433	Total	C	N	O	S	0	0	0
			3333	2127	576	620	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	MET	-	EXPRESSION TAG	UNP A8E0K8
B	27	MET	-	EXPRESSION TAG	UNP A8E0K8

- Molecule 2 is (2R,4S)-2-[(1R)-1-({(2R)-2-[(4-ETHYL-2,3-DIOXOPIPERAZIN-1-YL)AMINO]-2-PHENYLACETYL}AMINO)-2-OXOETHYL]-5,5-DIMETHYL-1,3-THIAZOLIDINE-4-CARBOXYLICACID (three-letter code: CMV) (formula: C₂₂H₂₉N₅O₆S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			34	22	5	6	1		
2	B	1	Total	C	N	O	S	0	0
			34	22	5	6	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	60	Total	O	0	0
			60	60		
3	B	26	Total	O	0	0
			26	26		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Penicillin-binding protein 4

Chain A:



• Molecule 1: Penicillin-binding protein 4

Chain B:



T460	R461	A462	A463	P464	E469	R470	N471	L472	Y473	L476	Y477	R478	Y479
------	------	------	------	------	------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.90Å 92.82Å 104.17Å 90.00° 107.79° 90.00°	Depositor
Resolution (Å)	49.57 – 2.40 49.60 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.57-2.40) 93.0 (49.60-2.40)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, R_{free}	0.217 , 0.280 0.218 , 0.274	Depositor DCC
R_{free} test set	2163 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	37.6	Xtriage
Anisotropy	0.277	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 30.4	EDS
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 42895 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6865	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CMV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	3/3437 (0.1%)	1.03	3/4644 (0.1%)
1	B	0.93	1/3391 (0.0%)	0.99	6/4582 (0.1%)
All	All	0.99	4/6828 (0.1%)	1.01	9/9226 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	180	CYS	CB-SG	-7.14	1.70	1.82
1	B	378	GLU	CG-CD	5.59	1.60	1.51
1	A	148	VAL	CB-CG1	5.46	1.64	1.52
1	A	427	VAL	CB-CG1	5.09	1.63	1.52

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	170	PRO	C-N-CD	6.83	142.75	128.40
1	A	249	LEU	CA-CB-CG	-6.40	100.58	115.30
1	B	177	ASP	CB-CG-OD1	6.31	123.98	118.30
1	B	306	MET	CG-SD-CE	-5.75	91.01	100.20
1	B	456	LEU	CA-CB-CG	5.75	128.51	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	146	THR	Peptide
1	B	147	SER	Peptide

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3378	0	3440	149	0
1	B	3333	0	3400	193	0
2	A	34	0	26	5	0
2	B	34	0	27	0	0
3	A	60	0	0	3	0
3	B	26	0	0	1	0
All	All	6865	0	6893	341	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 25.

The worst 5 of 341 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:212:TYR:CE2	1:B:239:CYS:HB3	1.66	1.29
1:B:218:GLU:C	1:B:220:PRO:HD2	1.53	1.29
1:B:218:GLU:C	1:B:220:PRO:CD	2.05	1.25
1:A:146:THR:HG22	1:A:147:SER:N	1.45	1.17
1:A:146:THR:CG2	1:A:147:SER:H	1.58	1.17

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	434/453 (96%)	406 (94%)	20 (5%)	8 (2%)	13	15
1	B	429/453 (95%)	384 (90%)	31 (7%)	14 (3%)	6	5
All	All	863/906 (95%)	790 (92%)	51 (6%)	22 (2%)	9	8

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	147	SER
1	B	171	PRO
1	B	215	ASP
1	B	216	SER
1	B	218	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/378 (97%)	329 (90%)	36 (10%)	11	16
1	B	358/378 (95%)	322 (90%)	36 (10%)	11	16
All	All	723/756 (96%)	651 (90%)	72 (10%)	11	16

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	458	SER
1	B	100	GLN
1	B	375	SER
1	A	459	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	36	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	366	ASN
1	B	71	GLN
1	B	383	ASN
1	A	446	GLN
1	B	51	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CMV	A	1	1	36,36,36	3.65	8 (22%)	52,52,52	5.69	25 (48%)
2	CMV	B	1	1	36,36,36	3.55	8 (22%)	52,52,52	3.36	24 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CMV	A	1	1	-	3/23/60/60	0/3/3/3
2	CMV	B	1	1	-	2/23/60/60	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	CMV	NAN-NAG	-14.72	1.23	1.41
2	B	1	CMV	NAN-NAG	-14.27	1.23	1.41
2	B	1	CMV	CAE-CAI	-11.39	1.31	1.53
2	A	1	CMV	CAE-CAI	-9.02	1.36	1.53
2	A	1	CMV	CAE-NAG	-8.18	1.30	1.37

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	CMV	CAQ-NAG-NAN	28.56	130.43	109.56
2	A	1	CMV	CAM-NAN-NAG	19.96	136.08	112.56
2	B	1	CMV	CAM-NAN-NAG	13.52	128.49	112.56
2	A	1	CMV	CAE-NAG-NAN	-10.04	104.20	118.08
2	B	1	CMV	CAQ-NAG-NAN	9.77	116.70	109.56

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	CMV	CAM-NAN-NAG-CAQ
2	A	1	CMV	CAM-NAN-NAG-CAQ
2	A	1	CMV	CBE-CBB-NAO-CAI
2	B	1	CMV	CAM-NAN-NAG-CAE
2	A	1	CMV	CAM-NAN-NAG-CAE

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	438/453 (96%)	-0.00	3 (0%) 84 84	16, 32, 54, 76	0
1	B	433/453 (95%)	0.27	24 (5%) 24 22	18, 40, 72, 86	0
All	All	871/906 (96%)	0.14	27 (3%) 47 44	16, 36, 67, 86	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	456	LEU	6.8
1	B	217	ASN	4.7
1	B	220	PRO	4.7
1	B	457	GLU	4.2
1	B	221	TYR	4.2

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CMV	B	1	34/34	0.13	-0.58	25,33,54,57	0
2	CMV	A	1	34/34	0.14	-0.86	15,25,46,47	0

6.5 Other polymers ⓘ

There are no such residues in this entry.