



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 26, 2014 – 07:44 PM GMT

PDB ID : 1A5L
Title : K217C VARIANT OF KLEBSIELLA AEROGENES UREASE
Authors : Pearson, M.A.; Schaller, R.A.; Michel, L.O.; Karplus, P.A.; Hausinger, R.P.
Deposited on : 1998-02-17
Resolution : 2.20 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3751 (2.20-2.20)
Ramachandran outliers	78287	3681 (2.20-2.20)
Sidechain outliers	78261	3682 (2.20-2.20)
RSRZ outliers	66119	2939 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	100	
2	B	101	
3	C	566	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5731 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UREASE (GAMMA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	0	0	0
			776	491	134	146	5			

- Molecule 2 is a protein called UREASE (BETA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	101	Total	C	N	O	S	0	0	0
			785	496	150	136	3			

- Molecule 3 is a protein called UREASE (ALPHA SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	536	Total	C	N	O	S	0	1	0
			3987	2502	700	764	21			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	217	CYS	LYS	ENGINEERED	UNP P18314

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	17	Total	O	0	0
			17	17		
4	B	12	Total	O	0	0
			12	12		
4	C	154	Total	O	0	0
			154	154		

4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	170.80Å 170.80Å 170.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.20 29.29 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.0 (10.00-2.20) 89.8 (29.29-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.20Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.186 , (Not available) 0.183 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	14.8	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 23.7	EDS
Estimated twinning fraction	0.037 for -l,-k,-h	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 37711 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5731	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/787	0.60	0/1061
2	B	0.38	0/805	0.65	0/1087
3	C	0.38	0/4071	0.70	1/5548 (0.0%)
All	All	0.38	0/5663	0.68	1/7696 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	372	LEU	CA-CB-CG	5.78	128.60	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	776	0	807	4	0
2	B	785	0	775	9	0
3	C	3987	0	3944	54	0
4	A	17	0	0	1	0
4	B	12	0	0	1	0
4	C	154	0	0	2	0
All	All	5731	0	5526	64	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

All (64) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:C:299:ASN:HB2	3:C:373:ARG:HD2	1.56	0.86
3:C:297:SER:HB2	3:C:349:HIS:HE1	1.44	0.82
3:C:300:PRO:HG3	3:C:364:MET:O	1.81	0.81
3:C:302:LEU:HD22	3:C:377:VAL:HG21	1.74	0.70
3:C:359:SER:O	3:C:365:GLY:HA3	1.93	0.68
3:C:274:GLU:HG3	3:C:345:GLU:HB2	1.76	0.67
3:C:299:ASN:HB2	3:C:373:ARG:CD	2.26	0.64
3:C:218:ILE:HD13	3:C:243:VAL:HG13	1.81	0.62
3:C:545:VAL:O	3:C:548:GLU:HG2	2.02	0.60
3:C:188:PRO:O	3:C:449:LYS:HD2	2.04	0.57
3:C:299:ASN:O	3:C:373:ARG:HD3	2.05	0.56
2:B:7:HIS:HB3	3:C:20:LYS:HB2	1.87	0.56
3:C:201:GLN:HE21	3:C:203:ASP:H	1.52	0.56
3:C:285:ILE:HD11	3:C:348:LEU:CD1	2.37	0.55
3:C:303:PRO:O	3:C:305:THR:HG23	2.07	0.54
3:C:380:ARG:NE	3:C:555:ALA:HB2	2.23	0.54
3:C:359:SER:HB3	3:C:370:VAL:CG2	2.37	0.54
2:B:89:ALA:HA	2:B:100:PRO:HA	1.90	0.53
3:C:173:GLY:O	3:C:177:ILE:HG13	2.08	0.53
3:C:166:HIS:HD2	4:C:688:HOH:O	1.92	0.53
3:C:363:ALA:O	3:C:364:MET:HB2	2.08	0.52
2:B:72:GLU:H	2:B:75:GLN:NE2	2.08	0.52
3:C:236:ALA:O	3:C:518:ARG:HD2	2.11	0.51
2:B:4:GLY:HA2	3:C:22:ARG:O	2.10	0.51
3:C:296:SER:HB3	3:C:356:LEU:HB2	1.92	0.51
3:C:297:SER:HB2	3:C:349:HIS:CE1	2.35	0.50
2:B:83:ALA:HB1	2:B:88:ARG:HH21	1.76	0.50
3:C:298:THR:HG21	3:C:360:ASP:HB2	1.93	0.50
3:C:297:SER:CB	3:C:349:HIS:HE1	2.22	0.49
2:B:3:PRO:HA	3:C:24:ALA:O	2.12	0.49
3:C:285:ILE:HD11	3:C:348:LEU:HD11	1.94	0.49
3:C:435:TRP:CZ3	3:C:444:PRO:HG3	2.47	0.49
3:C:395:ASN:C	3:C:395:ASN:HD22	2.17	0.48
3:C:246:HIS:CG	3:C:272[A]:HIS:CE1	3.01	0.48
1:A:72:GLY:O	1:A:76:MET:HG3	2.14	0.48
2:B:50:LYS:HB2	4:B:110:HOH:O	2.14	0.47
3:C:99:ALA:HA	3:C:110:ILE:O	2.15	0.47
3:C:419:HIS:HE1	4:C:628:HOH:O	1.98	0.47
1:A:53:VAL:O	1:A:57:MET:HG3	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:C:408:THR:O	3:C:411:PRO:HD2	2.15	0.47
3:C:447:VAL:HB	3:C:455:ILE:HG22	1.96	0.47
1:A:10:LYS:NZ	4:A:113:HOH:O	2.48	0.46
3:C:177:ILE:HD13	3:C:213:VAL:HG13	1.97	0.46
3:C:305:THR:HG21	3:C:558:LEU:HD11	1.96	0.46
3:C:303:PRO:HB3	3:C:377:VAL:HG22	1.97	0.46
3:C:36:LEU:N	3:C:36:LEU:HD12	2.31	0.45
3:C:246:HIS:CG	3:C:272[A]:HIS:HE1	2.34	0.45
3:C:347:VAL:HG11	3:C:550:ILE:HD12	1.99	0.45
2:B:100:PRO:O	2:B:101:LEU:CB	2.65	0.45
2:B:72:GLU:HB2	2:B:75:GLN:HE21	1.81	0.44
3:C:349:HIS:HD2	3:C:406:LYS:NZ	2.15	0.44
3:C:299:ASN:N	3:C:300:PRO:HD2	2.33	0.43
3:C:75:ILE:O	3:C:81:ILE:HA	2.17	0.43
3:C:65:CYS:HA	3:C:111:PRO:HG2	1.99	0.43
3:C:246:HIS:NE2	3:C:278:GLY:HA3	2.33	0.43
3:C:282:PRO:HG3	3:C:541:TYR:CE1	2.54	0.43
3:C:218:ILE:HD13	3:C:243:VAL:CG1	2.48	0.43
3:C:273:THR:O	3:C:285:ILE:HA	2.20	0.42
3:C:281:ALA:HA	3:C:282:PRO:HA	1.85	0.41
3:C:199:VAL:HG23	3:C:205:LEU:HD21	2.01	0.41
3:C:526:VAL:HG12	3:C:527:HIS:CD2	2.55	0.41
1:A:86:PHE:CE1	1:A:91:LYS:HB2	2.56	0.41
3:C:304:TYR:O	3:C:554:PRO:HA	2.21	0.41
3:C:474:ARG:HA	3:C:475:PRO:HD3	1.92	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
2	B	99/101 (98%)	94 (95%)	5 (5%)	0	100	100
3	C	533/566 (94%)	496 (93%)	34 (6%)	3 (1%)	33	32

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	730/767 (95%)	687 (94%)	40 (6%)	3 (0%)	43 45

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	364	MET
3	C	481	GLY
3	C	282	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	85/85 (100%)	84 (99%)	1 (1%)	82 90
2	B	78/78 (100%)	77 (99%)	1 (1%)	80 89
3	C	417/443 (94%)	404 (97%)	13 (3%)	52 63
All	All	580/606 (96%)	565 (97%)	15 (3%)	61 70

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	ILE
2	B	97	VAL
3	C	179	ARG
3	C	199	VAL
3	C	272[A]	HIS
3	C	272[B]	HIS
3	C	282	PRO
3	C	349	HIS
3	C	392	THR
3	C	394	ASP
3	C	395	ASN
3	C	449	LYS
3	C	507	LEU
3	C	508	ARG
3	C	520	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	GLN
2	B	12	GLN
2	B	16	ASN
2	B	75	GLN
3	C	3	ASN
3	C	142	GLN
3	C	166	HIS
3	C	201	GLN
3	C	349	HIS
3	C	362	GLN
3	C	395	ASN
3	C	419	HIS
3	C	469	GLN
3	C	486	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/100 (100%)	-0.70	0 100 100	3, 11, 30, 38	0
2	B	101/101 (100%)	-0.40	0 100 100	7, 18, 37, 46	0
3	C	536/566 (94%)	-0.56	17 (3%) 45 46	2, 10, 48, 85	0
All	All	737/767 (96%)	-0.56	17 (2%) 57 58	2, 11, 41, 85	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	302	LEU	5.6
3	C	277	GLY	5.4
3	C	307	ASN	5.3
3	C	306	LEU	4.5
3	C	304	TYR	4.4
3	C	300	PRO	4.2
3	C	280	HIS	3.8
3	C	279	GLY	3.7
3	C	538	ALA	3.1
3	C	541	TYR	3.1
3	C	305	THR	3.0
3	C	299	ASN	3.0
3	C	303	PRO	2.8
3	C	339	ARG	2.7
3	C	278	GLY	2.6
3	C	301	THR	2.6
3	C	547	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.