



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 08:00 PM GMT

PDB ID : 2A54
Title : fluorescent protein asFP595, A143S, on-state, 1min irradiation
Authors : Andresen, M.; Wahl, M.C.; Stiel, A.C.; Graeter, F.; Schaefer, L.; Trowitzsch, S.; Weber, G.; Eggeling, C.; Grubmueller, H.; Hell, S.W.; Jakobs, S.
Deposited on : 2005-06-30
Resolution : 1.45 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2476 (1.50-1.42)
Clashscore	79885	2834 (1.50-1.42)
Ramachandran outliers	78287	2769 (1.50-1.42)
Sidechain outliers	78261	2767 (1.50-1.42)
RSRZ outliers	66119	2477 (1.50-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	73	
1	C	73	
2	B	168	
2	D	168	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4313 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GFP-like non-fluorescent chromoprotein FP595 chain 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	64	Total	C	N	O	S	0	2	0
			501	320	81	96	4			
1	C	59	Total	C	N	O	S	2	0	0
			456	295	72	85	4			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	EXPRESSION TAG	UNP Q9GZ28
A	-9	ARG	-	EXPRESSION TAG	UNP Q9GZ28
A	-8	GLY	-	EXPRESSION TAG	UNP Q9GZ28
A	-7	SER	-	EXPRESSION TAG	UNP Q9GZ28
A	-6	HIS	-	EXPRESSION TAG	UNP Q9GZ28
A	-5	HIS	-	EXPRESSION TAG	UNP Q9GZ28
A	-4	HIS	-	EXPRESSION TAG	UNP Q9GZ28
A	-3	HIS	-	EXPRESSION TAG	UNP Q9GZ28
A	-2	HIS	-	EXPRESSION TAG	UNP Q9GZ28
A	-1	HIS	-	EXPRESSION TAG	UNP Q9GZ28
A	0	GLY	-	EXPRESSION TAG	UNP Q9GZ28
A	1	SER	-	EXPRESSION TAG	UNP Q9GZ28
C	-10	MET	-	EXPRESSION TAG	UNP Q9GZ28
C	-9	ARG	-	EXPRESSION TAG	UNP Q9GZ28
C	-8	GLY	-	EXPRESSION TAG	UNP Q9GZ28
C	-7	SER	-	EXPRESSION TAG	UNP Q9GZ28
C	-6	HIS	-	EXPRESSION TAG	UNP Q9GZ28
C	-5	HIS	-	EXPRESSION TAG	UNP Q9GZ28
C	-4	HIS	-	EXPRESSION TAG	UNP Q9GZ28
C	-3	HIS	-	EXPRESSION TAG	UNP Q9GZ28
C	-2	HIS	-	EXPRESSION TAG	UNP Q9GZ28
C	-1	HIS	-	EXPRESSION TAG	UNP Q9GZ28
C	0	GLY	-	EXPRESSION TAG	UNP Q9GZ28
C	1	SER	-	EXPRESSION TAG	UNP Q9GZ28

- Molecule 2 is a protein called GFP-like non-fluorescent chromoprotein FP595 chain 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	168	Total	C	N	O	S	9	4	0
			1392	885	237	259	11			
2	D	168	Total	C	N	O	S	6	3	0
			1389	885	235	258	11			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	65	NRQ	MET	CHROMOPHORE	UNP Q9GZ28
B	65	NRQ	TYR	CHROMOPHORE	UNP Q9GZ28
B	65	NRQ	GLY	CHROMOPHORE	UNP Q9GZ28
B	143	SER	ALA	ENGINEERED	UNP Q9GZ28
D	65	NRQ	MET	CHROMOPHORE	UNP Q9GZ28
D	65	NRQ	TYR	CHROMOPHORE	UNP Q9GZ28
D	65	NRQ	GLY	CHROMOPHORE	UNP Q9GZ28
D	143	SER	ALA	ENGINEERED	UNP Q9GZ28

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	94	Total	O	0	0
			94	94		
4	B	209	Total	O	0	0
			209	209		
4	C	62	Total	O	0	0
			62	62		
4	D	209	Total	O	0	0
			209	209		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GFP-like non-fluorescent chromoprotein FP595 chain 1

Chain A: 



- Molecule 1: GFP-like non-fluorescent chromoprotein FP595 chain 1

Chain C: 



- Molecule 2: GFP-like non-fluorescent chromoprotein FP595 chain 2

Chain B: 



- Molecule 2: GFP-like non-fluorescent chromoprotein FP595 chain 2

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	76.54Å 125.95Å 93.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.45 65.41 – 1.45	Depositor EDS
% Data completeness (in resolution range)	96.1 (20.00-1.45) 96.1 (65.41-1.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.176 , 0.200 0.175 , 0.200	Depositor DCC
R_{free} test set	3882 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	16.9	Xtriage
Anisotropy	0.294	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.2	EDS
Estimated twinning fraction	0.012 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.017 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 80415 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4313	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NRQ, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/514	0.63	0/690
1	C	1.09	1/468 (0.2%)	0.69	1/627 (0.2%)
2	B	0.67	2/1382 (0.1%)	0.81	2/1863 (0.1%)
2	D	0.78	3/1380 (0.2%)	0.71	2/1862 (0.1%)
All	All	0.77	6/3744 (0.2%)	0.74	5/5042 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	29	LYS	CD-CE	19.85	2.00	1.51
2	D	209	LYS	CG-CD	14.23	2.00	1.52
2	D	212	LYS	CG-CD	9.29	1.84	1.52
2	B	207	LYS	CB-CG	8.31	1.75	1.52
2	B	203	GLU	CA-CB	5.41	1.65	1.53
2	D	91	GLU	CD-OE1	5.23	1.31	1.25

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	203	GLU	N-CA-CB	11.67	131.61	110.60
1	C	29	LYS	CD-CE-NZ	-7.65	94.11	111.70
2	B	162	LEU	CA-CB-CG	6.33	129.86	115.30
2	D	212	LYS	CG-CD-CE	5.67	128.92	111.90
2	D	209	LYS	CB-CG-CD	-5.14	98.24	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	501	0	489	13	0
1	C	456	0	449	6	0
2	B	1392	0	1328	33	0
2	D	1389	0	1320	22	0
3	B	1	0	0	0	0
4	A	94	0	0	3	0
4	B	209	0	0	4	0
4	C	62	0	0	1	0
4	D	209	0	0	10	0
All	All	4313	0	3586	65	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (65) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:207:LYS:HB3	2:B:208:GLY:CA	1.73	1.16
2:B:207:LYS:CB	2:B:208:GLY:HA3	1.69	1.16
2:B:95[A]:THR:HG21	4:B:701:HOH:O	1.63	0.97
2:D:155:ARG:HD2	4:D:328:HOH:O	1.73	0.88
2:D:155:ARG:CD	4:D:328:HOH:O	2.28	0.82
1:A:13[A]:THR:HG21	1:A:62:CYS:O	1.84	0.77
1:A:62:CYS:OXT	2:B:65[B]:NRQ:N1	2.22	0.72
2:B:207:LYS:HB3	2:B:208:GLY:HA3	0.80	0.70
1:C:62:CYS:OXT	2:D:65[B]:NRQ:N1	2.26	0.68
2:B:141:GLU:OE2	2:B:169:HIS:HE1	1.77	0.67
2:D:208:GLY:HA2	2:D:211[B]:TYR:OH	1.95	0.66
2:D:163:LYS:HG2	2:D:169:HIS:CE1	2.34	0.63
2:B:116:VAL:HG12	2:B:118:LYS:HE2	1.80	0.63
2:D:65[A]:NRQ:HB12	2:D:215:GLU:OE1	1.99	0.62
1:A:29:LYS:HG3	4:A:114:HOH:O	2.02	0.60
2:B:200:GLU:OE1	2:B:202:MET:HE1	2.05	0.57
1:C:22:HIS:HE1	1:C:48:GLY:O	1.86	0.56
2:B:169:HIS:HD2	4:B:699:HOH:O	1.89	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:163:LYS:HE3	4:D:323:HOH:O	2.08	0.54
2:B:207:LYS:CB	2:B:208:GLY:CA	2.52	0.53
1:A:58:LEU:HD22	2:B:121:ILE:HD13	1.91	0.52
2:D:95[A]:THR:HG23	4:D:236:HOH:O	2.10	0.51
1:A:18[B]:THR:HG21	4:A:87:HOH:O	2.11	0.51
2:B:175:HIS:HE1	4:B:701:HOH:O	1.93	0.51
1:A:9:MET:HE3	1:A:34:PRO:HG3	1.93	0.51
1:A:41:MET:HG2	2:B:65[B]:NRQ:HE2A	1.93	0.50
2:B:95[A]:THR:CG2	2:B:173:HIS:HB3	2.41	0.50
2:D:175:HIS:HE1	4:D:320:HOH:O	1.93	0.50
1:A:33:ASN:C	1:A:33:ASN:HD22	2.14	0.50
2:D:65[B]:NRQ:O2	2:D:65[B]:NRQ:HD2	2.11	0.49
2:D:169:HIS:CE1	4:D:323:HOH:O	2.65	0.49
1:A:22:HIS:HE1	1:A:48:GLY:O	1.96	0.49
1:C:59:SER:O	2:D:65[A]:NRQ:HB11	2.14	0.48
2:B:137:ALA:O	2:B:162:LEU:HG	2.14	0.48
2:B:65[A]:NRQ:HB12	2:B:215:GLU:OE1	2.13	0.47
2:D:168:ARG:HD2	4:D:360:HOH:O	2.14	0.47
1:A:59:SER:O	2:B:65[A]:NRQ:HB11	2.14	0.47
2:D:197:HIS:CE1	4:D:441:HOH:O	2.66	0.47
2:D:231:HIS:HD2	4:D:340:HOH:O	1.97	0.47
2:D:65[A]:NRQ:N1	2:D:65[A]:NRQ:SD	2.88	0.46
2:B:198:ARG:O	2:B:215:GLU:HA	2.16	0.46
2:B:176:THR:HG21	2:B:178:TYR:CZ	2.51	0.46
1:A:33:ASN:ND2	1:A:36:GLU:H	2.14	0.46
1:C:11:PHE:HB3	2:D:115:LEU:HB2	1.97	0.46
2:B:107:ASP:OD1	2:B:118:LYS:NZ	2.45	0.45
1:A:41:MET:HG2	2:B:65[B]:NRQ:CE	2.47	0.45
2:B:162:LEU:O	2:B:169:HIS:HA	2.17	0.44
2:B:95[B]:THR:HG21	4:D:369:HOH:O	2.17	0.44
2:B:202:MET:HE2	2:B:214:TYR:HB3	1.98	0.44
2:B:134:GLN:HB3	2:B:136:LYS:HD3	1.99	0.43
2:B:65[B]:NRQ:O2	2:B:65[B]:NRQ:HD2	2.18	0.43
1:C:22:HIS:CE1	1:C:48:GLY:O	2.69	0.43
1:C:29:LYS:HG3	4:C:125:HOH:O	2.18	0.43
2:B:84:PHE:HB3	2:B:85:PRO:HA	2.01	0.43
1:A:-1:HIS:HD2	4:A:123:HOH:O	2.01	0.42
2:B:124:ASN:HB2	2:D:91:GLU:HG3	2.01	0.42
2:D:65[B]:NRQ:HB2	2:D:199:ILE:HD11	2.01	0.42
2:B:95[A]:THR:HG23	2:B:95[A]:THR:O	2.19	0.42
2:D:67:LYS:HE2	2:D:178:TYR:CZ	2.54	0.42
2:D:168:ARG:HG3	2:D:169:HIS:N	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:105:HIS:HE1	4:B:810:HOH:O	2.02	0.41
2:B:116:VAL:HG12	2:B:118:LYS:CE	2.49	0.41
2:B:79:TYR:CG	2:B:154:LEU:HD22	2.56	0.41
2:B:95[A]:THR:HG23	2:B:173:HIS:HB3	2.02	0.40
2:D:141:GLU:OE2	2:D:163:LYS:HD2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	64/73 (88%)	64 (100%)	0	0	100	100
1	C	57/73 (78%)	56 (98%)	1 (2%)	0	100	100
2	B	168/168 (100%)	161 (96%)	7 (4%)	0	100	100
2	D	167/168 (99%)	159 (95%)	8 (5%)	0	100	100
All	All	456/482 (95%)	440 (96%)	16 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	55/61 (90%)	54 (98%)	1 (2%)	71	37
1	C	50/61 (82%)	50 (100%)	0	100	100
2	B	144/141 (102%)	135 (94%)	9 (6%)	25	3
2	D	143/141 (101%)	142 (99%)	1 (1%)	91	75

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	392/404 (97%)	381 (97%)	11 (3%)	59 19

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
2	B	118	LYS
2	B	154	LEU
2	B	155	ARG
2	B	162	LEU
2	B	202	MET
2	B	203	GLU
2	B	207	LYS
2	B	228[A]	LYS
2	B	228[B]	LYS
2	D	154	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	HIS
1	A	33	ASN
2	B	105	HIS
2	B	135	ASN
2	B	169	HIS
2	B	175	HIS
2	B	197	HIS
2	B	232	ASN
1	C	22	HIS
2	D	135	ASN
2	D	157	GLN
2	D	169	HIS
2	D	175	HIS
2	D	197	HIS
2	D	231	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NRQ	B	65[A]	2	24,24,25	5.76	8 (33%)	30,32,34	3.19	7 (23%)
2	NRQ	B	65[B]	2	24,24,25	6.09	9 (37%)	30,32,34	3.74	9 (30%)
2	NRQ	D	65[A]	2	24,24,25	6.11	9 (37%)	30,32,34	3.40	8 (26%)
2	NRQ	D	65[B]	2	24,24,25	6.12	9 (37%)	30,32,34	3.93	10 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NRQ	B	65[A]	2	-	0/9/31/32	0/2/2/2
2	NRQ	B	65[B]	2	-	0/9/31/32	0/2/2/2
2	NRQ	D	65[A]	2	-	0/9/31/32	0/2/2/2
2	NRQ	D	65[B]	2	-	0/9/31/32	0/2/2/2

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	65[B]	NRQ	CB2-CA2	21.67	1.50	1.35
2	B	65[B]	NRQ	CB2-CA2	21.51	1.49	1.35
2	D	65[A]	NRQ	CB2-CA2	20.61	1.49	1.35
2	B	65[A]	NRQ	CB2-CA2	19.12	1.48	1.35
2	D	65[A]	NRQ	O3-C3	17.51	1.23	1.11
2	B	65[B]	NRQ	O3-C3	17.19	1.23	1.11
2	D	65[B]	NRQ	O3-C3	16.73	1.22	1.11
2	B	65[A]	NRQ	O3-C3	16.01	1.22	1.11
2	B	65[A]	NRQ	CA2-C2	-10.70	1.36	1.48
2	D	65[A]	NRQ	CA2-C2	-10.12	1.37	1.48
2	D	65[B]	NRQ	CA2-C2	-9.15	1.38	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	65[B]	NRQ	CA2-C2	-8.64	1.39	1.48
2	D	65[A]	NRQ	OH-CZ	-4.27	1.26	1.37
2	D	65[B]	NRQ	OH-CZ	-4.22	1.26	1.37
2	B	65[A]	NRQ	OH-CZ	-4.14	1.26	1.37
2	B	65[B]	NRQ	OH-CZ	-4.02	1.26	1.37
2	B	65[A]	NRQ	C2-N3	-3.82	1.31	1.39
2	D	65[B]	NRQ	CA3-C3	3.49	1.52	1.48
2	D	65[A]	NRQ	C2-N3	-3.32	1.32	1.39
2	B	65[B]	NRQ	C2-N3	-3.02	1.33	1.39
2	D	65[B]	NRQ	C2-N3	-3.02	1.33	1.39
2	B	65[B]	NRQ	CA3-C3	2.87	1.52	1.48
2	D	65[A]	NRQ	O2-C2	2.80	1.29	1.23
2	B	65[A]	NRQ	O2-C2	2.76	1.29	1.23
2	D	65[A]	NRQ	CA3-C3	2.55	1.51	1.48
2	B	65[A]	NRQ	C1-N2	2.51	1.38	1.33
2	D	65[A]	NRQ	CA1-N1	2.46	1.33	1.27
2	D	65[B]	NRQ	CA2-N2	-2.37	1.33	1.38
2	B	65[B]	NRQ	CA1-N1	2.36	1.33	1.27
2	B	65[A]	NRQ	CA1-N1	2.27	1.33	1.27
2	D	65[B]	NRQ	O2-C2	2.23	1.28	1.23
2	B	65[B]	NRQ	CA2-N2	-2.23	1.33	1.38
2	D	65[B]	NRQ	CA1-N1	2.21	1.33	1.27
2	B	65[B]	NRQ	O2-C2	2.14	1.27	1.23
2	D	65[A]	NRQ	C1-N2	2.13	1.37	1.33

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	65[B]	NRQ	CA2-C2-N3	13.72	111.26	103.44
2	D	65[B]	NRQ	CA2-C2-N3	13.67	111.23	103.44
2	D	65[A]	NRQ	CA2-C2-N3	13.04	110.87	103.44
2	B	65[A]	NRQ	CA2-C2-N3	12.82	110.74	103.44
2	D	65[B]	NRQ	CB2-CA2-C2	9.66	137.32	122.46
2	D	65[A]	NRQ	O2-C2-CA2	-9.41	125.52	130.96
2	B	65[B]	NRQ	CB2-CA2-C2	9.14	136.51	122.46
2	B	65[A]	NRQ	O2-C2-CA2	-8.29	126.16	130.96
2	D	65[B]	NRQ	O2-C2-CA2	-7.70	126.50	130.96
2	D	65[B]	NRQ	CB2-CA2-N2	-6.73	116.16	128.59
2	B	65[B]	NRQ	CB2-CA2-N2	-5.97	117.56	128.59
2	B	65[B]	NRQ	O2-C2-CA2	-5.80	127.61	130.96
2	B	65[A]	NRQ	CA3-N3-C1	5.04	131.61	124.10
2	D	65[B]	NRQ	CA3-N3-C1	4.41	130.67	124.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	65[B]	NRQ	CA3-N3-C1	4.28	130.47	124.10
2	D	65[A]	NRQ	N3-C1-N2	-4.16	109.99	113.24
2	B	65[B]	NRQ	C2-CA2-N2	-3.99	105.91	108.91
2	D	65[A]	NRQ	CA3-N3-C1	3.96	129.99	124.10
2	D	65[B]	NRQ	CA3-N3-C2	3.28	125.50	123.46
2	B	65[B]	NRQ	N3-C1-N2	-3.23	110.72	113.24
2	D	65[B]	NRQ	C2-CA2-N2	-3.21	106.50	108.91
2	D	65[B]	NRQ	C2-N3-C1	-3.10	103.74	109.01
2	D	65[B]	NRQ	N3-C1-N2	-2.80	111.06	113.24
2	B	65[B]	NRQ	C2-N3-C1	-2.69	104.43	109.01
2	D	65[A]	NRQ	CB2-CA2-C2	2.64	126.53	122.46
2	D	65[A]	NRQ	CA1-C1-N2	2.57	126.70	122.17
2	B	65[B]	NRQ	CA3-N3-C2	2.56	125.06	123.46
2	B	65[A]	NRQ	N3-C1-N2	-2.52	111.27	113.24
2	D	65[A]	NRQ	CE2-CD2-CG2	-2.41	118.26	121.30
2	B	65[A]	NRQ	CA1-C1-N2	2.32	126.27	122.17
2	B	65[A]	NRQ	C2-N3-C1	-2.31	105.08	109.01
2	B	65[A]	NRQ	CD1-CG2-CD2	2.17	121.03	117.66
2	D	65[A]	NRQ	C2-N3-C1	-2.17	105.32	109.01
2	D	65[B]	NRQ	CG1-CB1-CA1	-2.05	108.69	112.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	64/73 (87%)	-0.23	0 100 100	14, 18, 24, 25	0
1	C	59/73 (80%)	-0.12	1 (1%) 67 73	16, 19, 24, 32	1 (1%)
2	B	168/168 (100%)	-0.22	3 (1%) 65 72	13, 16, 30, 48	2 (1%)
2	D	168/168 (100%)	-0.25	0 100 100	14, 18, 30, 37	2 (1%)
All	All	459/482 (95%)	-0.22	4 (0%) 81 87	13, 18, 29, 48	5 (1%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	4	PHE	4.6
2	B	202	MET	3.6
2	B	208	GLY	3.0
2	B	206	GLU	2.5

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NRQ	B	65[B]	23/24	0.10	2.15	15,17,18,21	23
2	NRQ	B	65[A]	23/24	0.10	2.03	13,15,22,26	23
2	NRQ	D	65[B]	23/24	0.09	0.79	14,16,19,21	23
2	NRQ	D	65[A]	23/24	0.09	0.76	15,17,21,25	23

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	CL	B	601	1/1	0.04	-2.37	20,20,20,20	0

6.5 Other polymers

There are no such residues in this entry.