



# wwPDB X-ray Structure Validation Summary Report i

Feb 26, 2014 – 03:31 PM GMT

PDB ID : 2A5H  
Title : 2.1 Angstrom X-ray crystal structure of lysine-2,3-aminomutase from Clostridium subterminale SB4, with Michaelis analog (L-alpha-lysine external aldimine form of pyridoxal-5'-phosphate).  
Authors : Lepore, B.W.; Ruzicka, F.J.; Frey, P.A.; Ringe, D.  
Deposited on : 2005-06-30  
Resolution : 2.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	416	
1	B	416	
1	C	416	
1	D	416	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	SO4	A	592	-	X
3	SO4	B	495	-	X
3	SO4	C	593	-	X
3	SO4	D	494	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Geometry	Electron density
4	SAM	C	417	-	X
5	LYS	A	420[A]	X	-
5	LYS	A	420[B]	X	-
5	LYS	B	420[A]	X	-
5	LYS	B	420[B]	X	-
5	LYS	C	420[A]	X	-
5	LYS	C	420[B]	X	-
5	LYS	D	420[A]	X	-
5	LYS	D	420[B]	X	-

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14034 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-lysine 2,3-aminomutase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	409	Total	C	N	O	S	Se	28	9	0
			3285	2067	591	607	11	9			
1	B	410	Total	C	N	O	S	Se	21	8	0
			3288	2071	589	608	11	9			
1	C	409	Total	C	N	O	S	Se	23	8	0
			3280	2065	588	607	11	9			
1	D	410	Total	C	N	O	S	Se	17	9	0
			3297	2074	595	608	11	9			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	GB 5410603
A	57	MSE	MET	MODIFIED RESIDUE	GB 5410603
A	124	MSE	MET	MODIFIED RESIDUE	GB 5410603
A	127	MSE	MET	MODIFIED RESIDUE	GB 5410603
A	145	MSE	MET	MODIFIED RESIDUE	GB 5410603
A	147	MSE	MET	MODIFIED RESIDUE	GB 5410603
A	218	MSE	MET	MODIFIED RESIDUE	GB 5410603
A	272	MSE	MET	MODIFIED RESIDUE	GB 5410603
A	341	MSE	MET	MODIFIED RESIDUE	GB 5410603
A	400	MSE	MET	MODIFIED RESIDUE	GB 5410603
B	1	MSE	MET	MODIFIED RESIDUE	GB 5410603
B	57	MSE	MET	MODIFIED RESIDUE	GB 5410603
B	124	MSE	MET	MODIFIED RESIDUE	GB 5410603
B	127	MSE	MET	MODIFIED RESIDUE	GB 5410603
B	145	MSE	MET	MODIFIED RESIDUE	GB 5410603
B	147	MSE	MET	MODIFIED RESIDUE	GB 5410603
B	218	MSE	MET	MODIFIED RESIDUE	GB 5410603
B	272	MSE	MET	MODIFIED RESIDUE	GB 5410603
B	341	MSE	MET	MODIFIED RESIDUE	GB 5410603
B	400	MSE	MET	MODIFIED RESIDUE	GB 5410603
C	1	MSE	MET	MODIFIED RESIDUE	GB 5410603

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	57	MSE	MET	MODIFIED RESIDUE	GB 5410603
C	124	MSE	MET	MODIFIED RESIDUE	GB 5410603
C	127	MSE	MET	MODIFIED RESIDUE	GB 5410603
C	145	MSE	MET	MODIFIED RESIDUE	GB 5410603
C	147	MSE	MET	MODIFIED RESIDUE	GB 5410603
C	218	MSE	MET	MODIFIED RESIDUE	GB 5410603
C	272	MSE	MET	MODIFIED RESIDUE	GB 5410603
C	341	MSE	MET	MODIFIED RESIDUE	GB 5410603
C	400	MSE	MET	MODIFIED RESIDUE	GB 5410603
D	1	MSE	MET	MODIFIED RESIDUE	GB 5410603
D	57	MSE	MET	MODIFIED RESIDUE	GB 5410603
D	124	MSE	MET	MODIFIED RESIDUE	GB 5410603
D	127	MSE	MET	MODIFIED RESIDUE	GB 5410603
D	145	MSE	MET	MODIFIED RESIDUE	GB 5410603
D	147	MSE	MET	MODIFIED RESIDUE	GB 5410603
D	218	MSE	MET	MODIFIED RESIDUE	GB 5410603
D	272	MSE	MET	MODIFIED RESIDUE	GB 5410603
D	341	MSE	MET	MODIFIED RESIDUE	GB 5410603
D	400	MSE	MET	MODIFIED RESIDUE	GB 5410603

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is S-ADENOSYLMETHIONINE (three-letter code: SAM) (formula: C<sub>15</sub>H<sub>22</sub>N<sub>6</sub>O<sub>5</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
4	B	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
4	C	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
4	D	1	Total	C	N	O	S	0	0
			27	15	6	5	1		

- Molecule 5 is LYSINE (three-letter code: PLP, LYS) (formula:  $C_8H_{10}NO_6P$ ,  $C_6H_{15}N_2O_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	2	Total	C	N	O	P	0	1
			28	17	3	7	1		
5	B	2	Total	C	N	O	P	0	1
			28	17	3	7	1		
5	C	2	Total	C	N	O	P	0	1
			28	17	3	7	1		
5	D	2	Total	C	N	O	P	0	1
			28	17	3	7	1		

- Molecule 6 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula:  $Fe_4S_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	Fe	S	0	0
			8	4	4		
6	B	1	Total	Fe	S	0	0
			8	4	4		
6	C	1	Total	Fe	S	0	0
			8	4	4		
6	D	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	135	Total	O	0	0
			135	135		
7	B	116	Total	O	0	0
			116	116		
7	C	183	Total	O	0	0
			183	183		
7	D	174	Total	O	0	0
			174	174		




### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: L-lysine 2,3-aminomutase




- Molecule 1: L-lysine 2,3-aminomutase



- Molecule 1: L-lysine 2,3-aminomutase

Chain C: 

- Molecule 1: L-lysine 2,3-aminomutase

Chain D: 

## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.89Å 92.93Å 177.74Å 90.00° 96.74° 90.00°	Depositor
Resolution (Å)	50.00 – 2.10 46.61 – 3.00	Depositor EDS
% Data completeness (in resolution range)	81.5 (50.00-2.10) 96.8 (46.61-3.00)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.65 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.184 , 0.225 0.204 , 0.245	Depositor DCC
$R_{free}$ test set	8171 reflections (9.41%)	DCC
Wilson B-factor (Å <sup>2</sup> )	50.2	Xtriage
Anisotropy	1.080	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 11.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	2 of 76379 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	14034	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ZN, SF4, SAM, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.47	7/3391 (0.2%)	1.17	35/4590 (0.8%)
1	B	1.50	8/3388 (0.2%)	1.14	25/4586 (0.5%)
1	C	1.05	6/3380 (0.2%)	1.18	34/4576 (0.7%)
1	D	1.19	7/3402 (0.2%)	1.15	35/4604 (0.8%)
All	All	1.32	28/13561 (0.2%)	1.16	129/18356 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	2
1	D	0	2
All	All	0	8

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	408	GLU	CD-OE2	51.67	1.82	1.25
1	A	148	GLU	CG-CD	48.53	2.24	1.51
1	B	408	GLU	CD-OE1	-44.21	0.77	1.25
1	A	33	GLU	CG-CD	-38.65	0.94	1.51
1	A	46	GLU	CG-CD	-32.43	1.03	1.51

The worst 5 of 129 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	408	GLU	CG-CD-OE2	-25.33	67.64	118.30
1	A	33	GLU	CB-CG-CD	19.16	165.93	114.20
1	D	408	GLU	CG-CD-OE1	-16.68	84.94	118.30
1	B	408	GLU	CG-CD-OE1	15.71	149.73	118.30
1	B	263	ARG	NE-CZ-NH2	15.03	127.81	120.30

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	148	GLU	Sidechain
1	A	170	GLY	Peptide
1	B	170	GLY	Peptide
1	B	408	GLU	Sidechain
1	C	170	GLY	Peptide

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3285	0	0	18	0
1	B	3288	0	0	20	0
1	C	3280	0	0	20	0
1	D	3297	0	0	25	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
4	A	27	0	0	0	0
4	B	27	0	0	0	0
4	C	27	0	0	0	0
4	D	27	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	28	0	0	0	0
5	B	28	0	0	1	0
5	C	28	0	0	0	0
5	D	28	0	0	1	0
6	A	8	0	0	0	0
6	B	8	0	0	0	0
6	C	8	0	0	0	0
6	D	8	0	0	0	0
7	A	135	0	0	2	0
7	B	116	0	0	3	0
7	C	183	0	0	5	1
7	D	174	0	0	3	0
All	All	14034	0	0	73	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:285[B]:ARG:CD	1:D:285[B]:ARG:CD	2.09	1.31
1:A:214:GLU:OE1	7:A:713:HOH:O	1.68	1.10
1:C:27[B]:ARG:NH2	1:C:29:GLU:OE2	2.13	0.81
1:C:285[B]:ARG:CD	1:D:285[B]:ARG:NE	2.45	0.79
1:D:243[A]:ARG:CD	7:D:578:HOH:O	2.31	0.78

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
7:C:764:HOH:O	7:C:764:HOH:O[2_454]	1.88	0.32

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	416/416 (100%)	400 (96%)	13 (3%)	3 (1%)	30	23
1	B	416/416 (100%)	398 (96%)	15 (4%)	3 (1%)	30	23
1	C	415/416 (100%)	397 (96%)	15 (4%)	3 (1%)	30	23
1	D	417/416 (100%)	397 (95%)	17 (4%)	3 (1%)	30	23
All	All	1664/1664 (100%)	1592 (96%)	60 (4%)	12 (1%)	30	23

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	171	GLY
1	B	52	VAL
1	B	171	GLY
1	C	171	GLY
1	D	52	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/362 (103%)	316 (84%)	58 (16%)	4	2
1	B	374/362 (103%)	315 (84%)	59 (16%)	4	1
1	C	373/362 (103%)	320 (86%)	53 (14%)	5	2
1	D	375/362 (104%)	318 (85%)	57 (15%)	4	2
All	All	1496/1448 (103%)	1269 (85%)	227 (15%)	5	2

5 of 227 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	296	LEU
1	C	119	LEU
1	D	276	VAL
1	B	316[A]	ARG
1	B	407	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 28 ligands modelled in this entry, 4 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	SAM	A	417	6	26,29,29	2.49	6 (23%)	38,42,42	2.17	11 (28%)
6	SF4	A	418	1,4	12,12,12	25.52	11 (91%)	0,24,24	0.00	-
5	PLP	A	419	5	14,15,16	1.33	1 (7%)	20,22,23	1.27	4 (20%)
5	LYS	A	420[A]	-	2,2,9	7.49	2 (100%)	0,1,10	0.00	-
5	LYS	A	420[B]	-	2,2,9	6.28	2 (100%)	0,1,10	0.00	-
3	SO4	A	592	-	4,4,4	0.18	0	6,6,6	0.19	0
4	SAM	B	417	6	26,29,29	2.34	7 (26%)	38,42,42	2.05	13 (34%)
6	SF4	B	418	1,4	12,12,12	23.20	11 (91%)	0,24,24	0.00	-
5	PLP	B	419	5	14,15,16	1.26	1 (7%)	20,22,23	1.01	0
5	LYS	B	420[A]	-	2,2,9	4.90	2 (100%)	0,1,10	0.00	-
5	LYS	B	420[B]	-	2,2,9	4.72	2 (100%)	0,1,10	0.00	-
3	SO4	B	495	-	4,4,4	0.26	0	6,6,6	0.26	0
4	SAM	C	417	6	26,29,29	2.52	6 (23%)	38,42,42	1.91	10 (26%)
6	SF4	C	418	1,4	12,12,12	22.66	12 (100%)	0,24,24	0.00	-
5	PLP	C	419	5	14,15,16	1.19	1 (7%)	20,22,23	0.98	0



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	LYS	C	420[A]	-	2,2,9	3.97	2 (100%)	0,1,10	0.00	-
5	LYS	C	420[B]	-	2,2,9	4.34	2 (100%)	0,1,10	0.00	-
3	SO4	C	593	-	4,4,4	0.20	0	6,6,6	0.39	0
4	SAM	D	417	6	26,29,29	2.37	5 (19%)	38,42,42	2.28	11 (28%)
6	SF4	D	418	1,4	12,12,12	23.29	12 (100%)	0,24,24	0.00	-
5	PLP	D	419	5	14,15,16	1.36	2 (14%)	20,22,23	1.03	2 (10%)
5	LYS	D	420[A]	-	2,2,9	5.31	2 (100%)	0,1,10	0.00	-
5	LYS	D	420[B]	-	2,2,9	4.90	2 (100%)	0,1,10	0.00	-
3	SO4	D	494	-	4,4,4	0.24	0	6,6,6	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAM	A	417	6	-	0/13/33/33	0/1/3/3
6	SF4	A	418	1,4	-	0/0/48/48	0/0/5/5
5	PLP	A	419	5	-	0/6/6/8	0/1/1/1
5	LYS	A	420[A]	-	-	0/0/0/9	0/0/0/0
5	LYS	A	420[B]	-	-	0/0/0/9	0/0/0/0
3	SO4	A	592	-	-	0/0/0/0	0/0/0/0
4	SAM	B	417	6	-	0/13/33/33	0/1/3/3
6	SF4	B	418	1,4	-	0/0/48/48	0/0/5/5
5	PLP	B	419	5	-	0/6/6/8	0/1/1/1
5	LYS	B	420[A]	-	-	0/0/0/9	0/0/0/0
5	LYS	B	420[B]	-	-	0/0/0/9	0/0/0/0
3	SO4	B	495	-	-	0/0/0/0	0/0/0/0
4	SAM	C	417	6	-	0/13/33/33	0/1/3/3
6	SF4	C	418	1,4	-	0/0/48/48	0/0/5/5
5	PLP	C	419	5	-	0/6/6/8	0/1/1/1
5	LYS	C	420[A]	-	-	0/0/0/9	0/0/0/0
5	LYS	C	420[B]	-	-	0/0/0/9	0/0/0/0
3	SO4	C	593	-	-	0/0/0/0	0/0/0/0
4	SAM	D	417	6	-	0/13/33/33	0/1/3/3
6	SF4	D	418	1,4	-	0/0/48/48	0/0/5/5
5	PLP	D	419	5	-	0/6/6/8	0/1/1/1
5	LYS	D	420[A]	-	-	0/0/0/9	0/0/0/0
5	LYS	D	420[B]	-	-	0/0/0/9	0/0/0/0
3	SO4	D	494	-	-	0/0/0/0	0/0/0/0

The worst 5 of 91 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	418	SF4	S1-FE4	-30.64	2.12	2.33
6	A	418	SF4	S1-FE2	-30.22	2.12	2.33
6	B	418	SF4	S3-FE1	-30.08	2.13	2.33
6	A	418	SF4	S4-FE3	-30.03	2.13	2.33
6	D	418	SF4	S3-FE4	-29.60	2.13	2.33

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	417	SAM	O4'-C1'-N9	7.01	114.96	108.44
4	A	417	SAM	O4'-C1'-N9	6.73	114.70	108.44
4	B	417	SAM	N3-C2-N1	-5.45	124.15	128.71
4	D	417	SAM	N3-C4-N9	5.18	134.79	125.43
4	D	417	SAM	N3-C2-N1	-5.16	124.39	128.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.











## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	409/416 (98%)	1.45	106 (25%)  	18, 39, 61, 72	9 (2%)
1	B	410/416 (98%)	1.73	140 (34%)  	20, 44, 79, 98	7 (1%)
1	C	409/416 (98%)	1.47	107 (26%)  	16, 32, 59, 73	7 (1%)
1	D	410/416 (98%)	1.44	98 (23%)  	15, 33, 55, 69	6 (1%)
All	All	1638/1664 (98%)	1.52	451 (27%)  	15, 36, 65, 98	29 (1%)

The worst 5 of 451 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	38	ILE	8.6
1	A	48	VAL	8.0
1	C	40	LEU	7.5
1	B	85	LEU	7.3
1	C	64	LEU	7.0

### 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	C	593	5/5	0.48	7.35	79,80,81,81	0
3	SO4	B	495	5/5	0.41	3.04	101,101,101,101	0
3	SO4	D	494	5/5	0.51	2.87	88,88,88,89	0
3	SO4	A	592	5/5	0.31	2.68	98,98,99,99	0
4	SAM	C	417	27/27	0.21	2.17	21,23,25,31	1
5	LYS	C	420[B]	3/10	0.18	1.95	23,23,23,23	3
4	SAM	B	417	27/27	0.25	1.66	32,34,39,43	1
5	LYS	C	420[A]	3/10	0.18	1.64	23,23,23,23	3
5	LYS	D	420[A]	3/10	0.21	1.08	24,24,25,25	3
4	SAM	A	417	27/27	0.19	0.94	25,27,31,36	1
5	LYS	D	420[B]	3/10	0.21	0.52	24,24,25,25	3
4	SAM	D	417	27/27	0.19	0.52	21,23,26,33	1
5	LYS	B	420[B]	3/10	0.23	0.22	32,32,33,34	3
5	LYS	B	420[A]	3/10	0.23	0.22	33,33,33,34	3
5	PLP	C	419	15/16	0.16	-0.26	18,24,25,26	0
5	LYS	A	420[A]	3/10	0.18	-0.27	27,27,28,28	3
5	LYS	A	420[B]	3/10	0.18	-0.33	27,27,28,28	3
5	PLP	B	419	15/16	0.20	-0.36	27,34,36,37	0
5	PLP	D	419	15/16	0.18	-0.42	20,27,29,30	0
5	PLP	A	419	15/16	0.18	-0.65	24,31,32,33	0
2	ZN	B	421	1/1	0.09	-1.43	45,45,45,45	0
6	SF4	B	418	8/8	0.08	-1.61	35,38,40,40	0
2	ZN	C	421	1/1	0.10	-1.90	36,36,36,36	0
2	ZN	A	421	1/1	0.07	-1.99	36,36,36,36	0
2	ZN	D	421	1/1	0.09	-2.02	31,31,31,31	0
6	SF4	D	418	8/8	0.08	-2.67	21,22,25,26	0
6	SF4	C	418	8/8	0.07	-2.68	22,24,25,26	0
6	SF4	A	418	8/8	0.08	-2.69	24,27,29,29	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.