



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 08:01 PM GMT

PDB ID : 2A5X
Title : Crystal Structure of a Cross-linked Actin Dimer
Authors : Kudryashov, D.S.; Sawaya, M.R.; Adisetiyo, H.; Norcross, T.; Hegyi, G.;
Reisler, E.; Yeates, T.O.
Deposited on : 2005-07-01
Resolution : 2.49 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **FAILED**
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	375	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	CA	A	376	-	X
5	NSB	A	380	-	X
6	MPD	A	381	-	X

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 2812 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Actin, alpha skeletal muscle.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	342	2671	1698	445	509	19	0	2	0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		

- Molecule 3 is PHOSPHOAMINOPHOSPHONICACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	31	10	6	12	3	0	0

- Molecule 4 is LATRUNCULIN A (three-letter code: LAR) (formula: $C_{22}H_{31}NO_5S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			29	22	1	5	1		

- Molecule 5 is N,N,N-TRIMETHYL-3-SULFOPROPAN-1-AMINIUM (three-letter code: NSB) (formula: $C_6H_{16}NO_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			11	6	1	3	1		

- Molecule 6 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			8	6	2		
6	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	52	Total	O	0	0
			52	52		

3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Actin, alpha skeletal muscle

Chain A:



4 Data and refinement statistics

Xtriage (Phenix) failed to run properly - this section will therefore be incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	207.38Å 54.37Å 36.20Å 90.00° 98.62° 90.00°	Depositor
Resolution (Å)	90.00 – 2.49 51.26 – 2.49	Depositor EDS
% Data completeness (in resolution range)	98.8 (90.00-2.49) 98.8 (51.26-2.49)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.194 , 0.250 0.196 , 0.249	Depositor DCC
R_{free} test set	686 reflections (5.17%)	DCC
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 27.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2812	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, ANP, CA, LAR, NSB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/2733	0.60	0/3708

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2671	0	2629	33	0
2	A	2	0	0	0	0
3	A	31	0	13	1	0
4	A	29	0	31	1	0
5	A	11	0	16	0	0
6	A	16	0	28	1	0
7	A	52	0	0	0	0
All	All	2812	0	2717	34	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

All (34) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:368:SER:HA	1:A:371:HIS:CE1	2.19	0.78
1:A:371:HIS:C	1:A:373:LYS:H	1.89	0.75
1:A:194:THR:HA	1:A:198:TYR:O	1.88	0.74
1:A:37:ARG:NH1	1:A:68:LYS:HG3	2.03	0.73
1:A:156:GLY:O	1:A:181:ALA:HB1	1.87	0.73
1:A:368:SER:O	1:A:370:VAL:N	2.30	0.63
1:A:299:MET:CE	1:A:313:MET:HG3	2.28	0.63
1:A:368:SER:C	1:A:370:VAL:H	2.03	0.62
1:A:31:PHE:HB2	1:A:32:PRO:HD2	1.84	0.58
1:A:368:SER:C	1:A:370:VAL:N	2.58	0.56
1:A:196:ARG:O	1:A:196:ARG:HG2	2.07	0.55
1:A:371:HIS:C	1:A:373:LYS:N	2.57	0.53
1:A:299:MET:HE1	1:A:313:MET:HG3	1.90	0.53
1:A:367:PRO:O	1:A:370:VAL:HG12	2.09	0.51
1:A:31:PHE:CE1	1:A:93:GLU:HG3	2.45	0.51
1:A:37:ARG:HH11	1:A:68:LYS:HG3	1.75	0.51
1:A:157:ASP:HB2	3:A:378:ANP:H5'1	1.93	0.51
1:A:368:SER:O	1:A:371:HIS:ND1	2.44	0.50
1:A:335:ARG:HA	1:A:338:SER:OG	2.12	0.49
1:A:81:ASP:HA	1:A:84:LYS:HD2	1.94	0.49
1:A:153:LEU:HD13	1:A:162:ASN:ND2	2.27	0.49
1:A:313:MET:O	1:A:317:ILE:HG12	2.16	0.46
4:A:379:LAR:O2	4:A:379:LAR:H42	2.16	0.46
1:A:131:ALA:HA	1:A:357:ILE:O	2.15	0.45
1:A:368:SER:CA	1:A:371:HIS:CE1	2.94	0.45
1:A:135:ALA:HB1	6:A:382:MPD:H51	1.99	0.45
1:A:153:LEU:HD13	1:A:162:ASN:HD21	1.84	0.43
1:A:107:GLU:HG2	1:A:134:VAL:HG12	2.01	0.43
1:A:106:THR:HB	1:A:137:GLN:HG3	2.01	0.42
1:A:275:HIS:CE1	1:A:276:GLU:HG3	2.55	0.42
1:A:171:LEU:HA	1:A:172:PRO:HD3	1.95	0.41
1:A:372:ARG:O	1:A:373:LYS:HG2	2.20	0.41
1:A:368:SER:HA	1:A:371:HIS:HE1	1.79	0.41
1:A:124:PHE:CZ	1:A:132:MET:HG3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	339/375 (90%)	325 (96%)	12 (4%)	2 (1%)	33 55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	372	ARG
1	A	369	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	289/318 (91%)	279 (96%)	10 (4%)	48 74

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	12	ASN
1	A	33	SER
1	A	195	GLU
1	A	250	ILE
1	A	291	LYS
1	A	323	SER
1	A	351	THR
1	A	360	GLN
1	A	364	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	275	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ANP	A	378	2	33,33,33	2.42	6 (18%)	51,52,52	2.40	13 (25%)
4	LAR	A	379	-	31,31,31	1.52	3 (9%)	43,43,43	2.06	17 (39%)
5	NSB	A	380	-	10,10,10	2.27	1 (10%)	15,15,15	1.84	4 (26%)
6	MPD	A	381	-	7,7,7	0.33	0	10,10,10	0.41	0
6	MPD	A	382	-	7,7,7	0.35	0	10,10,10	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	A	378	2	-	1/18/38/38	0/1/3/3
4	LAR	A	379	-	-	2/28/51/51	0/1/3/3
5	NSB	A	380	-	-	0/8/8/8	0/0/0/0
6	MPD	A	381	-	-	0/5/5/5	0/0/0/0
6	MPD	A	382	-	-	0/5/5/5	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	378	ANP	PB-N3B	8.63	1.71	1.64
3	A	378	ANP	PG-N3B	7.11	1.70	1.64
5	A	380	NSB	C6-S8	-6.73	1.67	1.77
4	A	379	LAR	O2-C1	5.38	1.46	1.34
4	A	379	LAR	C20-S1	-4.09	1.67	1.78
3	A	378	ANP	PG-O1G	3.89	1.51	1.46
3	A	378	ANP	PB-O1B	3.74	1.50	1.46
4	A	379	LAR	C8-C7	-3.59	1.32	1.44
3	A	378	ANP	C5-C4	3.15	1.47	1.40
3	A	378	ANP	C4-N9	-2.55	1.34	1.37

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	378	ANP	O1G-PG-N3B	-8.75	98.61	111.83
3	A	378	ANP	N3-C2-N1	-7.24	122.65	128.71
4	A	379	LAR	C8-C7-C6	5.61	150.15	124.23
3	A	378	ANP	PB-N3B-PG	-5.46	120.88	130.07
3	A	378	ANP	N3-C4-N9	5.12	134.69	125.43
5	A	380	NSB	O46-S8-C6	4.94	111.04	106.81
4	A	379	LAR	O2-C1-C2	4.67	123.41	111.66
4	A	379	LAR	C10-C9-C8	-4.02	112.86	126.70
3	A	378	ANP	O4'-C1'-N9	3.69	111.87	108.44
5	A	380	NSB	O48-S8-C6	3.49	109.80	106.81
4	A	379	LAR	O1-C1-C2	-3.45	117.22	126.28
4	A	379	LAR	S1-C20-N1	3.09	114.30	111.26
4	A	379	LAR	O3-C17-C18	3.08	108.32	104.27
3	A	378	ANP	O2B-PB-O1B	3.01	116.83	109.89
3	A	378	ANP	O3A-PB-N3B	-2.95	98.40	106.59
3	A	378	ANP	C4-C5-N7	-2.94	107.00	109.52
3	A	378	ANP	O4'-C1'-C2'	-2.92	102.30	106.77
3	A	378	ANP	C5-C4-N3	-2.91	119.36	125.70
4	A	379	LAR	O5-C20-N1	-2.76	124.49	126.57
4	A	379	LAR	C16-C17-C18	-2.54	108.67	113.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	379	LAR	C7-C8-C9	2.49	144.58	125.33
3	A	378	ANP	C2-N3-C4	2.47	121.05	114.01
4	A	379	LAR	C14-C15-C16	-2.45	105.88	110.88
4	A	379	LAR	O2-C15-C16	2.39	113.80	107.63
4	A	379	LAR	C4-C5-C6	-2.29	106.80	112.74
3	A	378	ANP	C8-N9-C4	2.28	108.64	106.90
4	A	379	LAR	C21-C3-C4	2.26	118.82	115.39
5	A	380	NSB	C4-C6-S8	-2.21	110.38	113.24
4	A	379	LAR	C18-C19-S1	2.18	109.62	105.93
3	A	378	ANP	C2-N1-C6	2.16	122.66	118.77
4	A	379	LAR	O3-C17-C16	2.13	113.05	110.07
4	A	379	LAR	O2-C1-O1	-2.12	119.38	123.24
5	A	380	NSB	O50-S8-C6	2.05	108.52	105.93
4	A	379	LAR	C17-C16-C15	2.04	115.87	111.16

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	379	LAR	C19-C18-C17-C16
4	A	379	LAR	C7-C8-C9-C10
3	A	378	ANP	O1G-PG-N3B-PB

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	342/375 (91%)	-0.05	10 (2%)	49	51	22, 30, 40, 63	8 (2%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	374	CYS	13.3
1	A	372	ARG	8.2
1	A	373	LYS	6.2
1	A	370	VAL	5.8
1	A	371	HIS	5.5
1	A	169	TYR	5.0
1	A	369	ILE	3.6
1	A	367	PRO	3.5
1	A	360	GLN	2.1
1	A	368	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	MPD	A	381	8/8	0.24	4.61	70,70,71,71	0
2	CA	A	376	1/1	0.16	3.96	27,27,27,27	0
5	NSB	A	380	11/11	0.21	2.61	73,75,79,80	0
4	LAR	A	379	29/29	0.15	0.33	31,34,37,37	0
6	MPD	A	382	8/8	0.19	-0.10	49,52,54,54	0
3	ANP	A	378	31/31	0.12	-0.64	17,22,25,25	0
2	CA	A	377	1/1	0.05	-2.73	44,44,44,44	0

6.5 Other polymers ⓘ

There are no such residues in this entry.