



Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 09:37 PM GMT

PDB ID : 2A8H
Title : Crystal structure of catalytic domain of TACE with Thiomorpholine Sulfonamide Hydroxamate inhibitor
Authors : Levin, J.I.; Chen, J.M.; Laakso, L.M.; Du, M.; Schmid, J.; Xu, W.; Cummons, T.; Xu, J.; Jin, G.; Barone, D.; Skotnicki, J.S.
Deposited on : 2005-07-08
Resolution : 2.30 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	271	
1	B	271	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4295 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADAM 17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	259	Total	C	N	O	S	0	0	0
			2054	1290	347	404	13			
1	B	257	Total	C	N	O	S	0	0	0
			2038	1282	344	399	13			


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	266	ALA	SER	ENGINEERED	UNP P78536
A	452	GLN	ASN	ENGINEERED	UNP P78536
A	478	GLY	-	EXPRESSION TAG	UNP P78536
A	479	SER	-	EXPRESSION TAG	UNP P78536
A	480	HIS	-	EXPRESSION TAG	UNP P78536
A	481	HIS	-	EXPRESSION TAG	UNP P78536
A	482	HIS	-	EXPRESSION TAG	UNP P78536
A	483	HIS	-	EXPRESSION TAG	UNP P78536
A	484	HIS	-	EXPRESSION TAG	UNP P78536
A	485	HIS	-	EXPRESSION TAG	UNP P78536
B	266	ALA	SER	ENGINEERED	UNP P78536
B	452	GLN	ASN	ENGINEERED	UNP P78536
B	478	GLY	-	EXPRESSION TAG	UNP P78536
B	479	SER	-	EXPRESSION TAG	UNP P78536
B	480	HIS	-	EXPRESSION TAG	UNP P78536
B	481	HIS	-	EXPRESSION TAG	UNP P78536
B	482	HIS	-	EXPRESSION TAG	UNP P78536
B	483	HIS	-	EXPRESSION TAG	UNP P78536
B	484	HIS	-	EXPRESSION TAG	UNP P78536
B	485	HIS	-	EXPRESSION TAG	UNP P78536

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 4-({4-[(4-AMINO BUT-2-YNYL)OXY]PHENYL}SULFONYL)-N-HYDROXY-2,2-DIMETHYLTHIOMORPHOLINE-3-CARBOXAMIDE (three-letter code: 4NH) (formula: C₁₇H₂₃N₃O₅S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			27	17	3	5	2		
3	B	1	Total	C	N	O	S	0	0
			27	17	3	5	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	91	Total	O	0	0
			91	91		
4	B	56	Total	O	0	0
			56	56		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ADAM 17

Chain A:



• Molecule 1: ADAM 17

Chain B:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.37Å 59.21Å 198.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 19.80 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.1 (20.00-2.30) 99.1 (19.80-2.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	11.88 (at 2.30Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.215 , 0.281 0.216 , 0.279	Depositor DCC
R_{free} test set	1313 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.421	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 32.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 25906 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4295	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 4NH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/2101	0.67	0/2837
1	B	0.54	0/2085	0.63	0/2814
All	All	0.62	0/4186	0.65	0/5651

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	372	PRO	Peptide

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2054	0	1960	30	0
1	B	2038	0	1951	52	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	27	0	22	1	0
3	B	27	0	22	3	0
4	A	91	0	0	2	0
4	B	56	0	0	3	0
All	All	4295	0	3955	81	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

All (81) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:370:TYR:HA	1:B:371:SER:CB	1.52	1.37
1:B:370:TYR:CA	1:B:371:SER:HB2	1.68	1.21
1:B:371:SER:N	1:B:372:PRO:HD2	1.40	1.17
1:B:371:SER:OG	1:B:376:LYS:O	1.70	1.08
1:B:373:VAL:HG22	1:B:374:GLY:H	1.10	1.06
1:A:369:TYR:CE2	1:A:378:ILE:HG12	1.95	1.02
1:B:371:SER:H	1:B:372:PRO:HD2	1.19	1.01
1:A:378:ILE:HG13	1:A:379:TYR:N	1.77	0.99
1:A:283:ARG:HH11	1:A:283:ARG:HB2	1.31	0.95
1:B:278:GLN:HE22	1:B:474:SER:N	1.64	0.94
1:B:373:VAL:CG2	1:B:374:GLY:N	2.30	0.94
1:B:371:SER:N	1:B:372:PRO:CD	2.30	0.92
1:B:278:GLN:HE22	1:B:474:SER:H	0.94	0.92
1:B:373:VAL:HG22	1:B:374:GLY:N	1.75	0.92
1:B:278:GLN:NE2	1:B:474:SER:H	1.74	0.83
1:B:370:TYR:CA	1:B:371:SER:CB	2.37	0.76
1:A:415:HIS:HE1	3:A:158:4NH:O4	1.66	0.76
1:B:307:GLU:HG2	4:B:66:HOH:O	1.85	0.75
1:B:370:TYR:HA	1:B:371:SER:HB2	0.76	0.74
1:B:269:ASN:HD21	1:B:452:GLN:HE22	1.37	0.72
1:A:297:HIS:HD2	1:A:299:ASN:H	1.39	0.71
1:A:269:ASN:HD21	1:A:452:GLN:HE22	1.43	0.65
1:B:312:TRP:H	1:B:341:GLN:NE2	1.97	0.63
1:A:279:ILE:HG21	1:A:282:ILE:HG13	1.79	0.63
1:A:306:ASN:O	1:A:309:LYS:HG2	1.98	0.62
1:A:290:GLU:HB3	1:B:429:GLN:HG2	1.80	0.62
1:A:283:ARG:HH11	1:A:283:ARG:CB	2.09	0.62
1:A:297:HIS:CD2	1:A:299:ASN:H	2.17	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:428:ASP:N	1:B:428:ASP:OD1	2.27	0.61
1:A:378:ILE:HD11	1:A:380:LEU:CD2	2.32	0.60
1:B:415:HIS:CE1	3:B:159:4NH:O4	2.54	0.60
1:A:378:ILE:HD11	1:A:380:LEU:HD23	1.84	0.59
1:B:306:ASN:HD21	1:B:308:GLU:HG3	1.67	0.59
1:B:375:LYS:O	1:B:376:LYS:CB	2.51	0.58
1:B:361:HIS:HA	4:B:75:HOH:O	2.04	0.58
1:B:361:HIS:HB3	1:B:380:LEU:HD12	1.85	0.58
1:B:304:TYR:CD1	1:B:305:PRO:HA	2.39	0.57
1:B:439:ALA:HB1	3:B:159:4NH:HN21	1.70	0.56
1:A:234:ARG:O	1:A:238:TYR:HD1	1.88	0.56
1:B:409:HIS:CD2	3:B:159:4NH:O5	2.59	0.56
1:B:297:HIS:CD2	1:B:299:ASN:H	2.24	0.56
1:A:369:TYR:HE2	1:A:378:ILE:HG12	1.63	0.55
1:A:260:ASP:OD1	1:A:263:ARG:NH2	2.38	0.54
1:B:226:LYS:HB3	1:B:280:GLU:HB2	1.89	0.54
1:B:375:LYS:O	1:B:376:LYS:HB3	2.09	0.53
1:A:369:TYR:CZ	1:A:378:ILE:HG21	2.44	0.53
1:B:357:ARG:CZ	1:B:360:SER:HB2	2.39	0.53
1:A:436:TYR:CD1	1:A:437:PRO:HD2	2.45	0.52
1:B:370:TYR:HA	1:B:371:SER:HB3	1.75	0.50
1:B:306:ASN:HD22	1:B:306:ASN:C	2.14	0.50
1:A:289:GLN:HG2	1:A:297:HIS:CG	2.47	0.50
1:A:292:LYS:HB3	1:A:293:PRO:HD2	1.94	0.49
1:B:219:ASP:C	1:B:220:PRO:O	2.51	0.49
1:B:269:ASN:HD21	1:B:452:GLN:NE2	2.09	0.49
1:B:230:VAL:HG22	1:B:283:ARG:HB2	1.94	0.48
1:B:370:TYR:C	1:B:372:PRO:HD2	2.25	0.48
1:B:224:THR:HG21	1:B:278:GLN:HE21	1.77	0.48
1:B:255:ILE:HG23	1:B:277:ILE:HD12	1.95	0.47
1:B:220:PRO:C	1:B:222:LYS:H	2.17	0.47
1:B:312:TRP:H	1:B:341:GLN:HE21	1.61	0.46
1:A:348:LEU:HD21	1:A:389:ASN:HB2	1.98	0.46
1:B:330:SER:HB3	1:B:379:TYR:CZ	2.51	0.46
1:B:356:PRO:HD3	4:B:146:HOH:O	2.16	0.45
1:A:217:ASP:HB3	4:A:9:HOH:O	2.15	0.45
1:A:397:LYS:O	1:A:401:LEU:HG	2.17	0.45
1:B:389:ASN:O	1:B:392:LYS:HG3	2.17	0.44
1:B:373:VAL:HG23	1:B:374:GLY:N	2.27	0.44
1:A:416:ASP:HA	1:A:417:PRO:HD3	1.86	0.44
1:A:279:ILE:HG21	1:A:282:ILE:CG1	2.47	0.44
1:B:343:PHE:HB2	1:B:347:THR:HB	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:370:TYR:CG	1:B:371:SER:HB3	2.52	0.43
1:B:304:TYR:CG	1:B:305:PRO:HA	2.53	0.43
1:A:406:GLU:OE1	1:A:406:GLU:HA	2.18	0.43
1:B:371:SER:O	1:B:372:PRO:C	2.58	0.42
1:B:401:LEU:HD11	1:B:442:GLY:CA	2.49	0.42
1:A:306:ASN:HB2	1:A:309:LYS:HE2	2.02	0.42
1:B:352:TYR:HB3	1:B:362:GLY:HA2	2.02	0.42
1:A:369:TYR:O	1:A:378:ILE:HG23	2.20	0.41
1:A:308:GLU:HB2	4:A:21:HOH:O	2.21	0.41
1:B:458:ILE:HG22	1:B:462:ILE:HD12	2.01	0.41
1:A:378:ILE:HD11	1:A:380:LEU:HD21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/271 (95%)	247 (96%)	8 (3%)	2 (1%)	27	30
1	B	255/271 (94%)	240 (94%)	10 (4%)	5 (2%)	11	8
All	All	512/542 (94%)	487 (95%)	18 (4%)	7 (1%)	16	15

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	371	SER
1	B	372	PRO
1	B	376	LYS
1	A	312	TRP
1	A	365	CYS
1	B	365	CYS
1	B	220	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/232 (96%)	211 (95%)	11 (5%)	34	45
1	B	220/232 (95%)	207 (94%)	13 (6%)	28	35
All	All	442/464 (95%)	418 (95%)	24 (5%)	31	40

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	223	ASN
1	A	277	ILE
1	A	283	ARG
1	A	302	LYS
1	A	306	ASN
1	A	308	GLU
1	A	327	GLU
1	A	341	GLN
1	A	378	ILE
1	A	407	LEU
1	A	448	LYS
1	B	223	ASN
1	B	306	ASN
1	B	308	GLU
1	B	309	LYS
1	B	322	SER
1	B	392	LYS
1	B	407	LEU
1	B	414	GLU
1	B	420	LEU
1	B	422	GLU
1	B	426	ASN
1	B	428	ASP
1	B	460	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	223	ASN
1	A	297	HIS
1	A	341	GLN
1	A	452	GLN
1	B	223	ASN
1	B	278	GLN
1	B	297	HIS
1	B	306	ASN
1	B	320	GLN
1	B	341	GLN
1	B	426	ASN
1	B	429	GLN
1	B	452	GLN
1	B	471	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	4NH	A	158	2	28,28,28	2.06	5 (17%)	39,40,40	1.18	5 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	4NH	B	159	2	28,28,28	2.06	5 (17%)	39,40,40	1.18	5 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4NH	A	158	2	-	1/23/41/41	0/2/2/2
3	4NH	B	159	2	-	1/23/41/41	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	159	4NH	S2-N1	8.85	1.76	1.63
3	A	158	4NH	S2-N1	8.82	1.76	1.63
3	A	158	4NH	C3-N1	3.13	1.52	1.48
3	B	159	4NH	C3-N1	3.08	1.52	1.48
3	B	159	4NH	C16-N3	3.08	1.38	1.33
3	A	158	4NH	C16-N3	3.08	1.38	1.33
3	A	158	4NH	C4-N1	2.74	1.52	1.47
3	B	159	4NH	C4-N1	2.69	1.52	1.47
3	B	159	4NH	C2-C3	2.06	1.56	1.53
3	A	158	4NH	C2-C3	2.04	1.56	1.53

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	158	4NH	C6-S2-N1	-3.11	102.24	107.44
3	B	159	4NH	C6-S2-N1	-3.09	102.27	107.44
3	B	159	4NH	O5-N3-C16	-2.98	115.80	119.80
3	A	158	4NH	O5-N3-C16	-2.97	115.81	119.80
3	B	159	4NH	O1-S2-C6	-2.39	104.77	107.94
3	A	158	4NH	O1-S2-C6	-2.38	104.79	107.94
3	B	159	4NH	C17-C2-C3	2.23	114.31	110.64
3	A	158	4NH	C17-C2-C3	2.22	114.31	110.64
3	B	159	4NH	C3-N1-S2	2.15	124.32	119.08
3	A	158	4NH	C3-N1-S2	2.15	124.32	119.08

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	159	4NH	C15-C14-C13-C12
3	A	158	4NH	C15-C14-C13-C12

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/271 (95%)	0.01	6 (2%) 57 67	12, 25, 43, 56	1 (0%)
1	B	257/271 (94%)	0.48	21 (8%) 12 18	16, 36, 59, 72	0
All	All	516/542 (95%)	0.24	27 (5%) 26 36	12, 30, 53, 72	1 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	373	VAL	10.3
1	B	372	PRO	5.0
1	B	369	TYR	4.0
1	B	359	ASN	3.7
1	B	375	LYS	3.5
1	B	370	TYR	3.4
1	B	361	HIS	3.3
1	A	293	PRO	3.1
1	B	270	ALA	3.1
1	B	292	LYS	3.0
1	B	360	SER	2.9
1	A	308	GLU	2.9
1	B	371	SER	2.8
1	A	428	ASP	2.8
1	B	391	GLY	2.7
1	A	302	LYS	2.3
1	B	358	ALA	2.3
1	B	390	TYR	2.2
1	A	238	TYR	2.2
1	B	327	GLU	2.2
1	B	304	TYR	2.2
1	A	291	VAL	2.1
1	B	364	VAL	2.1
1	B	374	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	357	ARG	2.1
1	B	323	PHE	2.0
1	B	281	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	4NH	A	158	27/27	0.16	1.29	23,30,33,34	0
3	4NH	B	159	27/27	0.16	0.60	29,38,39,40	0
2	ZN	B	486	1/1	0.06	-2.12	27,27,27,27	0
2	ZN	A	486	1/1	0.05	-2.38	18,18,18,18	0

6.5 Other polymers

There are no such residues in this entry.