



# Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 10:22 PM GMT

PDB ID : 4A8S  
Title : Non-Catalytic Ions Direct the RNA-Dependent RNA Polymerase of Bacterial dsRNA virus phi6 from De Novo Initiation to Elongation  
Authors : Wright, S.; Poranen, M.M.; Bamford, D.H.; Stuart, D.I.; Grimes, J.M.  
Deposited on : 2011-11-21  
Resolution : 2.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1053 (2.90-2.90)
Clashscore	79885	1326 (2.90-2.90)
Ramachandran outliers	78287	1290 (2.90-2.90)
Sidechain outliers	78261	1292 (2.90-2.90)
RSRZ outliers	66119	1054 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	665	
1	B	665	
1	C	665	
2	F	13	
2	G	13	
2	H	13	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15955 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-DIRECTED RNA POLYMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	664	Total	C	N	O	S	0	0	0
			5265	3342	915	976	32			
1	B	664	Total	C	N	O	S	0	0	0
			5265	3342	915	976	32			
1	C	646	Total	C	N	O	S	0	0	0
			5123	3255	887	949	32			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	456	MET	ILE	CONFLICT	UNP P11124
A	634	GLN	GLU	ENGINEERED MUTATION	UNP P11124
B	456	MET	ILE	CONFLICT	UNP P11124
B	634	GLN	GLU	ENGINEERED MUTATION	UNP P11124
C	456	MET	ILE	CONFLICT	UNP P11124
C	634	GLN	GLU	ENGINEERED MUTATION	UNP P11124


- Molecule 2 is a DNA chain called 5'-D(\*TP\*TP\*TP\*TP\*CP\*GP\*CP\*GP\*TP\*AP\*GP\*C P\*GP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	3	Total	C	N	O	P	0	0	0
			60	29	13	16	2			
2	G	4	Total	C	N	O	P	0	0	0
			81	39	18	21	3			
2	H	6	Total	C	N	O	P	0	0	0
			118	59	16	38	5			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	2	Total	Mg	0	0
			2	2		

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			13	10	3		
4	B	1	Total	O	P	0	0
			13	10	3		
4	C	1	Total	O	P	0	0
			13	10	3		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	O	0	0
			1	1		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: RNA-DIRECTED RNA POLYMERASE

Chain A: 




#### • Molecule 1: RNA-DIRECTED RNA POLYMERASE

Chain B: 



#### • Molecule 1: RNA-DIRECTED RNA POLYMERASE

Chain C: 



- Molecule 2: 5'-D(\*TP\*TP\*TP\*TP\*CP\*GP\*CP\*GP\*TP\*AP\*GP\*CP\*GP)-3'

Chain F: 



- Molecule 2: 5'-D(\*TP\*TP\*TP\*TP\*CP\*GP\*CP\*GP\*TP\*AP\*GP\*CP\*GP)-3'

Chain G: 



- Molecule 2: 5'-D(\*TP\*TP\*TP\*TP\*CP\*GP\*CP\*GP\*TP\*AP\*GP\*CP\*GP)-3'

Chain H: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.51Å 91.71Å 142.08Å 90.00° 102.05° 90.00°	Depositor
Resolution (Å)	33.49 – 2.90 65.57 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.0 (33.49-2.90) 90.1 (65.57-2.90)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.96 (at 2.91Å)	Xtriage
Refinement program	BUSTER 2.9.2	Depositor
R, $R_{free}$	0.216 , 0.250 0.232 , 0.264	Depositor DCC
$R_{free}$ test set	2740 reflections (5.08%)	DCC
Wilson B-factor (Å <sup>2</sup> )	86.5	Xtriage
Anisotropy	0.672	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 53.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 53974 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	15955	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	97.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.45	0/5396	0.64	1/7297 (0.0%)
1	B	0.46	0/5396	0.67	0/7297
1	C	0.46	0/5249	0.68	0/7094
2	F	1.39	0/67	2.48	6/102 (5.9%)
2	G	1.35	0/91	2.24	7/139 (5.0%)
2	H	1.63	1/130 (0.8%)	2.89	16/199 (8.0%)
All	All	0.49	1/16329 (0.0%)	0.75	30/22128 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	5	DC	C1'-N1	5.32	1.56	1.49

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	DT	O4'-C1'-N1	14.61	118.22	108.00
2	H	1	DT	P-O3'-C3'	11.02	132.92	119.70
2	H	3	DT	P-O3'-C3'	11.01	132.91	119.70
2	H	5	DC	O4'-C1'-N1	9.12	114.38	108.00
2	H	3	DT	N3-C2-O2	-8.58	117.15	122.30
2	G	12	DC	O4'-C1'-N1	8.23	113.76	108.00
2	F	6	DG	O4'-C1'-N9	8.19	113.73	108.00
2	H	3	DT	O4'-C1'-N1	8.08	113.66	108.00
2	G	12	DC	P-O3'-C3'	7.57	128.78	119.70
2	H	5	DC	N1-C2-O2	6.73	122.94	118.90
2	F	6	DG	C8-N9-C4	-6.30	103.88	106.40
2	H	5	DC	C2-N1-C1'	6.15	125.56	118.80
2	H	4	DT	C1'-O4'-C4'	-6.13	103.97	110.10
2	H	4	DT	O4'-C1'-N1	5.60	111.92	108.00

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	6	DG	N7-C8-N9	5.50	115.85	113.10
2	H	2	DT	O4'-C1'-N1	5.40	111.78	108.00
2	G	11	DG	C8-N9-C4	-5.36	104.26	106.40
2	G	12	DC	C6-N1-C2	-5.34	118.16	120.30
2	G	11	DG	N7-C8-N9	5.34	115.77	113.10
2	F	7	DC	O4'-C1'-N1	5.30	111.71	108.00
2	G	13	DG	C2-N3-C4	5.25	114.53	111.90
2	H	5	DC	C6-N1-C2	-5.23	118.21	120.30
2	H	6	DG	O4'-C1'-N9	5.18	111.63	108.00
2	F	7	DC	P-O3'-C3'	5.17	125.91	119.70
2	F	7	DC	N1-C2-O2	5.13	121.98	118.90
2	G	11	DG	N3-C4-C5	-5.09	126.06	128.60
1	A	206	GLN	N-CA-C	-5.07	97.31	111.00
2	H	2	DT	N3-C2-O2	-5.05	119.27	122.30
2	H	5	DC	N3-C2-O2	-5.03	118.38	121.90
2	H	5	DC	O4'-C4'-C3'	-5.03	102.49	104.50

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5265	0	0	3	0
1	B	5265	0	0	2	0
1	C	5123	0	0	4	0
2	F	60	0	35	2	0
2	G	81	0	46	1	0
2	H	118	0	72	4	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
4	A	13	0	0	0	0
4	B	13	0	0	0	0
4	C	13	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	1	0	0	0	0
All	All	15955	0	153	12	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

All (12) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:393:SER:CB	2:G:11:DG:H21	2.16	0.59
2:H:3:DT:H2''	2:H:4:DT:H5'	1.88	0.54
1:A:392:SER:O	1:A:398:THR:CG2	2.58	0.51
1:C:86:GLY:O	1:C:89:HIS:CD2	2.66	0.49
1:C:136:SER:OG	1:C:293:LYS:NZ	2.47	0.48
1:B:504:ASP:OD2	1:B:506:ARG:NH1	2.47	0.47
1:C:629:GLN:NE2	2:H:4:DT:C4	2.84	0.46
2:H:4:DT:H2''	2:H:5:DC:O4'	2.16	0.46
1:A:204:ARG:CZ	2:F:7:DC:H41	2.27	0.45
2:H:1:DT:H1'	2:H:2:DT:OP2	2.19	0.43
1:A:204:ARG:NE	2:F:7:DC:H41	2.17	0.41
1:C:477:GLU:N	1:C:477:GLU:CD	2.73	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	662/665 (100%)	642 (97%)	18 (3%)	2 (0%)	50 85
1	B	662/665 (100%)	633 (96%)	25 (4%)	4 (1%)	33 76
1	C	640/665 (96%)	606 (95%)	28 (4%)	6 (1%)	25 66
All	All	1964/1995 (98%)	1881 (96%)	71 (4%)	12 (1%)	33 76

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	136	SER
1	A	607	ARG
1	B	607	ARG
1	B	136	SER
1	C	368	GLY
1	A	136	SER
1	B	368	GLY
1	C	393	SER
1	C	552	ALA
1	B	612	ALA
1	C	545	PRO
1	C	531	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	557/558 (100%)	536 (96%)	21 (4%)	44	83
1	B	557/558 (100%)	538 (97%)	19 (3%)	49	86
1	C	543/558 (97%)	512 (94%)	31 (6%)	29	66
All	All	1657/1674 (99%)	1586 (96%)	71 (4%)	40	78

All (71) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ARG
1	A	30	ARG
1	A	63	ASP
1	A	78	ASN
1	A	81	ARG
1	A	82	THR
1	A	102	LEU
1	A	113	ASP
1	A	163	LYS
1	A	202	VAL
1	A	204	ARG
1	A	269	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	391	LEU
1	A	404	LEU
1	A	452	SER
1	A	497	LEU
1	A	506	ARG
1	A	608	GLN
1	A	629	GLN
1	A	662	MET
1	A	664	ARG
1	B	3	ARG
1	B	30	ARG
1	B	63	ASP
1	B	78	ASN
1	B	82	THR
1	B	102	LEU
1	B	113	ASP
1	B	163	LYS
1	B	202	VAL
1	B	204	ARG
1	B	269	ARG
1	B	403	THR
1	B	404	LEU
1	B	452	SER
1	B	497	LEU
1	B	605	MET
1	B	607	ARG
1	B	647	HIS
1	B	662	MET
1	C	3	ARG
1	C	30	ARG
1	C	63	ASP
1	C	78	ASN
1	C	81	ARG
1	C	82	THR
1	C	102	LEU
1	C	113	ASP
1	C	163	LYS
1	C	202	VAL
1	C	269	ARG
1	C	307	ARG
1	C	324	ASP
1	C	398	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	404	LEU
1	C	452	SER
1	C	472	PHE
1	C	477	GLU
1	C	479	LYS
1	C	513	ILE
1	C	542	ARG
1	C	590	MET
1	C	594	ASP
1	C	597	GLU
1	C	600	ARG
1	C	605	MET
1	C	617	ILE
1	C	622	LEU
1	C	646	MET
1	C	662	MET
1	C	664	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ATP	A	1666	3	12,12,33	1.26	1 (8%)	20,20,52	1.18	1 (5%)
4	ATP	B	1666	-	12,12,33	1.38	1 (8%)	20,20,52	1.08	1 (5%)
4	ATP	C	1665	-	12,12,33	1.24	1 (8%)	20,20,52	1.30	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	1666	3	-	0/12/12/38	0/0/0/3
4	ATP	B	1666	-	-	0/12/12/38	0/0/0/3
4	ATP	C	1665	-	-	0/12/12/38	0/0/0/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1666	ATP	PA-O5'	3.48	1.62	1.50
4	C	1665	ATP	PA-O5'	3.44	1.62	1.50
4	B	1666	ATP	PA-O5'	3.20	1.61	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	1665	ATP	O3A-PB-O3B	4.24	110.28	101.66
4	A	1666	ATP	O3A-PB-O3B	3.63	109.05	101.66
4	B	1666	ATP	O2A-PA-O3A	3.54	115.55	104.29
4	C	1665	ATP	O2A-PA-O3A	2.06	110.85	104.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	664/665 (99%)	0.35	13 (1%)	62	71	59, 83, 107, 136	0
1	B	664/665 (99%)	0.51	46 (6%)	17	20	69, 92, 124, 157	0
1	C	646/665 (97%)	0.74	65 (10%)	7	10	61, 105, 159, 217	0
2	F	3/13 (23%)	0.88	0	100	100	108, 108, 112, 118	0
2	G	4/13 (30%)	1.34	1 (25%)	1	2	160, 166, 170, 181	0
2	H	6/13 (46%)	0.86	0	100	100	121, 131, 141, 142	0
All	All	1987/2034 (97%)	0.53	125 (6%)	19	23	59, 90, 141, 217	0

All (125) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	601	TYR	10.2
1	C	612	ALA	8.7
1	C	541	LYS	8.6
1	B	611	LEU	6.7
1	C	606	ALA	6.7
1	C	546	PHE	6.3
1	C	540	SER	5.9
1	C	602	VAL	5.6
1	C	594	ASP	5.1
1	C	630	TYR	5.1
1	B	661	VAL	4.9
1	C	614	LEU	4.8
1	B	135	PHE	4.8
1	C	567	VAL	4.6
1	C	603	ALA	4.5
1	B	608	GLN	4.5
1	C	539	ARG	4.3
1	C	457	LEU	4.2
1	C	415	LEU	4.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	604	SER	3.8
1	C	422	LEU	3.7
1	C	503	TYR	3.6
1	B	610	GLY	3.6
1	B	607	ARG	3.6
1	C	542	ARG	3.6
1	C	583	TYR	3.6
1	B	356	LEU	3.5
1	C	410	TYR	3.5
1	C	420	PRO	3.4
1	C	404	LEU	3.4
1	C	315	LYS	3.4
1	C	576	TRP	3.4
1	C	435	PHE	3.3
1	C	547	PRO	3.3
1	B	422	LEU	3.3
1	B	142	PRO	3.2
1	C	646	MET	3.2
1	B	400	LEU	3.2
1	C	598	LEU	3.1
1	C	551	TRP	3.1
1	B	634	GLN	3.1
1	C	282	PRO	3.1
1	C	418	THR	3.0
1	C	421	HIS	3.0
1	B	279	LEU	2.9
1	C	471	LEU	2.9
1	B	609	ALA	2.8
1	C	2	ARG	2.8
1	B	341	CYS	2.8
1	C	619	LEU	2.8
1	B	310	LYS	2.7
1	B	127	PHE	2.7
1	B	466	VAL	2.7
1	C	465	LEU	2.7
1	A	567	VAL	2.7
1	C	501	LEU	2.7
1	B	598	LEU	2.6
1	C	143	LEU	2.6
1	C	459	TRP	2.6
1	C	613	GLU	2.6
1	C	101	PRO	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	180	LEU	2.6
1	B	647	HIS	2.6
1	C	548	GLY	2.6
2	G	10	DA	2.6
1	B	363	LEU	2.6
1	C	549	LEU	2.5
1	B	203	TYR	2.5
1	B	249	ALA	2.5
1	C	470	ARG	2.5
1	A	471	LEU	2.5
1	A	664	ARG	2.5
1	B	321	VAL	2.5
1	C	463	ARG	2.5
1	C	75	VAL	2.5
1	A	171	LEU	2.4
1	B	622	LEU	2.4
1	B	658	LEU	2.4
1	B	344	LEU	2.4
1	A	357	PHE	2.4
1	B	637	VAL	2.4
1	C	436	LEU	2.4
1	C	449	ILE	2.4
1	B	614	LEU	2.3
1	C	536	VAL	2.3
1	C	599	SER	2.3
1	C	645	LEU	2.3
1	A	321	VAL	2.3
1	B	285	ALA	2.3
1	B	307	ARG	2.3
1	B	446	ILE	2.2
1	B	468	GLY	2.2
1	A	661	VAL	2.2
1	B	435	PHE	2.2
1	B	649	VAL	2.2
1	B	618	ASP	2.2
1	B	601	TYR	2.2
1	C	234	VAL	2.2
1	C	419	ALA	2.2
1	C	530	TYR	2.2
1	C	595	THR	2.2
1	C	461	LYS	2.2
1	B	657	PHE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	391	LEU	2.1
1	B	348	GLY	2.1
1	A	607	ARG	2.1
1	C	314	VAL	2.1
1	C	1	PRO	2.1
1	A	135	PHE	2.1
1	C	448	GLN	2.1
1	B	314	VAL	2.1
1	B	103	ALA	2.1
1	B	170	ALA	2.1
1	B	606	ALA	2.1
1	C	467	GLY	2.1
1	B	354	VAL	2.1
1	C	408	ILE	2.1
1	C	575	TRP	2.1
1	B	196	GLY	2.1
1	C	522	LEU	2.0
1	A	415	LEU	2.0
1	B	197	GLY	2.0
1	A	2	ARG	2.0
1	B	134	MET	2.0
1	C	600	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	ATP	C	1665	13/31	0.17	-0.91	201,206,211,212	0
4	ATP	A	1666	13/31	0.19	-0.94	180,184,185,186	0
3	MG	A	1665	1/1	0.14	-1.09	72,72,72,72	0
4	ATP	B	1666	13/31	0.17	-1.23	154,159,164,165	0
3	MG	B	1665	1/1	0.14	-1.38	71,71,71,71	0
3	MG	A	1667	1/1	0.11	-2.37	69,69,69,69	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.