



# wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 09:35 PM GMT

PDB ID : 1A9C  
Title : GTP CYCLOHYDROLASE I (C110S MUTANT) IN COMPLEX WITH GTP  
Authors : Auerbach, G.; Nar, H.; Bracher, A.; Bacher, A.; Huber, R.  
Deposited on : 1998-04-04  
Resolution : 2.90 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1326 (2.90-2.90)
Ramachandran outliers	78287	1290 (2.90-2.90)
Sidechain outliers	78261	1292 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	221	
1	B	221	
1	C	221	
1	D	221	
1	E	221	
1	F	221	
1	G	221	
1	H	221	
1	I	221	
1	J	221	
1	K	221	
1	L	221	
1	M	221	
1	N	221	
1	O	221	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 33165 atoms, of which 6495 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GTP CYCLOHYDROLASE I.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	221	Total	C	H	N	O	S	513	0	0
			2137	1088	405	309	327	8			
1	B	221	Total	C	H	N	O	S	510	0	0
			2137	1088	405	309	327	8			
1	C	221	Total	C	H	N	O	S	502	0	0
			2137	1088	405	309	327	8			
1	D	221	Total	C	H	N	O	S	502	0	0
			2137	1088	405	309	327	8			
1	E	221	Total	C	H	N	O	S	519	0	0
			2137	1088	405	309	327	8			
1	F	221	Total	C	H	N	O	S	527	0	0
			2137	1088	405	309	327	8			
1	G	221	Total	C	H	N	O	S	526	0	0
			2137	1088	405	309	327	8			
1	H	221	Total	C	H	N	O	S	513	0	0
			2137	1088	405	309	327	8			
1	I	221	Total	C	H	N	O	S	527	0	0
			2137	1088	405	309	327	8			
1	J	221	Total	C	H	N	O	S	502	0	0
			2137	1088	405	309	327	8			
1	K	221	Total	C	H	N	O	S	514	0	0
			2137	1088	405	309	327	8			
1	L	221	Total	C	H	N	O	S	531	0	0
			2137	1088	405	309	327	8			
1	M	221	Total	C	H	N	O	S	503	0	0
			2137	1088	405	309	327	8			
1	N	221	Total	C	H	N	O	S	519	0	0
			2137	1088	405	309	327	8			
1	O	221	Total	C	H	N	O	S	519	0	0
			2137	1088	405	309	327	8			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	110	SER	CYS	MUTATION	UNP P0A6T5
B	110	SER	CYS	MUTATION	UNP P0A6T5
C	110	SER	CYS	MUTATION	UNP P0A6T5
D	110	SER	CYS	MUTATION	UNP P0A6T5
E	110	SER	CYS	MUTATION	UNP P0A6T5
F	110	SER	CYS	MUTATION	UNP P0A6T5
G	110	SER	CYS	MUTATION	UNP P0A6T5
H	110	SER	CYS	MUTATION	UNP P0A6T5
I	110	SER	CYS	MUTATION	UNP P0A6T5
J	110	SER	CYS	MUTATION	UNP P0A6T5
K	110	SER	CYS	MUTATION	UNP P0A6T5
L	110	SER	CYS	MUTATION	UNP P0A6T5
M	110	SER	CYS	MUTATION	UNP P0A6T5
N	110	SER	CYS	MUTATION	UNP P0A6T5
O	110	SER	CYS	MUTATION	UNP P0A6T5

- Molecule 2 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	B	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	D	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	E	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	F	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	G	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	H	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	I	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	J	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	K	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	L	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	M	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	N	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	O	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	14	Total	H	O	30	0
			42	28	14		
3	B	15	Total	H	O	32	0
			45	30	15		
3	C	14	Total	H	O	30	0
			42	28	14		
3	D	14	Total	H	O	30	0
			42	28	14		
3	E	13	Total	H	O	28	0
			39	26	13		
3	F	14	Total	H	O	30	0
			42	28	14		
3	G	13	Total	H	O	28	0
			39	26	13		
3	H	14	Total	H	O	30	0
			42	28	14		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	I	13	Total 39	H 26	O 13	28	0
3	J	16	Total 48	H 32	O 16	34	0
3	K	15	Total 45	H 30	O 15	32	0
3	L	13	Total 39	H 26	O 13	28	0
3	M	15	Total 45	H 30	O 15	33	0
3	N	14	Total 42	H 28	O 14	30	0
3	O	13	Total 39	H 26	O 13	28	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: GTP CYCLOHYDROLASE I

Chain A:




#### • Molecule 1: GTP CYCLOHYDROLASE I

Chain B:



#### • Molecule 1: GTP CYCLOHYDROLASE I

Chain C:




#### • Molecule 1: GTP CYCLOHYDROLASE I

Chain D:



#### • Molecule 1: GTP CYCLOHYDROLASE I

Chain E:




- Molecule 1: GTP CYCLOHYDROLASE I

Chain F:



- Molecule 1: GTP CYCLOHYDROLASE I

Chain G:




- Molecule 1: GTP CYCLOHYDROLASE I

Chain H:



- Molecule 1: GTP CYCLOHYDROLASE I

Chain I:



- Molecule 1: GTP CYCLOHYDROLASE I

Chain J:





- Molecule 1: GTP CYCLOHYDROLASE I

Chain K: 




- Molecule 1: GTP CYCLOHYDROLASE I

Chain L: 




- Molecule 1: GTP CYCLOHYDROLASE I

Chain M: 




- Molecule 1: GTP CYCLOHYDROLASE I

Chain N: 



- Molecule 1: GTP CYCLOHYDROLASE I

Chain O: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	313.90Å 226.80Å 131.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.90	Depositor
% Data completeness (in resolution range)	84.4 (8.00-2.90)	Depositor
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, $R_{free}$	0.206 , 0.288	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	33165	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.47	0/1760	0.78	1/2384 (0.0%)
1	B	0.50	0/1760	0.78	1/2384 (0.0%)
1	C	0.51	0/1760	0.79	1/2384 (0.0%)
1	D	0.48	0/1760	0.78	2/2384 (0.1%)
1	E	0.46	0/1760	0.76	1/2384 (0.0%)
1	F	0.44	0/1760	0.74	1/2384 (0.0%)
1	G	0.44	0/1760	0.75	1/2384 (0.0%)
1	H	0.46	0/1760	0.77	2/2384 (0.1%)
1	I	0.43	0/1760	0.75	1/2384 (0.0%)
1	J	0.46	0/1760	0.76	1/2384 (0.0%)
1	K	0.44	0/1760	0.77	1/2384 (0.0%)
1	L	0.44	0/1760	0.75	1/2384 (0.0%)
1	M	0.45	0/1760	0.77	1/2384 (0.0%)
1	N	0.45	0/1760	0.76	1/2384 (0.0%)
1	O	0.45	0/1760	0.76	1/2384 (0.0%)
All	All	0.46	0/26400	0.76	17/35760 (0.0%)

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	K	150	VAL	N-CA-C	-6.25	94.14	111.00
1	C	150	VAL	N-CA-C	-6.16	94.37	111.00
1	B	150	VAL	N-CA-C	-6.10	94.53	111.00
1	L	150	VAL	N-CA-C	-6.04	94.68	111.00
1	J	150	VAL	N-CA-C	-5.95	94.93	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1732	405	1363	18	0
1	B	1732	405	1363	20	0
1	C	1732	405	1363	23	0
1	D	1732	405	1363	18	0
1	E	1732	405	1363	19	0
1	F	1732	405	1363	16	0
1	G	1732	405	1363	17	0
1	H	1732	405	1363	22	0
1	I	1732	405	1363	20	0
1	J	1732	405	1363	19	0
1	K	1732	405	1363	19	0
1	L	1732	405	1363	19	0
1	M	1732	405	1363	20	0
1	N	1732	405	1363	20	0
1	O	1732	405	1363	21	0
2	A	32	0	12	0	0
2	B	32	0	12	1	0
2	C	32	0	12	1	0
2	D	32	0	12	0	0
2	E	32	0	12	0	0
2	F	32	0	12	0	0
2	G	32	0	12	1	0
2	H	32	0	12	0	0
2	I	32	0	12	1	0
2	J	32	0	12	1	0
2	K	32	0	12	1	0
2	L	32	0	12	0	0
2	M	32	0	12	1	0
2	N	32	0	12	1	0
2	O	32	0	12	1	0
3	A	14	28	0	0	0
3	B	15	30	0	0	0
3	C	14	28	0	0	0
3	D	14	28	0	0	0
3	E	13	26	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	14	28	0	0	0
3	G	13	26	0	0	0
3	H	14	28	0	0	0
3	I	13	26	0	0	0
3	J	16	32	0	0	0
3	K	15	30	0	0	0
3	L	13	26	0	0	0
3	M	15	30	0	0	0
3	N	14	28	0	0	0
3	O	13	26	0	0	0
All	All	26670	6495	20625	254	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

The worst 5 of 254 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:142:GLN:O	1:B:146:GLN:HG2	1.93	0.68
1:H:142:GLN:O	1:H:146:GLN:HG2	1.95	0.66
1:J:86:ILE:HG12	1:J:165:LEU:HD13	1.78	0.66
1:L:86:ILE:HG12	1:L:165:LEU:HD13	1.78	0.66
1:F:139:ARG:HG2	1:J:187:ILE:HD12	1.77	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/221 (99%)	207 (94%)	8 (4%)	4 (2%)	13	44
1	B	219/221 (99%)	208 (95%)	6 (3%)	5 (2%)	10	36
1	C	219/221 (99%)	206 (94%)	8 (4%)	5 (2%)	10	36
1	D	219/221 (99%)	203 (93%)	11 (5%)	5 (2%)	10	36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	219/221 (99%)	204 (93%)	11 (5%)	4 (2%)	13	44
1	F	219/221 (99%)	206 (94%)	8 (4%)	5 (2%)	10	36
1	G	219/221 (99%)	206 (94%)	8 (4%)	5 (2%)	10	36
1	H	219/221 (99%)	205 (94%)	9 (4%)	5 (2%)	10	36
1	I	219/221 (99%)	203 (93%)	12 (6%)	4 (2%)	13	44
1	J	219/221 (99%)	204 (93%)	10 (5%)	5 (2%)	10	36
1	K	219/221 (99%)	203 (93%)	12 (6%)	4 (2%)	13	44
1	L	219/221 (99%)	207 (94%)	7 (3%)	5 (2%)	10	36
1	M	219/221 (99%)	207 (94%)	8 (4%)	4 (2%)	13	44
1	N	219/221 (99%)	204 (93%)	10 (5%)	5 (2%)	10	36
1	O	219/221 (99%)	203 (93%)	12 (6%)	4 (2%)	13	44
All	All	3285/3315 (99%)	3076 (94%)	140 (4%)	69 (2%)	11	39

5 of 69 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	C	2	SER
1	D	2	SER
1	E	2	SER
1	J	2	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/194 (100%)	190 (98%)	4 (2%)	66	92
1	B	194/194 (100%)	190 (98%)	4 (2%)	66	92
1	C	194/194 (100%)	191 (98%)	3 (2%)	76	95
1	D	194/194 (100%)	190 (98%)	4 (2%)	66	92
1	E	194/194 (100%)	190 (98%)	4 (2%)	66	92
1	F	194/194 (100%)	190 (98%)	4 (2%)	66	92

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	194/194 (100%)	190 (98%)	4 (2%)	66	92
1	H	194/194 (100%)	190 (98%)	4 (2%)	66	92
1	I	194/194 (100%)	191 (98%)	3 (2%)	76	95
1	J	194/194 (100%)	191 (98%)	3 (2%)	76	95
1	K	194/194 (100%)	191 (98%)	3 (2%)	76	95
1	L	194/194 (100%)	191 (98%)	3 (2%)	76	95
1	M	194/194 (100%)	190 (98%)	4 (2%)	66	92
1	N	194/194 (100%)	190 (98%)	4 (2%)	66	92
1	O	194/194 (100%)	190 (98%)	4 (2%)	66	92
All	All	2910/2910 (100%)	2855 (98%)	55 (2%)	69	93

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	93	MET
1	H	185	ARG
1	N	185	ARG
1	G	168	THR
1	H	20	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	82	ASN
1	J	82	ASN
1	O	82	ASN
1	I	82	ASN
1	I	138	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	GTP	A	401	-	34,34,34	1.42	5 (14%)	51,54,54	3.20	13 (25%)
2	GTP	B	402	-	34,34,34	1.39	5 (14%)	51,54,54	3.78	11 (21%)
2	GTP	C	403	-	34,34,34	1.39	5 (14%)	51,54,54	3.65	14 (27%)
2	GTP	D	404	-	34,34,34	1.40	4 (11%)	51,54,54	3.89	12 (23%)
2	GTP	E	405	-	34,34,34	1.52	4 (11%)	51,54,54	3.51	14 (27%)
2	GTP	F	406	-	34,34,34	1.38	4 (11%)	51,54,54	3.66	13 (25%)
2	GTP	G	407	-	34,34,34	1.47	4 (11%)	51,54,54	3.33	14 (27%)
2	GTP	H	408	-	34,34,34	1.52	4 (11%)	51,54,54	3.31	13 (25%)
2	GTP	I	409	-	34,34,34	1.50	5 (14%)	51,54,54	3.20	14 (27%)
2	GTP	J	410	-	34,34,34	1.38	5 (14%)	51,54,54	3.05	13 (25%)
2	GTP	K	411	-	34,34,34	1.49	5 (14%)	51,54,54	3.26	13 (25%)
2	GTP	L	412	-	34,34,34	1.41	6 (17%)	51,54,54	3.48	15 (29%)
2	GTP	M	413	-	34,34,34	1.38	4 (11%)	51,54,54	3.27	12 (23%)
2	GTP	N	414	-	34,34,34	1.42	5 (14%)	51,54,54	3.39	12 (23%)
2	GTP	O	415	-	34,34,34	1.41	4 (11%)	51,54,54	3.52	14 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTP	A	401	-	-	0/22/38/38	0/1/3/3

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTP	B	402	-	-	0/22/38/38	0/1/3/3
2	GTP	C	403	-	-	0/22/38/38	0/1/3/3
2	GTP	D	404	-	-	0/22/38/38	0/1/3/3
2	GTP	E	405	-	-	0/22/38/38	0/1/3/3
2	GTP	F	406	-	-	0/22/38/38	0/1/3/3
2	GTP	G	407	-	-	0/22/38/38	0/1/3/3
2	GTP	H	408	-	-	0/22/38/38	0/1/3/3
2	GTP	I	409	-	-	0/22/38/38	0/1/3/3
2	GTP	J	410	-	-	0/22/38/38	0/1/3/3
2	GTP	K	411	-	-	0/22/38/38	0/1/3/3
2	GTP	L	412	-	-	0/22/38/38	0/1/3/3
2	GTP	M	413	-	-	0/22/38/38	0/1/3/3
2	GTP	N	414	-	-	0/22/38/38	0/1/3/3
2	GTP	O	415	-	-	0/22/38/38	0/1/3/3

The worst 5 of 69 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	408	GTP	PG-O3B	4.51	1.68	1.60
2	G	407	GTP	PA-O3A	4.28	1.67	1.59
2	E	405	GTP	PA-O3A	4.26	1.67	1.59
2	D	404	GTP	PA-O3A	4.19	1.67	1.59
2	E	405	GTP	PG-O3B	3.94	1.67	1.60

The worst 5 of 197 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	404	GTP	C6-C5-N7	-24.87	130.79	134.14
2	B	402	GTP	C6-C5-N7	-24.07	130.90	134.14
2	F	406	GTP	C6-C5-N7	-23.04	131.04	134.14
2	C	403	GTP	C6-C5-N7	-22.89	131.06	134.14
2	O	415	GTP	C6-C5-N7	-21.93	131.19	134.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers

EDS was not executed - this section will therefore be empty.