



# wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 02:25 AM GMT

PDB ID : 2AAR  
Title : Structure of trigger factor binding domain in biologically homologous complex with eubacterial ribosome.  
Authors : Baram, D.; Pyetan, E.; Sittner, A.; Auerbach-Nevo, T.; Bashan, A.; Yonath, A.  
Deposited on : 2005-07-14  
Resolution : 3.50 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1243 (3.70-3.30)
Clashscore	79885	1039 (3.66-3.34)
RSRZ outliers	66119	1243 (3.70-3.30)
RNA backbone	1838	1007 (4.22-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	0	2880	
2	R	95	
3	W	67	
4	7	113	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 59630 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 23S ribosomal RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	0	2766	Total	C	N	O	P	0	0	0
			59359	26479	10949	19166	2765			

- Molecule 2 is a protein called 50S ribosomal protein L23.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
2	R	93	Total	C	0	0	93
			93	93			

- Molecule 3 is a protein called 50S ribosomal protein L29.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
3	W	65	Total	C	0	0	65
			65	65			


- Molecule 4 is a protein called Trigger Factor.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
4	7	113	Total	C	0	0	113
			113	113			

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: 23S ribosomal RNA











• Molecule 2: 50S ribosomal protein L23

Chain R:




• Molecule 3: 50S ribosomal protein L29

Chain W:



• Molecule 4: Trigger Factor

Chain 7:



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	169.39Å 407.06Å 692.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 3.50 19.99 – 3.50	Depositor EDS
% Data completeness (in resolution range)	99.0 (8.00-3.50) 100.0 (19.99-3.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.83 (at 3.52Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.251 , 0.320 0.336 , 0.367	Depositor DCC
$R_{free}$ test set	13460 reflections (4.97%)	DCC
Wilson B-factor (Å <sup>2</sup> )	98.5	Xtriage
Anisotropy	0.659	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.17 , 35.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	2 of 297349 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.80	EDS
Total number of atoms	59630	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.



## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	0	0.65	7/66467 (0.0%)	0.84	119/103673 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	0	0	128

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	0	700	C	N1-C2	10.10	1.50	1.40
1	0	538	A	C5-C6	-5.66	1.35	1.41
1	0	788	G	N9-C4	5.49	1.42	1.38
1	0	2593	A	C5-C6	-5.46	1.36	1.41
1	0	774	A	C5-C6	-5.29	1.36	1.41

The worst 5 of 119 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	0	1279	G	N9-C1'-C2'	10.85	128.10	114.00
1	0	1266	G	N9-C1'-C2'	10.62	127.81	114.00
1	0	765	C	O4'-C1'-N1	9.33	115.66	108.20
1	0	2237	C	N1-C1'-C2'	9.28	126.06	114.00
1	0	985	G	N9-C1'-C2'	9.24	126.01	114.00

There are no chirality outliers.

5 of 128 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	0	118	U	Sidechain
1	0	123	A	Sidechain
1	0	154	U	Sidechain
1	0	67	G	Sidechain
1	0	71	A	Sidechain

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	59359	0	29917	4315	0
2	R	93	0	0	0	0
3	W	65	0	0	0	0
4	7	113	0	0	0	0
All	All	59630	0	29917	4315	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 49.

The worst 5 of 4315 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:0:541:C:O2'	1:0:2018:G:N2	1.63	1.29
1:0:788:G:H22	1:0:801:A:P	1.58	1.26
1:0:1066:G:H3'	1:0:1067:G:H4'	1.23	1.20
1:0:2170:C:H2'	1:0:2171:U:H5'	1.23	1.17
1:0:1314:A:O2'	1:0:1315:A:H3'	1.42	1.15

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

There are no protein backbone outliers to report in this entry.

### 5.3.2 Protein sidechains ⓘ

There are no protein residues with a non-rotameric sidechain to report in this entry.

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	0	2757/2880 (95%)	657 (23%)	223 (8%)

5 of 657 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	0	13	A
1	0	14	A
1	0	34	U
1	0	45	C
1	0	48	A

5 of 223 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	0	1263	G
1	0	1469	U
1	0	2608	A
1	0	1278	A
1	0	1333	G

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	0	2766/2880 (96%)	0.45	271 (9%) 8 5	21, 75, 199, 205	0
2	R	93/95 (97%)	1.94	36 (38%) 1 1	16, 63, 157, 200	0
3	W	65/67 (97%)	1.71	20 (30%) 1 1	21, 86, 159, 200	0
4	7	113/113 (100%)	2.94	51 (45%) 1 1	16, 120, 200, 200	0
All	All	3037/3155 (96%)	0.61	378 (12%) 5 3	16, 76, 200, 205	0

The worst 5 of 378 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	7	110	TYR	13.4
4	7	74	TYR	13.2
2	R	90	ALA	11.8
4	7	108	GLU	11.4
1	0	1120	C	10.5

### 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.