



Full wwPDB X-ray Structure Validation Report i

Feb 28, 2014 – 06:50 PM GMT

PDB ID : 2ACS
Title : AN ANION BINDING SITE IN HUMAN ALDOSE REDUCTASE: MECHANISTIC IMPLICATIONS FOR THE BINDING OF CITRATE, CACODYLATE, AND GLUCOSE-6-PHOSPHATE
Authors : Harrison, D.H.; Bohren, K.M.; Gabbay, K.H.; Petsko, G.A.; Ringe, D.
Deposited on : 1994-04-15
Resolution : 1.76 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1304 (1.76-1.76)
Ramachandran outliers	78287	1288 (1.76-1.76)
Sidechain outliers	78261	1288 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	315	

2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 2768 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALDOSE REDUCTASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	314	2512	1614	424	462	12	0	0	0

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0

- Molecule 3 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	195	Total	O	0	0
			195	195		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: ALDOSE REDUCTASE

Chain A:



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.11Å 67.20Å 92.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.76	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-1.76)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.169 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2768	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/2574	0.67	0/3495

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	9

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	209	TYR	Sidechain
1	A	220	ALA	Peptide
1	A	221	LYS	Peptide
1	A	224	ASP	Peptide
1	A	232	ARG	Sidechain
1	A	268	ARG	Sidechain
1	A	293	ARG	Sidechain
1	A	296	ARG	Sidechain
1	A	3	ARG	Sidechain

5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2512	0	2525	34	0
2	A	48	0	25	3	0
3	A	13	0	5	2	0
4	A	195	0	0	3	0
All	All	2768	0	2555	34	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (34) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:224:ASP:HB3	1:A:225:PRO:HD3	1.41	0.98
1:A:224:ASP:HB3	1:A:225:PRO:CD	2.12	0.78
1:A:220:ALA:HB1	1:A:222:PRO:HA	1.78	0.64
1:A:110:HIS:HE1	3:A:317:CIT:O5	1.84	0.60
1:A:223:GLU:H	1:A:223:GLU:CD	2.06	0.58
1:A:220:ALA:HB1	1:A:222:PRO:CA	2.34	0.58
1:A:172:LYS:HE3	4:A:906:HOH:O	2.04	0.58
1:A:251:PHE:HZ	1:A:289:LEU:HD13	1.69	0.58
1:A:183:GLN:NE2	2:A:316:NAP:H2N	2.19	0.57
1:A:8:ASN:HD22	1:A:8:ASN:H	1.52	0.57
1:A:136:ASN:HD22	1:A:138:LEU:H	1.54	0.56
1:A:300:LEU:HD13	1:A:303:CYS:HB2	1.86	0.55
1:A:20:TRP:CE3	1:A:21:LYS:HE3	2.42	0.55
1:A:162:ASN:C	1:A:162:ASN:HD22	2.10	0.54
1:A:83:HIS:HD2	4:A:767:HOH:O	1.92	0.52
1:A:83:HIS:HE1	4:A:782:HOH:O	1.93	0.51
1:A:234:LYS:HG2	1:A:244:THR:HG21	1.94	0.48
1:A:183:GLN:NE2	2:A:316:NAP:H71N	2.13	0.47
1:A:110:HIS:CE1	3:A:317:CIT:O5	2.68	0.47
1:A:217:ARG:HD2	1:A:297:VAL:HG13	1.98	0.46
1:A:187:HIS:HD2	1:A:189:TYR:H	1.62	0.46
1:A:220:ALA:C	1:A:222:PRO:HA	2.38	0.44
1:A:217:ARG:CD	1:A:297:VAL:HG13	2.48	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:94:LYS:HD3	1:A:94:LYS:O	2.19	0.43
1:A:251:PHE:CZ	1:A:289:LEU:HD13	2.52	0.43
1:A:220:ALA:HB1	1:A:222:PRO:CG	2.50	0.42
1:A:262:LYS:O	2:A:316:NAP:H8A	2.20	0.42
1:A:4:LEU:HG	1:A:14:ILE:HG12	2.01	0.42
1:A:119:LYS:NZ	1:A:119:LYS:H	2.18	0.41
1:A:136:ASN:HD22	1:A:138:LEU:N	2.16	0.41
1:A:144:MET:O	1:A:147:LEU:HB2	2.20	0.41
1:A:242:LYS:HE3	1:A:246:GLN:OE1	2.21	0.41
1:A:220:ALA:CB	1:A:222:PRO:HA	2.50	0.40
1:A:5:LEU:HD23	1:A:11:LYS:HG3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	312/315 (99%)	305 (98%)	6 (2%)	1 (0%)	50 28

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	224	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	280/280 (100%)	257 (92%)	23 (8%)	17 2

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LEU
1	A	7	ASN
1	A	8	ASN
1	A	50	ASN
1	A	83	HIS
1	A	94	LYS
1	A	119	LYS
1	A	136	ASN
1	A	146	GLU
1	A	147	LEU
1	A	150	GLU
1	A	162	ASN
1	A	175	LEU
1	A	178	LYS
1	A	182	ASN
1	A	192	GLN
1	A	221	LYS
1	A	223	GLU
1	A	224	ASP
1	A	229	GLU
1	A	289	LEU
1	A	300	LEU
1	A	301	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	ASN
1	A	50	ASN
1	A	83	HIS
1	A	110	HIS
1	A	136	ASN
1	A	160	ASN
1	A	162	ASN
1	A	182	ASN
1	A	183	GLN
1	A	187	HIS
1	A	192	GLN
1	A	197	GLN
1	A	283	GLN
1	A	312	HIS

5.3.3 RNA i

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

There are no carbohydrates in this entry.

5.6 Ligand geometry i

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	A	316	-	52,52,52	1.20	4 (7%)	80,80,80	1.95	15 (18%)
3	CIT	A	317	-	12,12,12	3.44	3 (25%)	17,17,17	2.07	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	316	-	-	0/35/67/67	0/3/5/5
3	CIT	A	317	-	-	0/16/16/16	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	317	CIT	C2-C3	10.65	1.67	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	316	NAP	C2N-N1N	3.54	1.39	1.35
2	A	316	NAP	C4A-N9A	-2.99	1.33	1.37
3	A	317	CIT	C2-C1	2.63	1.60	1.50
2	A	316	NAP	P2B-O3X	-2.41	1.45	1.54
3	A	317	CIT	O6-C6	-2.25	1.21	1.30
2	A	316	NAP	PA-O2A	-2.18	1.45	1.55

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	316	NAP	N3A-C2A-N1A	-9.69	120.61	128.71
2	A	316	NAP	N3A-C4A-N9A	6.38	136.96	125.43
3	A	317	CIT	O6-C6-C3	5.28	120.57	112.89
2	A	316	NAP	C4A-C5A-N7A	3.88	112.84	109.52
3	A	317	CIT	C3-C2-C1	3.54	122.35	113.77
2	A	316	NAP	O3-PN-O1N	-3.48	100.52	108.83
2	A	316	NAP	C5A-C4A-N3A	-3.35	118.40	125.70
3	A	317	CIT	O2-C1-O1	-2.93	115.84	123.30
2	A	316	NAP	PN-O3-PA	2.92	145.50	132.95
2	A	316	NAP	C2A-N3A-C4A	2.73	121.78	114.01
2	A	316	NAP	O4B-C1B-N9A	2.66	110.91	108.44
2	A	316	NAP	C2D-C1D-N1N	-2.59	109.47	113.86
2	A	316	NAP	C6N-N1N-C2N	-2.56	119.14	122.04
3	A	317	CIT	O6-C6-O5	-2.52	115.83	123.76
2	A	316	NAP	O2X-P2B-O2B	-2.45	100.03	107.09
3	A	317	CIT	C3-C4-C5	-2.27	108.27	113.77
2	A	316	NAP	C2N-C3N-C4N	2.11	120.70	118.31
2	A	316	NAP	C1B-N9A-C4A	-2.10	123.01	126.64
2	A	316	NAP	O2X-P2B-O1X	2.09	117.28	110.44
2	A	316	NAP	O3D-C3D-C4D	2.08	117.20	111.08

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

6.4 Ligands

EDS was not executed - this section will therefore be empty.

6.5 Other polymers

EDS was not executed - this section will therefore be empty.