



Full wwPDB X-ray Structure Validation Report i

Mar 1, 2014 – 02:30 AM GMT

PDB ID : 3AE9
Title : Crystal structure of porcine heart mitochondrial complex II bound with N-(3-Pentafluorophenoxy-phenyl)-2-trifluoromethyl-benzamide
Authors : Harada, S.; Sasaki, T.; Shindo, M.; Kido, Y.; Inaoka, D.K.; Omori, J.; Osanai, A.; Sakamoto, K.; Mao, J.; Matsuoka, S.; Inoue, M.; Honma, T.; Tanaka, A.; Kita, K.
Deposited on : 2010-02-04
Resolution : 3.31 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1372 (3.44-3.20)
Clashscore	79885	1016 (3.42-3.22)
Ramachandran outliers	78287	1699 (3.44-3.20)
Sidechain outliers	78261	1697 (3.44-3.20)
RSRZ outliers	66119	1373 (3.44-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	622	
2	B	252	
3	C	140	
4	D	103	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
11	FD8	C	1201	-	X
12	EPH	D	1306	-	X

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 8677 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	613	Total	C	N	O	S	0	0	0
			4729	2954	848	895	32			

- Molecule 2 is a protein called Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	239	Total	C	N	O	S	0	0	0
			1922	1214	326	360	22			

- Molecule 3 is a protein called Succinate dehydrogenase cytochrome b560 subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	138	Total	C	N	O	S	0	0	0
			1064	695	179	183	7			

- Molecule 4 is a protein called Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	102	Total	C	N	O	S	0	0	0
			765	499	128	133	5			

- Molecule 5 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 6 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 7 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 8 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe_3S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 9 is MALONATE ION (three-letter code: MLI) (formula: $\text{C}_3\text{H}_2\text{O}_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			7	3	4		

- Molecule 10 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 11 is N-[3-(PENTAFLUOROPHENOXY)PHENYL]-2-(TRIFLUOROMETHYL) BENZAMIDE (three-letter code: FD8) (formula: $C_{20}H_9F_8NO_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	C	1	Total	C	F	N	O	0	0
			31	20	8	1	2		

- Molecule 12 is L-ALPHA-PHOSPHATIDYL-BETA-OLEOYL-GAMMA-PALMITOYL-PHOSPHATIDYLETHANOLAMINE (three-letter code: EPH) (formula: $C_{39}H_{68}NO_8P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	D	1	Total	C	N	O	P	0	0
			44	34	1	8	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial

Chain A: 



- Molecule 2: Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial

Chain B: 



- Molecule 3: Succinate dehydrogenase cytochrome b560 subunit, mitochondrial

Chain C: 



- Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.61Å 84.25Å 295.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.76 – 3.31 37.76 – 3.31	Depositor EDS
% Data completeness (in resolution range)	97.8 (37.76-3.31) 97.8 (37.76-3.31)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 3.32Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.257 , 0.307 0.257 , 0.304	Depositor DCC
R_{free} test set	1367 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	97.3	Xtriage
Anisotropy	0.506	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 55.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 27276 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8677	wwPDB-VP
Average B, all atoms (Å ²)	124.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, MLI, F3S, FES, EPH, HEM, FD8, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.31	0/4828	0.47	0/6531
2	B	0.31	0/1964	0.46	0/2648
3	C	0.32	0/1091	0.43	0/1483
4	D	0.29	0/784	0.42	0/1066
All	All	0.31	0/8667	0.46	0/11728

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4729	0	0	5	0
2	B	1922	0	0	0	0
3	C	1064	0	0	0	0
4	D	765	0	0	0	0
5	A	53	0	31	4	0
6	B	4	0	0	0	0
7	B	8	0	0	0	0
8	B	7	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	A	7	0	2	0	0
10	C	43	0	30	0	0
11	C	31	0	9	2	0
12	D	44	0	53	0	0
All	All	8677	0	125	7	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

All (7) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:57:HIS:NE2	5:A:700:FAD:HM82	1.36	1.41
1:A:57:HIS:CE1	5:A:700:FAD:HM82	2.00	0.96
1:A:57:HIS:NE2	5:A:700:FAD:HM81	2.10	0.61
11:C:1201:FD8:H14	11:C:1201:FD8:O1	2.03	0.58
1:A:415:LEU:N	5:A:700:FAD:O2	2.51	0.43
11:C:1201:FD8:C14	11:C:1201:FD8:O1	2.69	0.41
1:A:57:HIS:CG	1:A:57:HIS:O	2.75	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	611/622 (98%)	571 (94%)	38 (6%)	2 (0%)	50	92
2	B	237/252 (94%)	222 (94%)	15 (6%)	0	100	100
3	C	136/140 (97%)	129 (95%)	7 (5%)	0	100	100
4	D	100/103 (97%)	97 (97%)	3 (3%)	0	100	100
All	All	1084/1117 (97%)	1019 (94%)	63 (6%)	2 (0%)	56	94

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	290	PRO
1	A	260	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	499/506 (99%)	482 (97%)	17 (3%)	49	86
2	B	214/220 (97%)	203 (95%)	11 (5%)	33	77
3	C	117/118 (99%)	115 (98%)	2 (2%)	73	94
4	D	76/76 (100%)	74 (97%)	2 (3%)	59	90
All	All	906/920 (98%)	874 (96%)	32 (4%)	48	86

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	THR
1	A	65	ILE
1	A	74	GLU
1	A	121	THR
1	A	129	ARG
1	A	131	PHE
1	A	157	SER
1	A	165	ARG
1	A	167	LEU
1	A	313	ARG
1	A	368	MET
1	A	418	LEU
1	A	458	ARG
1	A	507	LEU
1	A	556	LYS
1	A	561	GLU
1	A	597	LEU
2	B	10	ARG
2	B	51	ILE
2	B	63	ARG
2	B	66	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	87	ARG
2	B	159	ILE
2	B	189	ARG
2	B	192	ILE
2	B	214	ARG
2	B	217	THR
2	B	230	ASN
3	C	52	LEU
3	C	122	LEU
4	D	78	LEU
4	D	108	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	FAD	A	700	1	58,58,58	1.06	5 (8%)	85,89,89	2.04	13 (15%)
9	MLI	A	701	-	2,6,6	2.68	2 (100%)	0,7,7	0.00	-
6	FES	B	302	2	0,4,4	0.00	-	0,4,4	0.00	-
7	SF4	B	303	2	12,12,12	4.13	12 (100%)	0,24,24	0.00	-
8	F3S	B	304	2	3,9,9	18.97	3 (100%)	0,15,15	0.00	-
11	FD8	C	1201	-	33,33,33	3.38	8 (24%)	49,49,49	0.73	0
10	HEM	C	1305	3,4	49,50,50	2.39	15 (30%)	46,82,82	2.04	7 (15%)
12	EPH	D	1306	-	42,43,48	2.05	7 (16%)	46,48,53	2.07	9 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	FAD	A	700	1	-	0/34/50/50	0/1/6/6
9	MLI	A	701	-	-	0/0/4/4	0/0/0/0
6	FES	B	302	2	-	0/0/4/4	0/0/1/1
7	SF4	B	303	2	-	0/0/48/48	0/0/5/5
8	F3S	B	304	2	-	0/0/24/24	0/0/3/3
11	FD8	C	1201	-	-	0/18/18/18	0/3/3/3
10	HEM	C	1305	3,4	-	0/14/114/114	0/0/8/8
12	EPH	D	1306	-	-	0/47/47/52	0/0/0/0

All (52) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	304	F3S	S3-FE1	-19.71	2.20	2.33
8	B	304	F3S	S3-FE3	-19.33	2.20	2.33
8	B	304	F3S	S3-FE4	-17.83	2.21	2.33
12	D	1306	EPH	C31-C30	-8.49	1.51	1.55
11	C	1201	FD8	C17-C16	7.87	1.53	1.37
11	C	1201	FD8	C18-C17	7.84	1.53	1.37
11	C	1201	FD8	C20-C19	7.82	1.53	1.37
11	C	1201	FD8	C19-C18	7.82	1.53	1.37
11	C	1201	FD8	C15-C20	7.35	1.53	1.38
11	C	1201	FD8	C15-C16	7.20	1.53	1.38
10	C	1305	HEM	C3D-C2D	5.84	1.54	1.43
10	C	1305	HEM	C3B-C2B	-5.40	1.34	1.43
10	C	1305	HEM	C3C-C2C	-5.39	1.34	1.43
10	C	1305	HEM	C3C-CAC	4.67	1.55	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	C	1305	HEM	C3B-CAB	4.58	1.54	1.40
10	C	1305	HEM	C4A-C3A	4.49	1.45	1.40
7	B	303	SF4	S3-FE2	-4.45	2.30	2.33
7	B	303	SF4	S4-FE3	-4.35	2.30	2.33
7	B	303	SF4	S3-FE4	-4.32	2.30	2.33
7	B	303	SF4	S1-FE2	-4.23	2.30	2.33
12	D	1306	EPH	O1-C3	4.17	1.47	1.34
12	D	1306	EPH	C16-C15	4.13	1.52	1.30
7	B	303	SF4	S4-FE2	-4.12	2.30	2.33
12	D	1306	EPH	O2-C4	4.05	1.46	1.33
7	B	303	SF4	S4-FE1	-4.05	2.30	2.33
7	B	303	SF4	S1-FE4	-4.03	2.30	2.33
7	B	303	SF4	S1-FE3	-4.02	2.30	2.33
7	B	303	SF4	S2-FE1	-4.02	2.30	2.33
10	C	1305	HEM	C2B-C1B	4.01	1.45	1.44
7	B	303	SF4	S2-FE4	-4.01	2.30	2.33
7	B	303	SF4	S2-FE3	-3.99	2.30	2.33
7	B	303	SF4	S3-FE1	-3.89	2.30	2.33
10	C	1305	HEM	FE-NA	3.85	2.09	1.92
12	D	1306	EPH	C25-C24	3.84	1.52	1.31
12	D	1306	EPH	C29-C28	3.84	1.52	1.31
12	D	1306	EPH	C13-C12	3.84	1.52	1.31
10	C	1305	HEM	FE-ND	3.74	2.11	1.97
10	C	1305	HEM	C3D-C4D	3.61	1.45	1.44
5	A	700	FAD	C2A-N3A	3.36	1.38	1.32
5	A	700	FAD	C1'-C2'	2.83	1.54	1.51
11	C	1201	FD8	C9-N	-2.78	1.36	1.41
9	A	701	MLI	C1-C3	2.71	1.51	1.49
10	C	1305	HEM	C2D-C1D	2.70	1.45	1.44
5	A	700	FAD	C1'-N10	2.68	1.51	1.48
9	A	701	MLI	C1-C2	2.65	1.51	1.49
10	C	1305	HEM	CMB-C2B	2.53	1.55	1.47
10	C	1305	HEM	CMC-C2C	2.51	1.55	1.47
5	A	700	FAD	C2A-N1A	2.45	1.38	1.33
10	C	1305	HEM	CMD-C2D	2.41	1.54	1.47
5	A	700	FAD	C5X-N5	2.20	1.38	1.35
10	C	1305	HEM	FE-NB	2.14	2.05	1.97
11	C	1201	FD8	O2-C15	2.07	1.43	1.39

All (29) bond angle outliers are listed below:

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	700	FAD	N3A-C2A-N1A	-10.86	119.63	128.71
12	D	1306	EPH	C14-C15-C16	-10.21	111.10	126.73
10	C	1305	HEM	C3B-C4B-NB	-8.82	107.69	114.00
5	A	700	FAD	O4B-C1B-N9A	6.85	114.81	108.44
10	C	1305	HEM	C4D-ND-C1D	6.45	111.76	105.16
5	A	700	FAD	C2-N1-C10	5.83	120.86	114.98
12	D	1306	EPH	O1-C3-C5	4.31	120.99	111.56
5	A	700	FAD	C2'-C1'-N10	-4.14	106.96	112.45
12	D	1306	EPH	C30-C29-C28	-4.06	110.89	125.02
5	A	700	FAD	N3A-C4A-N9A	3.98	132.63	125.43
10	C	1305	HEM	C2D-C1D-ND	-3.69	108.57	112.93
5	A	700	FAD	C4X-C10-N1	-3.52	119.21	122.73
5	A	700	FAD	C4X-N5-C5X	3.48	120.60	116.69
10	C	1305	HEM	C1B-NB-C4B	3.12	108.36	105.16
5	A	700	FAD	P-O3P-PA	-3.12	122.55	131.68
5	A	700	FAD	C5X-C9A-N10	3.10	119.86	116.80
10	C	1305	HEM	C4C-NC-C1C	2.66	108.30	105.53
5	A	700	FAD	C4-N3-C2	-2.64	119.97	125.39
12	D	1306	EPH	C11-C12-C13	-2.55	110.98	125.43
12	D	1306	EPH	C26-C25-C24	-2.55	111.00	125.43
12	D	1306	EPH	O2-C4-C18	2.51	119.82	111.94
12	D	1306	EPH	C23-C24-C25	-2.49	111.31	125.43
12	D	1306	EPH	C27-C28-C29	-2.40	111.81	125.43
5	A	700	FAD	C5A-C4A-N3A	-2.38	120.51	125.70
10	C	1305	HEM	CAD-C3D-C4D	2.32	128.70	124.53
5	A	700	FAD	C4B-O4B-C1B	-2.31	107.24	109.75
5	A	700	FAD	C2A-N3A-C4A	2.22	120.32	114.01
12	D	1306	EPH	C14-C13-C12	-2.21	111.25	124.86
10	C	1305	HEM	CMA-C3A-C4A	-2.01	125.53	128.62

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	613/622 (98%)	0.11	9 (1%)	70 24	89, 120, 184, 205	0
2	B	239/252 (94%)	0.10	2 (0%)	83 39	91, 113, 152, 184	0
3	C	138/140 (98%)	-0.04	2 (1%)	72 25	94, 117, 183, 207	0
4	D	102/103 (99%)	-0.08	1 (0%)	79 33	98, 120, 160, 176	0
All	All	1092/1117 (97%)	0.07	14 (1%)	72 27	89, 118, 178, 207	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	35	SER	3.1
1	A	326	LEU	3.0
1	A	327	HIS	2.8
3	C	81	CYS	2.8
1	A	356	GLU	2.6
1	A	357	PRO	2.4
1	A	622	TYR	2.3
2	B	34	GLU	2.3
3	C	143	MET	2.1
2	B	11	ILE	2.1
1	A	325	GLN	2.1
1	A	277	ILE	2.0
1	A	280	GLN	2.0
1	A	321	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
12	EPH	D	1306	44/49	0.62	3.91	150,153,156,156	0
11	FD8	C	1201	31/31	0.36	3.67	136,137,138,138	0
10	HEM	C	1305	43/43	0.27	1.77	101,102,103,104	0
5	FAD	A	700	53/53	0.37	0.95	98,100,101,101	0
9	MLI	A	701	7/7	0.33	0.55	222,222,222,222	0
8	F3S	B	304	7/7	0.22	-0.46	99,99,99,99	0
7	SF4	B	303	8/8	0.21	-0.93	85,85,86,86	0
6	FES	B	302	4/4	0.22	-0.95	100,100,100,101	0

6.5 Other polymers

There are no such residues in this entry.