



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 10:46 PM GMT

PDB ID : 4AID
Title : Crystal structure of C. crescentus PNPase bound to RNase E recognition peptide
Authors : Hardwick, S.W.; Gubbey, T.; Hug, I.; Jenal, U.; Luisi, B.F.
Deposited on : 2012-02-09
Resolution : 2.60 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1718 (2.60-2.60)
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)
RSRZ outliers	66119	1718 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	726	
1	B	726	
1	C	726	
2	F	14	
2	G	14	
2	H	14	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12655 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called POLYRIBONUCLEOTIDE NUCLEOTIDYLTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	573	Total	C	N	O	S	0	1	0
			4248	2686	720	819	23			
1	B	558	Total	C	N	O	S	0	0	0
			4170	2639	706	802	23			
1	C	539	Total	C	N	O	S	0	0	0
			3836	2425	654	735	22			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	EXPRESSION TAG	UNP Q9AC32
A	-12	GLY	-	EXPRESSION TAG	UNP Q9AC32
A	-11	SER	-	EXPRESSION TAG	UNP Q9AC32
A	-10	SER	-	EXPRESSION TAG	UNP Q9AC32
A	-9	HIS	-	EXPRESSION TAG	UNP Q9AC32
A	-8	HIS	-	EXPRESSION TAG	UNP Q9AC32
A	-7	HIS	-	EXPRESSION TAG	UNP Q9AC32
A	-6	HIS	-	EXPRESSION TAG	UNP Q9AC32
A	-5	HIS	-	EXPRESSION TAG	UNP Q9AC32
A	-4	HIS	-	EXPRESSION TAG	UNP Q9AC32
A	-3	SER	-	EXPRESSION TAG	UNP Q9AC32
A	-2	GLN	-	EXPRESSION TAG	UNP Q9AC32
A	-1	ASP	-	EXPRESSION TAG	UNP Q9AC32
A	0	PRO	-	EXPRESSION TAG	UNP Q9AC32
B	-13	MET	-	EXPRESSION TAG	UNP Q9AC32
B	-12	GLY	-	EXPRESSION TAG	UNP Q9AC32
B	-11	SER	-	EXPRESSION TAG	UNP Q9AC32
B	-10	SER	-	EXPRESSION TAG	UNP Q9AC32
B	-9	HIS	-	EXPRESSION TAG	UNP Q9AC32
B	-8	HIS	-	EXPRESSION TAG	UNP Q9AC32
B	-7	HIS	-	EXPRESSION TAG	UNP Q9AC32
B	-6	HIS	-	EXPRESSION TAG	UNP Q9AC32

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	HIS	-	EXPRESSION TAG	UNP Q9AC32
B	-4	HIS	-	EXPRESSION TAG	UNP Q9AC32
B	-3	SER	-	EXPRESSION TAG	UNP Q9AC32
B	-2	GLN	-	EXPRESSION TAG	UNP Q9AC32
B	-1	ASP	-	EXPRESSION TAG	UNP Q9AC32
B	0	PRO	-	EXPRESSION TAG	UNP Q9AC32
C	-13	MET	-	EXPRESSION TAG	UNP Q9AC32
C	-12	GLY	-	EXPRESSION TAG	UNP Q9AC32
C	-11	SER	-	EXPRESSION TAG	UNP Q9AC32
C	-10	SER	-	EXPRESSION TAG	UNP Q9AC32
C	-9	HIS	-	EXPRESSION TAG	UNP Q9AC32
C	-8	HIS	-	EXPRESSION TAG	UNP Q9AC32
C	-7	HIS	-	EXPRESSION TAG	UNP Q9AC32
C	-6	HIS	-	EXPRESSION TAG	UNP Q9AC32
C	-5	HIS	-	EXPRESSION TAG	UNP Q9AC32
C	-4	HIS	-	EXPRESSION TAG	UNP Q9AC32
C	-3	SER	-	EXPRESSION TAG	UNP Q9AC32
C	-2	GLN	-	EXPRESSION TAG	UNP Q9AC32
C	-1	ASP	-	EXPRESSION TAG	UNP Q9AC32
C	0	PRO	-	EXPRESSION TAG	UNP Q9AC32

- Molecule 2 is a protein called RIBONUCLEASE, RNE/RNG FAMILY PROTEIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	F	7	Total	C	N	O	0	0	0
			66	44	15	7			
2	G	8	Total	C	N	O	0	0	0
			77	50	19	8			
2	H	5	Total	C	N	O	0	0	0
			42	30	7	5			


- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	C	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	96	Total	O	0	0
			96	96		
4	B	80	Total	O	0	0
			80	80		
4	C	24	Total	O	0	0
			24	24		
4	G	1	Total	O	0	0
			1	1		




• Molecule 1: POLYRIBONUCLEOTIDE NUCLEOTIDYLTRANSFERASE

Chain C:




• Molecule 2: RIBONUCLEASE, RNE/RNG FAMILY PROTEIN

Chain F:



• Molecule 2: RIBONUCLEASE, RNE/RNG FAMILY PROTEIN

Chain G:



• Molecule 2: RIBONUCLEASE, RNE/RNG FAMILY PROTEIN

Chain H: 

THR	ALA	PRO	PRO	GLU	LYS	PRO	ARG	E893	+	W896	E897	ARG
-----	-----	-----	-----	-----	-----	-----	-----	------	---	------	------	-----

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	157.44Å 157.44Å 302.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.60 29.75 – 2.60	Depositor EDS
% Data completeness (in resolution range)	90.0 (30.00-2.60) 90.0 (29.75-2.60)	Depositor EDS
R_{merge}	0.22	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.209 , 0.254 0.190 , 0.209	Depositor DCC
R_{free} test set	2319 reflections (3.09%)	DCC
Wilson B-factor (Å ²)	46.7	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 19.5	EDS
Estimated twinning fraction	0.357 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 77316 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12655	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	8/4324 (0.2%)	1.15	27/5859 (0.5%)
1	B	1.03	9/4242 (0.2%)	1.21	33/5744 (0.6%)
1	C	0.66	2/3900 (0.1%)	0.87	3/5305 (0.1%)
2	F	1.59	2/70 (2.9%)	1.32	1/94 (1.1%)
2	G	1.30	1/81 (1.2%)	1.14	0/109
2	H	1.49	1/45 (2.2%)	1.03	0/62
All	All	0.94	23/12662 (0.2%)	1.10	64/17173 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	3
1	C	0	1
2	F	0	1
All	All	0	7

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	896	TRP	CD2-CE2	7.50	1.50	1.41
1	B	81	GLU	CD-OE1	6.69	1.33	1.25
2	G	896	TRP	CD2-CE2	6.34	1.49	1.41
1	A	432	GLU	CD-OE1	6.33	1.32	1.25
1	A	332	GLU	CD-OE1	6.26	1.32	1.25

The worst 5 of 64 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	177	LEU	CB-CG-CD1	-10.79	92.66	111.00
1	B	28	ASP	CB-CG-OD2	10.72	127.95	118.30
1	B	3	ASP	CB-CG-OD1	-10.37	108.97	118.30
1	B	153	ARG	NE-CZ-NH1	9.91	125.26	120.30
1	B	486	ASP	CB-CG-OD1	-9.45	109.80	118.30

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	554	ALA	Peptide
1	A	8	THR	Peptide
1	B	554	ALA	Peptide
1	B	6	ARG	Peptide
1	B	9	ILE	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4248	0	2	85	1
1	B	4170	0	0	84	1
1	C	3836	0	0	47	2
2	F	66	0	58	1	0
2	G	77	0	70	1	0
2	H	42	0	26	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
4	A	96	0	0	3	0
4	B	80	0	0	3	0
4	C	24	0	0	2	0
4	G	1	0	0	0	0
All	All	12655	0	156	215	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including

hydrogens) of the entry. The overall clashscore for this entry is 17.

The worst 5 of 215 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:287:PRO:CB	1:A:288:THR:CG2	1.84	1.53
1:A:281:GLY:CA	1:A:289:GLY:O	1.68	1.37
1:B:291:ASP:O	1:B:293:LEU:N	1.61	1.29
1:B:279:ALA:O	1:B:282:LEU:CD2	1.97	1.13
1:A:287:PRO:CA	1:A:288:THR:CG2	2.28	1.10

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:8:THR:OG1	1:C:540:ASN:O[2_555]	1.87	0.33
1:B:6:ARG:NH2	1:C:224:GLU:OE2[3_565]	2.10	0.10

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	570/726 (78%)	502 (88%)	37 (6%)	31 (5%)	3	3
1	B	556/726 (77%)	490 (88%)	41 (7%)	25 (4%)	4	4
1	C	535/726 (74%)	476 (89%)	32 (6%)	27 (5%)	3	4
2	F	5/14 (36%)	4 (80%)	0	1 (20%)	0	0
2	G	6/14 (43%)	5 (83%)	0	1 (17%)	0	0
2	H	3/14 (21%)	2 (67%)	1 (33%)	0	100	100
All	All	1675/2220 (76%)	1479 (88%)	111 (7%)	85 (5%)	3	3

5 of 85 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	ILE
1	A	55	ASP
1	A	239	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	248	LEU
1	A	262	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	432/578 (75%)	401 (93%)	31 (7%)	21	39
1	B	428/578 (74%)	397 (93%)	31 (7%)	21	39
1	C	364/578 (63%)	348 (96%)	16 (4%)	39	68
2	F	5/12 (42%)	3 (60%)	2 (40%)	0	0
2	G	6/12 (50%)	5 (83%)	1 (17%)	3	5
2	H	2/12 (17%)	2 (100%)	0	100	100
All	All	1237/1770 (70%)	1156 (94%)	81 (6%)	24	46

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	64	GLN
1	B	242	LYS
1	C	496	LYS
1	B	83	ARG
1	B	188	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	A	1614	-	4,4,4	0.14	0	6,6,6	0.35	0
3	PO4	B	1557	-	4,4,4	0.08	0	6,6,6	0.31	0
3	PO4	C	1556	-	4,4,4	0.18	0	6,6,6	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	A	1614	-	-	0/0/0/0	0/0/0/0
3	PO4	B	1557	-	-	0/0/0/0	0/0/0/0
3	PO4	C	1556	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	573/726 (78%)	-0.24	15 (2%) 53 50	24, 34, 123, 161	0
1	B	558/726 (76%)	-0.29	3 (0%) 88 90	24, 33, 109, 147	0
1	C	539/726 (74%)	0.04	16 (2%) 48 45	53, 87, 184, 253	0
2	F	7/14 (50%)	-0.13	0 100 100	40, 67, 84, 94	0
2	G	8/14 (57%)	-0.11	0 100 100	37, 68, 86, 93	0
2	H	5/14 (35%)	0.07	0 100 100	82, 97, 103, 107	0
All	All	1690/2220 (76%)	-0.16	34 (2%) 62 60	24, 49, 135, 253	0

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	251	ALA	5.3
1	A	278	ALA	4.6
1	C	268	ALA	4.2
1	C	269	VAL	3.9
1	C	278	ALA	3.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	PO4	C	1556	5/5	0.19	1.15	86,89,109,117	0
3	PO4	B	1557	5/5	0.11	-1.61	76,82,92,104	0
3	PO4	A	1614	5/5	0.10	-3.21	58,60,69,75	0

6.5 Other polymers

There are no such residues in this entry.