



Full wwPDB X-ray Structure Validation Report i

Feb 26, 2014 – 03:16 PM GMT

PDB ID : 3AJX
Title : Crystal Structure of 3-Hexulose-6-PhosphateSynthase
Authors : Kita, A.; Orita, I.; Yurimoto, H.; Kato, N.; Sakai, Y.; Miki, K.
Deposited on : 2010-06-24
Resolution : 1.60 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1872 (1.60-1.60)
Clashscore	79885	2199 (1.60-1.60)
Ramachandran outliers	78287	2126 (1.60-1.60)
Sidechain outliers	78261	2125 (1.60-1.60)
RSRZ outliers	66119	1872 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	207	
1	B	207	
1	C	207	
1	D	207	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	MG	A	9100	-	X
3	MG	B	9200	-	X
3	MG	C	9300	-	X
3	MG	D	9400	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
4	SO4	A	6001	-	X
4	SO4	B	6002	-	X
4	SO4	C	6003	-	X
4	SO4	D	6004	-	X

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6796 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-hexulose-6-phosphatesynthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	207	1471	932	249	286	4	0	0	0
1	B	207	1471	932	249	286	4	0	0	0
1	C	207	1471	932	249	286	4	0	0	0
1	D	207	1471	932	249	286	4	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1037	LYS	GLU	ENGINEERED MUTATION	UNP Q9LBW4
B	2037	LYS	GLU	ENGINEERED MUTATION	UNP Q9LBW4
C	3037	LYS	GLU	ENGINEERED MUTATION	UNP Q9LBW4
D	4037	LYS	GLU	ENGINEERED MUTATION	UNP Q9LBW4

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	C	2	Total	Cl	0	0
			2	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total Mg 1 1	0	0
3	C	1	Total Mg 1 1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	228	Total O 228 228	0	0
5	B	209	Total O 209 209	0	0
5	C	241	Total O 241 241	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	207	Total 207	O 207	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-hexulose-6-phosphatesynthase

Chain A: 




- Molecule 1: 3-hexulose-6-phosphatesynthase

Chain B: 



- Molecule 1: 3-hexulose-6-phosphatesynthase

Chain C: 



- Molecule 1: 3-hexulose-6-phosphatesynthase

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.04Å 57.35Å 113.62Å 90.00° 90.69° 90.00°	Depositor
Resolution (Å)	43.04 – 1.60 43.04 – 1.60	Depositor EDS
% Data completeness (in resolution range)	97.5 (43.04-1.60) 97.7 (43.04-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 1.60Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.194 , 0.215 0.195 , 0.215	Depositor DCC
R_{free} test set	5442 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	0.340	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 38.4	EDS
Estimated twinning fraction	0.019 for h,-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	5 of 108052 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6796	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 74.51 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.5078e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/1486	0.57	0/2010
1	B	0.28	0/1486	0.57	0/2010
1	C	0.27	0/1486	0.57	0/2010
1	D	0.28	0/1486	0.57	0/2010
All	All	0.27	0/5944	0.57	0/8040

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1471	0	1527	11	0
1	B	1471	0	1527	21	0
1	C	1471	0	1527	12	0
1	D	1471	0	1527	14	0
2	A	1	0	0	0	0
2	C	2	0	0	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
4	C	5	0	0	0	0
4	D	5	0	0	0	0
5	A	228	0	0	3	0
5	B	209	0	0	3	0
5	C	241	0	0	2	0
5	D	207	0	0	5	0
All	All	6796	0	6108	55	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

All (55) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:1134:HIS:CE1	5:A:5854:HOH:O	1.91	1.19
1:D:4134:HIS:CE1	5:D:5880:HOH:O	2.10	1.03
1:B:2134:HIS:CE1	5:B:5863:HOH:O	2.09	1.03
1:A:1134:HIS:HE1	5:A:5854:HOH:O	1.28	1.01
1:B:2134:HIS:HE1	5:B:5863:HOH:O	1.44	0.95
1:D:4031:LEU:HD11	1:D:4047:VAL:HG21	1.62	0.81
1:D:4063:MET:HG2	1:D:4084:LEU:HD12	1.62	0.79
1:A:1031:LEU:HD11	1:A:1047:VAL:HG21	1.68	0.74
1:B:2031:LEU:HD11	1:B:2047:VAL:HG21	1.72	0.72
1:C:3169:VAL:HG21	1:C:3201:GLU:HG2	1.78	0.66
1:C:3031:LEU:HD11	1:C:3047:VAL:HG21	1.80	0.64
1:B:2197:ALA:O	1:B:2201:GLU:HG3	2.01	0.61
1:B:2113:ILE:HD13	1:B:2119:ARG:HG2	1.83	0.59
1:B:2075:LYS:HB2	1:B:2075:LYS:NZ	2.17	0.59
1:B:2168:LYS:HE3	5:B:5847:HOH:O	2.03	0.57
1:D:4113:ILE:HD13	1:D:4119:ARG:HG2	1.88	0.54
1:A:1132:GLU:HG2	1:A:1134:HIS:HB2	1.90	0.52
1:B:2146:ASP:OD2	1:B:2148:ASN:HB2	2.11	0.51
1:A:1033:THR:N	1:A:1034:PRO:HD2	2.27	0.50
1:B:2132:GLU:HG2	1:B:2134:HIS:HB2	1.93	0.50
1:A:1168:LYS:HE3	5:A:5071:HOH:O	2.13	0.49
1:B:2159:VAL:HG13	1:B:2160:PRO:HD2	1.95	0.49
1:B:2082:THR:HA	1:B:2107:VAL:O	2.13	0.48
2:C:7002:CL:CL	1:D:4138:ASP:OD1	2.68	0.48
1:D:4134:HIS:NE2	5:D:5884:HOH:O	2.35	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:3033:THR:N	1:C:3034:PRO:HD2	2.28	0.48
1:C:3132:GLU:HG2	1:C:3134:HIS:HB2	1.96	0.47
1:B:2147:LEU:HD12	1:B:2150:LEU:HD12	1.97	0.46
1:C:3100:GLN:NE2	5:C:5657:HOH:O	2.48	0.46
1:D:4134:HIS:HE1	5:D:5880:HOH:O	1.73	0.46
1:A:1041:LEU:HG	1:B:2038:ALA:HA	1.98	0.46
1:D:4132:GLU:HG2	1:D:4134:HIS:HB2	1.98	0.45
1:C:3031:LEU:N	1:C:3031:LEU:HD22	2.30	0.45
1:A:1082:THR:HA	1:A:1107:VAL:O	2.17	0.45
1:B:2172:ILE:HB	1:B:2173:PRO:HD3	1.98	0.45
1:C:3030:GLU:HG3	1:C:3057:PHE:CD1	2.52	0.45
1:A:1030:GLU:HG3	1:A:1057:PHE:CD2	2.52	0.45
1:B:2118:THR:O	1:B:2122:GLU:HG3	2.17	0.45
1:C:3074:PHE:CE2	1:C:3081:VAL:HG13	2.52	0.44
1:B:2151:LEU:O	1:B:2155:GLU:HG3	2.18	0.44
1:D:4157:ALA:HB3	5:D:5096:HOH:O	2.18	0.44
1:C:3172:ILE:HB	1:C:3173:PRO:HD3	2.00	0.44
1:D:4030:GLU:HG3	1:D:4057:PHE:CD1	2.53	0.44
1:B:2030:GLU:HG3	1:B:2057:PHE:CD1	2.53	0.43
1:C:3134:HIS:HE1	5:C:5872:HOH:O	2.01	0.43
1:D:4082:THR:HA	1:D:4107:VAL:O	2.19	0.43
1:C:3041:LEU:HG	1:D:4038:ALA:HA	1.99	0.43
1:A:1147:LEU:HD12	1:A:1150:LEU:HD12	1.98	0.43
1:D:4201:GLU:HG3	5:D:5315:HOH:O	2.18	0.43
1:C:3082:THR:HA	1:C:3107:VAL:O	2.17	0.43
1:D:4074:PHE:CE2	1:D:4081:VAL:HG13	2.55	0.42
1:B:2158:ARG:HG3	1:B:2158:ARG:HH11	1.84	0.42
1:B:2075:LYS:HB2	1:B:2075:LYS:HZ2	1.85	0.42
1:B:2008:ASP:OD2	1:B:2032:GLY:HA3	2.20	0.41
1:A:1038:ALA:HA	1:B:2041:LEU:HG	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/207 (99%)	200 (98%)	4 (2%)	1 (0%)	38	13
1	B	205/207 (99%)	201 (98%)	3 (2%)	1 (0%)	38	13
1	C	205/207 (99%)	201 (98%)	3 (2%)	1 (0%)	38	13
1	D	205/207 (99%)	201 (98%)	3 (2%)	1 (0%)	38	13
All	All	820/828 (99%)	803 (98%)	13 (2%)	4 (0%)	38	13

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2165	GLY
1	D	4165	GLY
1	A	1165	GLY
1	C	3165	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/141 (100%)	141 (100%)	0	100	100
1	B	141/141 (100%)	140 (99%)	1 (1%)	91	81
1	C	141/141 (100%)	141 (100%)	0	100	100
1	D	141/141 (100%)	141 (100%)	0	100	100
All	All	564/564 (100%)	563 (100%)	1 (0%)	96	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2075	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1004	GLN
1	A	1100	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1121	GLN
1	A	1134	HIS
1	B	2004	GLN
1	B	2134	HIS
1	C	3004	GLN
1	C	3100	GLN
1	C	3121	GLN
1	D	4004	GLN
1	D	4121	GLN

5.3.3 RNA [i](#)

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 7 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	6001	-	4,4,4	0.29	0	6,6,6	0.13	0
4	SO4	B	6002	-	4,4,4	0.20	0	6,6,6	0.07	0
4	SO4	C	6003	-	4,4,4	0.31	0	6,6,6	0.17	0
4	SO4	D	6004	-	4,4,4	0.21	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SO4	A	6001	-	-	0/0/0/0	0/0/0/0
4	SO4	B	6002	-	-	0/0/0/0	0/0/0/0
4	SO4	C	6003	-	-	0/0/0/0	0/0/0/0
4	SO4	D	6004	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/207 (100%)	-0.10	2 (0%) 79 81	6, 13, 23, 31	0
1	B	207/207 (100%)	-0.20	3 (1%) 72 73	8, 14, 24, 33	0
1	C	207/207 (100%)	-0.24	1 (0%) 88 90	5, 11, 19, 31	0
1	D	207/207 (100%)	-0.19	2 (0%) 79 81	8, 13, 23, 32	0
All	All	828/828 (100%)	-0.18	8 (0%) 79 81	5, 12, 22, 33	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2144	GLY	4.4
1	D	4144	GLY	4.2
1	D	4143	PRO	3.5
1	A	1144	GLY	3.1
1	A	1143	PRO	2.7
1	B	2143	PRO	2.3
1	B	2146	ASP	2.2
1	C	3144	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SO4	C	6003	5/5	0.28	15.27	23,25,30,31	0
3	MG	A	9100	1/1	0.23	14.79	34,34,34,34	0
3	MG	B	9200	1/1	0.30	13.30	28,28,28,28	0
3	MG	D	9400	1/1	0.34	8.05	35,35,35,35	0
4	SO4	D	6004	5/5	0.17	6.07	30,32,33,34	0
4	SO4	B	6002	5/5	0.19	4.30	27,30,32,33	0
4	SO4	A	6001	5/5	0.14	4.09	21,21,23,24	0
3	MG	C	9300	1/1	0.12	2.80	42,42,42,42	0
2	CL	C	7002	1/1	0.12	1.57	22,22,22,22	0
2	CL	A	7001	1/1	0.04	-3.24	20,20,20,20	0
2	CL	C	7003	1/1	0.03	-8.44	18,18,18,18	0

6.5 Other polymers (i)

There are no such residues in this entry.