



# wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 08:15 PM GMT

PDB ID : 1ALX  
Title : GRAMICIDIN D FROM BACILLUS BREVIS (METHANOL SOLVATE)  
Authors : Burkhart, B.M.; Langs, D.A.; Smith, G.D.; Courseille, C.; Precigoux, G.;  
Hospital, M.; Pangborn, W.A.; Duax, W.L.  
Deposited on : 1997-06-05  
Resolution : 1.20 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	<b>FAILED</b>
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1158 (1.26-1.14)
Ramachandran outliers	78287	1106 (1.26-1.14)
Sidechain outliers	78261	1104 (1.26-1.14)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	17	
2	B	16	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 866 atoms, of which 483 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GRAMICIDIN A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	16	Total	C	H	N	O	0	3	0
			327	115	168	23	21			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	11	TYR	TRP	MICROHETEROGENEITY	NOR NOR00243

- Molecule 2 is a protein called GRAMICIDIN A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	16	Total	C	H	N	O	0	4	0
			329	113	175	22	19			

- Molecule 3 is METHANOL (three-letter code: MOH) (formula: CH<sub>4</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 6	C 1	H 4	O 1	0	0
3	A	1	Total 6	C 1	H 4	O 1	0	0
3	A	1	Total 6	C 1	H 4	O 1	0	0
3	A	1	Total 6	C 1	H 4	O 1	0	0
3	A	1	Total 6	C 1	H 4	O 1	0	0
3	A	1	Total 12	C 2	H 8	O 2	0	1
3	A	1	Total 12	C 2	H 8	O 2	0	1
3	A	1	Total 12	C 2	H 8	O 2	0	1
3	A	1	Total 24	C 4	H 16	O 4	0	1
3	B	1	Total 12	C 2	H 8	O 2	0	1
3	B	1	Total 6	C 1	H 4	O 1	0	0
3	B	1	Total 6	C 1	H 4	O 1	0	0
3	B	1	Total 6	C 1	H 4	O 1	0	0
3	B	1	Total 6	C 1	H 4	O 1	0	0
3	B	1	Total 6	C 1	H 4	O 1	0	1
3	B	1	Total 12	C 2	H 8	O 2	0	1
3	B	1	Total 12	C 2	H 8	O 2	0	1
3	B	1	Total 12	C 2	H 8	O 2	0	1
3	B	1	Total 12	C 2	H 8	O 2	0	1
3	B	1	Total 12	C 2	H 8	O 2	0	1
3	B	1	Total 30	C 5	H 20	O 5	0	1


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.


- Molecule 1: GRAMICIDIN A

Chain A: 



- Molecule 2: GRAMICIDIN A

Chain B: 



## 4 Data and refinement statistics i

EDS failed to run properly - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	14.91Å 26.01Å 31.91Å 90.00° 92.03° 90.00°	Depositor
Resolution (Å)	100.00 – 1.20	Depositor
% Data completeness (in resolution range)	99.9 (100.00-1.20)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	32.69 (at 1.20Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.102 , 0.168	Depositor
Wilson B-factor (Å <sup>2</sup> )	7.2	Xtriage
Anisotropy	0.235	Xtriage
Estimated twinning fraction	0.055 for h,-k,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 7726 reflections	Xtriage
Total number of atoms	866	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.09 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.5489e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ETA, DVA, DLE, FVA, MOH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.74	0/96	1.47	0/122
2	B	0.83	0/78	1.43	0/100
All	All	0.79	0/174	1.45	0/222

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	159	168	0	17	2
2	B	154	175	0	23	0
3	A	30	60	0	25	1
3	B	40	80	0	23	1
All	All	383	483	0	50	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 58.

The worst 5 of 50 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:A:505:MOH:H1	3:B:511[A]:MOH:H2	1.44	0.98
1:A:9:TRP:HE1	3:A:515[A]:MOH:H1	1.31	0.95
3:A:501:MOH:H2	3:A:519[C]:MOH:H3	1.46	0.93
3:A:505:MOH:H1	3:B:511[A]:MOH:C	2.01	0.89
2:B:6[A]:DVA:CG1	3:B:510:MOH:H2	2.08	0.84

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:11[A]:TYR:OH	3:A:515[A]:MOH:O[2.745]	2.10	0.10
1:A:16[A]:ETA:O	3:B:513[A]:MOH:C[1.556]	2.19	0.01

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	8/17 (47%)	5 (62%)	3 (38%)	0	100	100
2	B	8/16 (50%)	4 (50%)	3 (38%)	1 (12%)	1	0
All	All	16/33 (48%)	9 (56%)	6 (38%)	1 (6%)	2	0

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	7	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	7/6 (117%)	7 (100%)	0	100	100
2	B	5/5 (100%)	5 (100%)	0	100	100
All	All	12/11 (109%)	12 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

23 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	FVA	A	1	1	8,8,9	7.13	1 (12%)	6,9,11	1.04	0
1	DLE	A	10	1	7,7,8	6.66	1 (14%)	6,8,10	2.49	1 (16%)
1	DLE	A	12	1	7,7,8	7.95	2 (28%)	6,8,10	1.14	1 (16%)
1	DLE	A	14	1	7,7,8	6.88	2 (28%)	6,8,10	0.81	0
1	ETA	A	16[A]	1	3,3,3	0.56	0	2,2,2	0.70	0
1	ETA	A	16[B]	1	3,3,3	0.47	0	2,2,2	0.77	0
1	ETA	A	16[C]	1	3,3,3	0.49	0	2,2,2	0.12	0
1	DLE	A	4	1	7,7,8	7.19	2 (28%)	6,8,10	2.72	1 (16%)
1	DVA	A	6	1	6,6,7	8.51	2 (33%)	5,7,9	0.94	0
1	DVA	A	8	1	6,6,7	6.27	2 (33%)	5,7,9	0.72	0
2	FVA	B	1	2	8,8,9	6.60	2 (25%)	6,9,11	0.53	0
2	DLE	B	10[A]	-	2,3,8	0.94	0	1,3,10	0.52	0
2	DLE	B	10[B]	-	2,3,8	0.85	0	1,3,10	0.36	0
2	DLE	B	12	2	7,7,8	6.45	1 (14%)	6,8,10	0.65	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	DLE	B	14	2	7,7,8	6.14	1 (14%)	6,8,10	1.02	1 (16%)
2	ETA	B	16[A]	2	3,3,3	0.34	0	2,2,2	0.47	0
2	ETA	B	16[B]	2	3,3,3	0.37	0	2,2,2	0.72	0
2	ETA	B	16[C]	2	3,3,3	0.54	0	2,2,2	0.93	0
2	DLE	B	4	2	7,7,8	7.08	1 (14%)	6,8,10	2.03	1 (16%)
2	DVA	B	6[A]	-	2,2,7	1.77	1 (50%)	1,1,9	0.14	0
2	DVA	B	6[B]	-	2,2,7	2.14	2 (100%)	1,1,9	1.23	0
2	DVA	B	8[A]	-	2,2,7	2.02	1 (50%)	1,1,9	1.41	0
2	DVA	B	8[B]	-	2,2,7	2.51	1 (50%)	1,1,9	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	FVA	A	1	1	-	0/7/9/11	0/0/0/0
1	DLE	A	10	1	-	0/4/6/8	0/0/0/0
1	DLE	A	12	1	-	0/4/6/8	0/0/0/0
1	DLE	A	14	1	-	0/4/6/8	0/0/0/0
1	ETA	A	16[A]	1	-	0/1/1/1	0/0/0/0
1	ETA	A	16[B]	1	-	0/1/1/1	0/0/0/0
1	ETA	A	16[C]	1	-	0/1/1/1	0/0/0/0
1	DLE	A	4	1	-	0/4/6/8	0/0/0/0
1	DVA	A	6	1	-	0/4/6/8	0/0/0/0
1	DVA	A	8	1	-	0/4/6/8	0/0/0/0
2	FVA	B	1	2	-	0/7/9/11	0/0/0/0
2	DLE	B	10[A]	-	-	0/0/0/8	0/0/0/0
2	DLE	B	10[B]	-	-	0/0/0/8	0/0/0/0
2	DLE	B	12	2	-	0/4/6/8	0/0/0/0
2	DLE	B	14	2	-	0/4/6/8	0/0/0/0
2	ETA	B	16[A]	2	-	0/1/1/1	0/0/0/0
2	ETA	B	16[B]	2	-	0/1/1/1	0/0/0/0
2	ETA	B	16[C]	2	-	0/1/1/1	0/0/0/0
2	DLE	B	4	2	-	0/4/6/8	0/0/0/0
2	DVA	B	6[A]	-	-	0/0/0/8	0/0/0/0
2	DVA	B	6[B]	-	-	0/0/0/8	0/0/0/0
2	DVA	B	8[A]	-	-	0/0/0/8	0/0/0/0
2	DVA	B	8[B]	-	-	0/0/0/8	0/0/0/0

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	12	DLE	O-C	20.76	1.25	1.11
1	A	6	DVA	O-C	20.61	1.25	1.11
1	A	1	FVA	O-C	20.02	1.25	1.11
1	A	4	DLE	O-C	18.85	1.24	1.11
2	B	4	DLE	O-C	18.66	1.24	1.11

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4	DLE	C-CA-N	-6.56	107.27	113.83
1	A	10	DLE	C-CA-N	-5.97	107.86	113.83
2	B	4	DLE	C-CA-N	-4.83	109.01	113.83
1	A	12	DLE	C-CA-N	-2.72	111.11	113.83
2	B	14	DLE	C-CA-N	-2.33	111.50	113.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

35 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	MOH	A	501	-	1,1,1	0.37	0	0,0,0	0.00	-
3	MOH	A	502	-	1,1,1	0.48	0	0,0,0	0.00	-
3	MOH	A	505	-	1,1,1	0.32	0	0,0,0	0.00	-
3	MOH	A	506	-	1,1,1	0.35	0	0,0,0	0.00	-
3	MOH	A	508	-	1,1,1	0.29	0	0,0,0	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	MOH	A	515[A]	-	1,1,1	0.26	0	0,0,0	0.00	-
3	MOH	A	515[B]	-	1,1,1	0.47	0	0,0,0	0.00	-
3	MOH	A	516[A]	-	1,1,1	0.01	0	0,0,0	0.00	-
3	MOH	A	516[C]	-	1,1,1	0.32	0	0,0,0	0.00	-
3	MOH	A	518[A]	-	1,1,1	0.30	0	0,0,0	0.00	-
3	MOH	A	518[B]	-	1,1,1	0.28	0	0,0,0	0.00	-
3	MOH	A	519[A]	-	1,1,1	0.28	0	0,0,0	0.00	-
3	MOH	A	519[B]	-	1,1,1	0.24	0	0,0,0	0.00	-
3	MOH	A	519[C]	-	1,1,1	0.38	0	0,0,0	0.00	-
3	MOH	A	519[D]	-	1,1,1	0.29	0	0,0,0	0.00	-
3	MOH	B	503[A]	-	1,1,1	0.32	0	0,0,0	0.00	-
3	MOH	B	503[B]	-	1,1,1	0.23	0	0,0,0	0.00	-
3	MOH	B	504	-	1,1,1	0.33	0	0,0,0	0.00	-
3	MOH	B	507	-	1,1,1	0.17	0	0,0,0	0.00	-
3	MOH	B	509	-	1,1,1	0.35	0	0,0,0	0.00	-
3	MOH	B	510	-	1,1,1	0.10	0	0,0,0	0.00	-
3	MOH	B	511[A]	-	1,1,1	0.28	0	0,0,0	0.00	-
3	MOH	B	512[A]	-	1,1,1	0.24	0	0,0,0	0.00	-
3	MOH	B	512[B]	-	1,1,1	0.20	0	0,0,0	0.00	-
3	MOH	B	513[A]	-	1,1,1	0.26	0	0,0,0	0.00	-
3	MOH	B	513[B]	-	1,1,1	0.25	0	0,0,0	0.00	-
3	MOH	B	514[A]	-	1,1,1	0.31	0	0,0,0	0.00	-
3	MOH	B	514[B]	-	1,1,1	0.19	0	0,0,0	0.00	-
3	MOH	B	517[A]	-	1,1,1	0.17	0	0,0,0	0.00	-
3	MOH	B	517[B]	-	1,1,1	0.34	0	0,0,0	0.00	-
3	MOH	B	520[A]	-	1,1,1	0.26	0	0,0,0	0.00	-
3	MOH	B	520[B]	-	1,1,1	0.24	0	0,0,0	0.00	-
3	MOH	B	520[C]	-	1,1,1	0.16	0	0,0,0	0.00	-
3	MOH	B	520[D]	-	1,1,1	0.09	0	0,0,0	0.00	-
3	MOH	B	520[E]	-	1,1,1	0.24	0	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MOH	A	501	-	-	0/0/0/0	0/0/0/0
3	MOH	A	502	-	-	0/0/0/0	0/0/0/0
3	MOH	A	505	-	-	0/0/0/0	0/0/0/0
3	MOH	A	506	-	-	0/0/0/0	0/0/0/0
3	MOH	A	508	-	-	0/0/0/0	0/0/0/0
3	MOH	A	515[A]	-	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MOH	A	515[B]	-	-	0/0/0/0	0/0/0/0
3	MOH	A	516[A]	-	-	0/0/0/0	0/0/0/0
3	MOH	A	516[C]	-	-	0/0/0/0	0/0/0/0
3	MOH	A	518[A]	-	-	0/0/0/0	0/0/0/0
3	MOH	A	518[B]	-	-	0/0/0/0	0/0/0/0
3	MOH	A	519[A]	-	-	0/0/0/0	0/0/0/0
3	MOH	A	519[B]	-	-	0/0/0/0	0/0/0/0
3	MOH	A	519[C]	-	-	0/0/0/0	0/0/0/0
3	MOH	A	519[D]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	503[A]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	503[B]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	504	-	-	0/0/0/0	0/0/0/0
3	MOH	B	507	-	-	0/0/0/0	0/0/0/0
3	MOH	B	509	-	-	0/0/0/0	0/0/0/0
3	MOH	B	510	-	-	0/0/0/0	0/0/0/0
3	MOH	B	511[A]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	512[A]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	512[B]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	513[A]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	513[B]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	514[A]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	514[B]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	517[A]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	517[B]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	520[A]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	520[B]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	520[C]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	520[D]	-	-	0/0/0/0	0/0/0/0
3	MOH	B	520[E]	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS failed to run properly - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS failed to run properly - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS failed to run properly - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS failed to run properly - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS failed to run properly - this section will therefore be empty.