



# wwPDB X-ray Structure Validation Summary Report i

Mar 1, 2014 – 03:28 AM GMT

PDB ID : 2AN6  
Title : Protein-peptide complex  
Authors : House, C.M.; Hancock, N.C.; Moller, A.; Cromer, B.A.; Fedorov, V.; Bowtell, D.D.L.; Parker, M.W.; Polekhina, G.  
Deposited on : 2005-08-11  
Resolution : 3.00 Å (reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	191	
1	B	191	
1	C	191	
1	D	191	
2	E	24	
2	F	24	
2	G	24	
2	H	24	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	ZN	A	601	-	X

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6372 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin ligase SIAH1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	191	1506	953	258	279	16	0	0	0
1	B	191	1506	953	258	279	16	0	0	0
1	C	191	1506	953	258	279	16	0	0	0
1	D	191	1506	953	258	279	16	0	0	0

- Molecule 2 is a protein called peptide from Phyllopod.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	11	85	55	17	12	1	0	0	0
2	F	11	85	55	17	12	1	0	0	0
2	G	11	85	55	17	12	1	0	0	0
2	H	11	85	55	17	12	1	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Zn	0	0
			2	2		
3	A	2	Total	Zn	0	0
			2	2		
3	D	2	Total	Zn	0	0
			2	2		
3	C	2	Total	Zn	0	0
			2	2		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ubiquitin ligase SIAH1A

Chain A:




- Molecule 1: Ubiquitin ligase SIAH1A

Chain B:



- Molecule 1: Ubiquitin ligase SIAH1A


Chain C:



- Molecule 1: Ubiquitin ligase SIAH1A

Chain D:





- Molecule 2: peptide from Phyllopod

Chain E:



- Molecule 2: peptide from Phyllopod

Chain F:



- Molecule 2: peptide from Phyllopod

Chain G:



- Molecule 2: peptide from Phyllopod

Chain H:



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.30Å 100.00Å 103.40Å 90.00° 104.30° 90.00°	Depositor
Resolution (Å)	30.00 – 3.00 100.20 – 2.99	Depositor EDS
% Data completeness (in resolution range)	99.0 (30.00-3.00) 98.5 (100.20-2.99)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.00 (at 3.01Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.237 , 0.305 0.235 , 0.299	Depositor DCC
$R_{free}$ test set	1266 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	73.8	Xtrriage
Anisotropy	0.638	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 47.9	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Outliers	0 of 25343 reflections	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6372	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.81	0/1543	0.88	1/2088 (0.0%)
1	B	0.79	1/1543 (0.1%)	0.86	2/2088 (0.1%)
1	C	0.83	0/1543	0.86	0/2088
1	D	0.79	2/1543 (0.1%)	0.86	5/2088 (0.2%)
2	E	0.82	0/86	0.99	0/117
2	F	0.71	0/86	0.83	0/117
2	G	0.82	0/86	0.84	0/117
2	H	0.94	0/86	0.96	0/117
All	All	0.81	3/6516 (0.0%)	0.87	8/8820 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
All	All	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	236	TRP	CE3-CZ3	7.52	1.51	1.38
1	D	165	PHE	CE2-CZ	5.13	1.47	1.37
1	B	236	TRP	CA-CB	5.11	1.65	1.53

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	225	LEU	CB-CG-CD1	-6.52	99.92	111.00
1	D	148	LEU	CA-CB-CG	6.05	129.22	115.30
1	B	225	LEU	CB-CG-CD1	-5.81	101.13	111.00
1	D	191	LEU	CA-CB-CG	5.66	128.31	115.30
1	D	225	LEU	CA-CB-CG	5.61	128.19	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	102	SER	Peptide
1	D	191	LEU	Peptide

## 5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1506	0	1451	87	0
1	B	1506	0	1454	91	0
1	C	1506	0	1451	75	0
1	D	1506	0	1454	108	0
2	E	85	0	98	5	0
2	F	85	0	98	7	0
2	G	85	0	98	3	0
2	H	85	0	98	6	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	1	0
All	All	6372	0	6202	346	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 28.

The worst 5 of 346 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:212:ILE:HA	1:C:243:ILE:HD11	1.36	1.08
1:B:168:THR:HG22	2:F:120:VAL:O	1.56	1.06
1:C:107:ILE:H	1:C:107:ILE:HD13	1.20	1.04
1:B:162:ASP:OD1	1:B:278:THR:HG22	1.58	1.00
1:A:254:SER:O	1:B:263:ILE:HD12	1.68	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	189/191 (99%)	159 (84%)	25 (13%)	5 (3%)	8	39
1	B	189/191 (99%)	160 (85%)	22 (12%)	7 (4%)	5	28
1	C	189/191 (99%)	157 (83%)	25 (13%)	7 (4%)	5	28
1	D	189/191 (99%)	153 (81%)	29 (15%)	7 (4%)	5	28
2	E	9/24 (38%)	8 (89%)	1 (11%)	0	100	100
2	F	9/24 (38%)	7 (78%)	2 (22%)	0	100	100
2	G	9/24 (38%)	8 (89%)	1 (11%)	0	100	100
2	H	9/24 (38%)	7 (78%)	0	2 (22%)	0	0
All	All	792/860 (92%)	659 (83%)	105 (13%)	28 (4%)	6	30

5 of 28 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	243	ILE
1	B	244	HIS
1	B	245	GLU
1	B	256	CYS
1	C	155	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/165 (100%)	132 (80%)	33 (20%)	2	10
1	B	165/165 (100%)	135 (82%)	30 (18%)	2	13
1	C	165/165 (100%)	141 (86%)	24 (14%)	5	22
1	D	165/165 (100%)	133 (81%)	32 (19%)	2	11
2	E	10/23 (44%)	8 (80%)	2 (20%)	2	10
2	F	10/23 (44%)	9 (90%)	1 (10%)	11	39
2	G	10/23 (44%)	8 (80%)	2 (20%)	2	10
2	H	10/23 (44%)	7 (70%)	3 (30%)	0	2
All	All	700/752 (93%)	573 (82%)	127 (18%)	2	13

5 of 127 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	254	SER
1	C	169	ASP
1	D	265	GLN
1	B	261	THR
1	C	100	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	171	ASN
1	C	228	ASN
1	D	253	ASN
1	C	182	GLN
1	C	253	ASN

### 5.3.3 RNA [i](#)

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	191/191 (100%)	-0.13	0 <a href="#">100</a> <a href="#">100</a>	13, 66, 99, 133	0
1	B	191/191 (100%)	-0.14	0 <a href="#">100</a> <a href="#">100</a>	14, 71, 112, 126	0
1	C	191/191 (100%)	-0.11	1 (0%) <a href="#">88</a> <a href="#">36</a>	14, 72, 109, 148	0
1	D	191/191 (100%)	-0.10	0 <a href="#">100</a> <a href="#">100</a>	25, 83, 137, 143	0
2	E	11/24 (45%)	0.07	0 <a href="#">100</a> <a href="#">100</a>	48, 56, 69, 75	0
2	F	11/24 (45%)	0.05	0 <a href="#">100</a> <a href="#">100</a>	42, 68, 92, 93	0
2	G	11/24 (45%)	0.06	0 <a href="#">100</a> <a href="#">100</a>	52, 71, 94, 101	0
2	H	11/24 (45%)	0.03	0 <a href="#">100</a> <a href="#">100</a>	65, 77, 95, 107	0
All	All	808/860 (93%)	-0.11	1 (0%) <a href="#">93</a> <a href="#">63</a>	13, 72, 126, 148	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	201	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ZN	A	601	1/1	0.19	3.21	80,80,80,80	0
3	ZN	A	602	1/1	0.19	0.79	78,78,78,78	0
3	ZN	B	604	1/1	0.20	0.16	97,97,97,97	0
3	ZN	C	605	1/1	0.16	0.04	78,78,78,78	0
3	ZN	B	603	1/1	0.16	-0.09	89,89,89,89	0
3	ZN	D	607	1/1	0.17	-0.21	144,144,144,144	0
3	ZN	C	606	1/1	0.14	-0.55	95,95,95,95	0
3	ZN	D	608	1/1	0.13	-0.82	115,115,115,115	0

## 6.5 Other polymers i

There are no such residues in this entry.