



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 04:59 AM GMT

PDB ID : 3ANY
Title : Crystal structure of ethanolamine ammonia-lyase from escherichia coli complexed with CN-CBL and (R)-2-amino-1-propanol
Authors : Shibata, N.
Deposited on : 2010-09-16
Resolution : 2.10 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	453	
1	C	453	
2	B	263	
2	D	263	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11700 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ethanolamine ammonia-lyase heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	453	Total	C	N	O	S	0	2	0
			3470	2174	593	681	22			
1	C	453	Total	C	N	O	S	0	6	0
			3495	2192	599	682	22			

- Molecule 2 is a protein called Ethanolamine ammonia-lyase light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	252	Total	C	N	O	S	0	0	0
			1916	1197	347	362	10			
2	D	252	Total	C	N	O	S	0	0	0
			1916	1197	347	362	10			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	33	MET	-	EXPRESSION TAG	UNP P19636
B	34	ASP	-	EXPRESSION TAG	UNP P19636
B	35	GLN	-	EXPRESSION TAG	UNP P19636
B	36	SER	-	EXPRESSION TAG	UNP P19636
B	37	SER	-	EXPRESSION TAG	UNP P19636
B	38	HIS	-	EXPRESSION TAG	UNP P19636
B	39	HIS	-	EXPRESSION TAG	UNP P19636
B	40	HIS	-	EXPRESSION TAG	UNP P19636
B	41	HIS	-	EXPRESSION TAG	UNP P19636
B	42	HIS	-	EXPRESSION TAG	UNP P19636
B	43	HIS	-	EXPRESSION TAG	UNP P19636
D	33	MET	-	EXPRESSION TAG	UNP P19636
D	34	ASP	-	EXPRESSION TAG	UNP P19636
D	35	GLN	-	EXPRESSION TAG	UNP P19636
D	36	SER	-	EXPRESSION TAG	UNP P19636
D	37	SER	-	EXPRESSION TAG	UNP P19636

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	38	HIS	-	EXPRESSION TAG	UNP P19636
D	39	HIS	-	EXPRESSION TAG	UNP P19636
D	40	HIS	-	EXPRESSION TAG	UNP P19636
D	41	HIS	-	EXPRESSION TAG	UNP P19636
D	42	HIS	-	EXPRESSION TAG	UNP P19636
D	43	HIS	-	EXPRESSION TAG	UNP P19636

- Molecule 3 is (2R)-2-AMINOPROPAN-1-OL (three-letter code: 2A3) (formula: C₃H₉NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			5	3	1	1		
3	C	1	Total	C	N	O	0	0
			5	3	1	1		

- Molecule 4 is COBALAMIN (three-letter code: B12) (formula: C₆₂H₈₉CoN₁₃O₁₄P).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	B	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		
4	D	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	274	Total	O	0	0
			274	274		
5	B	75	Total	O	0	0
			75	75		
5	C	272	Total	O	0	0
			272	272		
5	D	90	Total	O	0	0
			90	90		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ethanolamine ammonia-lyase heavy chain

Chain A: 




- Molecule 1: Ethanolamine ammonia-lyase heavy chain

Chain C: 




- Molecule 2: Ethanolamine ammonia-lyase light chain

Chain B: 



- Molecule 2: Ethanolamine ammonia-lyase light chain

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	244.07Å 244.07Å 76.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.84 – 2.10 47.82 – 2.10	Depositor EDS
% Data completeness (in resolution range)	92.8 (47.84-2.10) 92.8 (47.82-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.88 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.246 , 0.276 0.242 , 0.252	Depositor DCC
R_{free} test set	7043 reflections (5.22%)	DCC
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 16.9	EDS
Estimated twinning fraction	0.059 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Outliers	0 of 141886 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	11700	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B12, 2A3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.56	0/3530	0.34	0/4780
1	C	0.57	0/3567	0.35	0/4829
2	B	0.53	0/1943	0.34	0/2633
2	D	0.53	0/1943	0.34	0/2633
All	All	0.55	0/10983	0.34	0/14875

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	C	0	1
2	B	0	2
2	D	0	2
All	All	0	7

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	326	VAL	Peptide
1	A	423	PRO	Peptide
2	B	120	LEU	Peptide
2	B	206	ARG	Peptide
1	C	326	VAL	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	D	110	VAL	Peptide
2	D	206	ARG	Peptide

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3470	0	0	2	0
1	C	3495	0	0	3	0
2	B	1916	0	0	3	0
2	D	1916	0	0	2	0
3	A	5	0	9	0	0
3	C	5	0	9	0	0
4	B	91	0	88	10	0
4	D	91	0	88	12	0
5	A	274	0	0	0	0
5	B	75	0	0	0	0
5	C	272	0	0	0	0
5	D	90	0	0	0	0
All	All	11700	0	194	29	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (29) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:B:601:B12:H531	4:B:601:B12:H552	1.58	0.86
4:B:601:B12:H362	4:B:601:B12:H351	1.56	0.85
4:D:601:B12:H552	4:D:601:B12:H531	1.63	0.81
4:D:601:B12:H351	4:D:601:B12:H362	1.67	0.77
1:C:329:PHE:O	2:D:229:ARG:NH2	2.20	0.75
1:C:269[B]:ARG:NH1	1:C:269[B]:ARG:CG	2.50	0.73
4:D:601:B12:H601	4:D:601:B12:H262	1.87	0.55
4:B:601:B12:H262	4:B:601:B12:H601	1.89	0.53
4:D:601:B12:H552	4:D:601:B12:C53	2.36	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:B:601:B12:H482	4:B:601:B12:H2B	1.89	0.53
4:B:601:B12:C35	4:B:601:B12:H362	2.32	0.53
2:B:229:ARG:NH1	4:B:601:B12:H543	2.25	0.52
4:B:601:B12:C53	4:B:601:B12:H552	2.36	0.49
4:D:601:B12:H362	4:D:601:B12:C35	2.40	0.49
2:B:110:VAL:O	2:B:203:ARG:NH2	2.45	0.49
4:D:601:B12:H2B	4:D:601:B12:H482	1.93	0.49
1:C:187:ASP:N	1:C:187:ASP:OD1	2.46	0.48
2:D:110:VAL:O	2:D:203:ARG:NH2	2.46	0.48
4:D:601:B12:H602	4:D:601:B12:H541	1.78	0.47
4:B:601:B12:H481	4:B:601:B12:H473	1.52	0.46
4:D:601:B12:H473	4:D:601:B12:H481	1.63	0.45
4:B:601:B12:H261	4:B:601:B12:H91	1.92	0.44
4:D:601:B12:H10	4:D:601:B12:H472	1.79	0.44
1:A:187:ASP:N	1:A:187:ASP:OD1	2.51	0.44
4:D:601:B12:H301	4:D:601:B12:H253	1.79	0.43
1:A:443:THR:O	2:B:82:ARG:NH1	2.52	0.42
4:D:601:B12:H91	4:D:601:B12:H261	1.93	0.41
4:B:601:B12:H481	4:B:601:B12:H521	1.45	0.41
4:D:601:B12:H562	4:D:601:B12:H18	1.86	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	453/453 (100%)	436 (96%)	17 (4%)	0	100	100
1	C	457/453 (101%)	438 (96%)	19 (4%)	0	100	100
2	B	250/263 (95%)	243 (97%)	7 (3%)	0	100	100
2	D	250/263 (95%)	243 (97%)	7 (3%)	0	100	100
All	All	1410/1432 (98%)	1360 (96%)	50 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	372/370 (100%)	352 (95%)	20 (5%)	31	27
1	C	376/370 (102%)	357 (95%)	19 (5%)	33	29
2	B	206/217 (95%)	187 (91%)	19 (9%)	13	8
2	D	206/217 (95%)	182 (88%)	24 (12%)	8	4
All	All	1160/1174 (99%)	1078 (93%)	82 (7%)	21	16

All (82) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	A	15	LYS
1	A	16	ASP
1	A	27	LEU
1	A	60	ARG
1	A	65	ILE
1	A	110	ILE
1	A	140	LYS
1	A	161	LEU
1	A	166	THR
1	A	171	GLN
1	A	196	THR
1	A	197	ASP
1	A	217	ASN
1	A	262	LEU
1	A	269	ARG
1	A	333	GLU
1	A	380	MET
1	A	401	MET
1	A	451	LEU
2	B	70	LEU
2	B	74	THR
2	B	77	ARG
2	B	96	LEU
2	B	107	LEU
2	B	108	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	118	GLN
2	B	121	LEU
2	B	123	VAL
2	B	152	LYS
2	B	182	LEU
2	B	183	PRO
2	B	190	LYS
2	B	194	LEU
2	B	196	VAL
2	B	220	LYS
2	B	229	ARG
2	B	234	GLN
2	B	245	SER
1	C	7	LEU
1	C	27	LEU
1	C	41	SER
1	C	60	ARG
1	C	75	LEU
1	C	161	LEU
1	C	166	THR
1	C	167	ARG
1	C	202	LEU
1	C	221	GLN
1	C	229	THR
1	C	230	THR
1	C	253	LYS
1	C	294	LEU
1	C	333	GLU
1	C	401	MET
1	C	405	GLN
1	C	450	SER
1	C	451	LEU
2	D	107	LEU
2	D	110	VAL
2	D	112	GLU
2	D	113	GLU
2	D	121	LEU
2	D	123	VAL
2	D	124	ARG
2	D	129	ASP
2	D	132	LEU
2	D	134	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	141	ARG
2	D	161	VAL
2	D	182	LEU
2	D	195	LYS
2	D	206	ARG
2	D	220	LYS
2	D	223	ILE
2	D	229	ARG
2	D	232	LEU
2	D	234	GLN
2	D	243	VAL
2	D	259	ILE
2	D	267	THR
2	D	291	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	2A3	A	602	-	4,4,4	0.45	0	4,4,4	0.64	0
4	B12	B	601	-	101,101,101	7.23	14 (13%)	152,166,166	1.16	12 (7%)
3	2A3	C	602	-	4,4,4	0.50	0	4,4,4	0.58	0
4	B12	D	601	-	101,101,101	7.17	15 (14%)	152,166,166	1.15	13 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2A3	A	602	-	-	0/2/2/2	0/0/0/0
4	B12	B	601	-	-	0/56/223/223	0/1/11/11
3	2A3	C	602	-	-	0/2/2/2	0/0/0/0
4	B12	D	601	-	-	0/56/223/223	0/1/11/11

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	601	B12	CO-N24	52.48	1.90	1.79
4	D	601	B12	CO-N21	50.05	1.89	1.79
4	D	601	B12	CO-N24	49.75	1.89	1.79
4	B	601	B12	CO-N21	48.06	1.89	1.79
4	B	601	B12	C19-N24	-6.42	1.39	1.48
4	D	601	B12	C19-N24	-6.22	1.39	1.48
4	D	601	B12	C11-C10	-6.12	1.30	1.41
4	B	601	B12	C11-C10	-5.52	1.31	1.41
4	B	601	B12	C8B-N1B	-4.32	1.33	1.39
4	D	601	B12	C8B-N1B	-4.14	1.34	1.39
4	B	601	B12	C47-C12	-3.93	1.46	1.54
4	D	601	B12	C46-C12	-3.92	1.46	1.54
4	B	601	B12	C46-C12	-3.40	1.47	1.54
4	D	601	B12	C8B-C9B	3.15	1.46	1.40
4	D	601	B12	C6B-C5B	3.07	1.49	1.40
4	B	601	B12	C6B-C5B	3.04	1.49	1.40
4	D	601	B12	C12-C11	-3.03	1.44	1.52
4	B	601	B12	C8B-C9B	2.81	1.46	1.40
4	B	601	B12	C4-N21	2.73	1.38	1.31
4	B	601	B12	C14-C15	2.28	1.48	1.41
4	D	601	B12	C16-N24	2.27	1.37	1.31
4	B	601	B12	C16-N24	2.27	1.37	1.31
4	B	601	B12	C6-N22	2.21	1.43	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	601	B12	C41-C8	2.20	1.57	1.53
4	D	601	B12	C6-N22	2.19	1.43	1.34
4	D	601	B12	C14-C15	2.15	1.47	1.41
4	B	601	B12	C17-C16	2.07	1.57	1.52
4	D	601	B12	O6R-C1R	2.06	1.44	1.41
4	D	601	B12	C17-C16	2.03	1.57	1.52

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	601	B12	C19-N24-C16	6.25	112.42	109.08
4	B	601	B12	C19-N24-C16	6.14	112.36	109.08
4	B	601	B12	C9-C10-C11	-3.74	122.78	132.53
4	B	601	B12	C1-C19-N24	3.57	110.49	106.27
4	D	601	B12	C9-C10-C11	-3.39	123.70	132.53
4	D	601	B12	C2-C1-N21	3.26	102.71	101.05
4	B	601	B12	C2-C1-N21	2.82	102.49	101.05
4	B	601	B12	C30-C3-C2	-2.75	113.39	119.04
4	D	601	B12	C54-C17-C18	-2.69	108.57	112.91
4	D	601	B12	C30-C3-C2	-2.66	113.57	119.04
4	B	601	B12	C54-C17-C18	-2.46	108.94	112.91
4	D	601	B12	C7-C8-C9	2.27	103.45	100.77
4	B	601	B12	C7-C8-C9	2.24	103.42	100.77
4	B	601	B12	O6R-C1R-N1B	2.23	110.52	108.44
4	D	601	B12	C3-C4-C5	-2.21	123.21	131.83
4	B	601	B12	C25-C2-C1	-2.19	110.60	113.80
4	D	601	B12	C1-C19-N24	2.19	108.85	106.27
4	D	601	B12	C19-C1-N21	2.17	104.47	102.11
4	B	601	B12	C13-C14-C15	-2.14	123.46	131.83
4	D	601	B12	C13-C14-C15	-2.10	123.63	131.83
4	D	601	B12	C20-C1-C19	-2.09	107.24	109.36
4	B	601	B12	C19-C1-N21	2.07	104.36	102.11
4	D	601	B12	C25-C2-C1	-2.04	110.81	113.80
4	D	601	B12	O6R-C1R-N1B	2.04	110.33	108.44
4	B	601	B12	C48-C13-C12	-2.01	110.76	116.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	453/453 (100%)	0.17	1 (0%) 93 95	18, 31, 49, 59	0
1	C	453/453 (100%)	0.15	1 (0%) 93 95	16, 31, 54, 67	0
2	B	252/263 (95%)	0.75	22 (8%) 10 11	26, 60, 85, 98	0
2	D	252/263 (95%)	1.25	58 (23%) 1 1	21, 72, 110, 122	0
All	All	1410/1432 (98%)	0.46	82 (5%) 22 25	16, 39, 90, 122	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	161	VAL	7.1
2	D	123	VAL	6.0
2	D	148	VAL	5.8
2	D	153	ALA	5.7
2	D	200	PHE	5.5
2	D	145	ALA	5.2
2	D	111	PRO	4.6
2	D	216	ILE	4.5
2	D	187	ALA	4.5
2	D	192	ALA	4.5
2	D	156	VAL	4.3
2	D	202	VAL	4.2
2	D	119	GLY	4.1
2	D	191	GLN	4.0
2	B	270	VAL	3.7
2	B	249	ALA	3.6
2	D	190	LYS	3.6
2	B	191	GLN	3.5
2	D	174	ILE	3.4
2	B	192	ALA	3.3
2	D	194	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	189	LEU	3.3
2	D	165	ILE	3.2
2	D	273	ALA	3.2
2	D	196	VAL	3.2
2	B	72	ARG	3.1
2	D	244	TYR	3.1
2	D	107	LEU	3.1
2	B	113	GLU	3.0
2	D	181	ILE	3.0
2	D	157	ALA	2.9
2	D	164	VAL	2.9
2	D	121	LEU	2.9
2	D	155	CYS	2.8
2	D	225	LEU	2.8
2	D	158	ASN	2.8
2	D	246	PRO	2.8
2	B	44	ALA	2.7
2	D	220	LYS	2.6
2	D	295	ARG	2.6
2	D	146	GLU	2.6
2	B	56	ILE	2.6
2	B	183	PRO	2.6
2	D	129	ASP	2.5
2	D	178	TYR	2.5
2	D	213	ILE	2.5
2	D	242	ALA	2.5
2	D	212	GLN	2.5
2	D	223	ILE	2.4
2	D	224	LEU	2.4
2	B	156	VAL	2.4
2	B	265	GLY	2.4
2	B	194	LEU	2.4
2	D	151	LEU	2.4
2	D	69	GLU	2.4
2	D	127	ILE	2.4
2	D	59	GLU	2.4
2	D	209	ILE	2.3
2	D	143	LEU	2.3
2	B	295	ARG	2.3
2	B	179	GLU	2.3
2	D	162	GLN	2.3
2	D	179	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	120	LEU	2.3
2	B	178	TYR	2.2
2	D	141	ARG	2.2
2	B	201	PHE	2.2
1	C	64	VAL	2.2
2	D	132	LEU	2.2
2	D	270	VAL	2.2
2	B	239	SER	2.2
2	D	258	CYS	2.2
2	D	183	PRO	2.2
2	D	206	ARG	2.1
2	B	234	GLN	2.1
1	A	336	TYR	2.1
2	B	200	PHE	2.1
2	B	225	LEU	2.0
2	D	250	THR	2.0
2	D	286	GLN	2.0
2	B	182	LEU	2.0
2	D	182	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	2A3	A	602	5/5	0.15	0.70	24,24,30,30	0
4	B12	B	601	91/91	0.15	0.22	28,35,45,53	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	B12	D	601	91/91	0.17	0.18	29,44,52,59	0
3	2A3	C	602	5/5	0.11	-0.48	23,26,29,29	0

6.5 Other polymers ⓘ

There are no such residues in this entry.