



wwPDB X-ray Structure Validation Summary Report (i)

Feb 27, 2014 – 12:45 AM GMT

PDB ID : 1AP2
Title : SINGLE CHAIN FV OF C219
Authors : Hoedemaeker, P.J.; Rose, D.R.
Deposited on : 1997-07-23
Resolution : 2.36 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance





The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1064 (2.38-2.34)
Ramachandran outliers	78287	1048 (2.38-2.34)
Sidechain outliers	78261	1049 (2.38-2.34)
RSRZ outliers	66119	3330 (2.40-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	113	
1	C	113	
2	B	123	
2	D	123	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3720 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MONOCLONAL ANTIBODY C219.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	112	Total	C	N	O	S	0	0	0
			859	541	138	176	4			
1	C	113	Total	C	N	O	S	0	0	0
			864	543	139	178	4			

- Molecule 2 is a protein called MONOCLONAL ANTIBODY C219.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	120	Total	C	N	O	S	0	0	0
			939	592	157	187	3			
2	D	123	Total	C	N	O	S	0	0	0
			956	600	160	193	3			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	33	PHE	TYR	CONFLICT	EMBL Z22078
B	56	ASP	GLY	CONFLICT	EMBL Z22078
B	71	ILE	THR	CONFLICT	EMBL Z22078
B	100	GLU	-	INSERTION	EMBL Z22078
B	101	VAL	-	INSERTION	EMBL Z22078
B	102	TYR	ASP	CONFLICT	EMBL Z22078
B	103	SER	ASN	CONFLICT	EMBL Z22078
B	106	SER	-	INSERTION	EMBL Z22078
B	107	PRO	ALA	CONFLICT	EMBL Z22078
B	108	LEU	MET	CONFLICT	EMBL Z22078
B	110	VAL	TYR	CONFLICT	EMBL Z22078
B	113	ALA	GLN	CONFLICT	EMBL Z22078
B	116	THR	SER	CONFLICT	EMBL Z22078
B	120	PRO	SER	CONFLICT	EMBL Z22078
D	33	PHE	TYR	CONFLICT	EMBL Z22078
D	56	ASP	GLY	CONFLICT	EMBL Z22078

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	71	ILE	THR	CONFLICT	EMBL Z22078
D	100	GLU	-	INSERTION	EMBL Z22078
D	101	VAL	-	INSERTION	EMBL Z22078
D	102	TYR	ASP	CONFLICT	EMBL Z22078
D	103	SER	ASN	CONFLICT	EMBL Z22078
D	106	SER	-	INSERTION	EMBL Z22078
D	107	PRO	ALA	CONFLICT	EMBL Z22078
D	108	LEU	MET	CONFLICT	EMBL Z22078
D	110	VAL	TYR	CONFLICT	EMBL Z22078
D	113	ALA	GLN	CONFLICT	EMBL Z22078
D	116	THR	SER	CONFLICT	EMBL Z22078
D	120	PRO	SER	CONFLICT	EMBL Z22078


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	28	Total O 28 28	0	0
3	B	19	Total O 19 19	0	0
3	C	29	Total O 29 29	0	0
3	D	26	Total O 26 26	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: MONOCLONAL ANTIBODY C219




• Molecule 1: MONOCLONAL ANTIBODY C219



• Molecule 2: MONOCLONAL ANTIBODY C219



• Molecule 2: MONOCLONAL ANTIBODY C219



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.04Å 64.35Å 154.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.36 23.78 – 2.38	Depositor EDS
% Data completeness (in resolution range)	78.4 (6.00-2.36) 73.9 (23.78-2.38)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 2.39Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.204 , 0.284 0.216 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	14.4	Xtriage
Anisotropy	0.607	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	0 of 22598 reflections	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	3720	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/879	0.64	1/1197 (0.1%)
1	C	0.40	0/884	0.59	0/1202
2	B	0.31	0/961	0.52	0/1310
2	D	0.35	0/978	0.58	0/1331
All	All	0.37	0/3702	0.58	1/5040 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	89	LEU	CB-CG-CD2	-5.01	102.48	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	859	0	831	49	0
1	C	864	0	834	45	0
2	B	939	0	904	42	0
2	D	956	0	917	73	0
3	A	28	0	0	2	0
3	B	19	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	29	0	0	3	0
3	D	26	0	0	0	0
All	All	3720	0	3486	201	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 28.

The worst 5 of 201 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:3:VAL:H	1:A:26:SER:HB2	1.32	0.94
1:C:31:ASN:HB3	1:C:34:ASN:HD21	1.37	0.88
2:B:38:LYS:HB2	2:B:48:ILE:HD11	1.61	0.83
1:C:1:ASP:HB2	3:C:125:HOH:O	1.83	0.78
2:D:12:VAL:HG23	2:D:119:VAL:HG22	1.65	0.78

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	110/113 (97%)	99 (90%)	8 (7%)	3 (3%)	8	4
1	C	111/113 (98%)	100 (90%)	11 (10%)	0	100	100
2	B	118/123 (96%)	101 (86%)	13 (11%)	4 (3%)	6	3
2	D	121/123 (98%)	105 (87%)	12 (10%)	4 (3%)	6	3
All	All	460/472 (98%)	405 (88%)	44 (10%)	11 (2%)	9	7

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	42	GLU
2	B	43	GLN
2	D	14	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	74	GLY
2	D	120	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/98 (100%)	77 (79%)	21 (21%)	1	1
1	C	98/98 (100%)	70 (71%)	28 (29%)	0	0
2	B	102/104 (98%)	71 (70%)	31 (30%)	0	0
2	D	104/104 (100%)	68 (65%)	36 (35%)	0	0
All	All	402/404 (100%)	286 (71%)	116 (29%)	0	0

5 of 116 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	1	ASP
1	C	32	SER
2	D	83	LEU
1	C	9	SER
1	C	18	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	55	ASN
1	C	27	GLN
2	D	43	GLN
2	B	43	GLN
2	D	35	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	112/113 (99%)	-0.11	0 100 100	2, 7, 20, 25	0
1	C	113/113 (100%)	-0.13	2 (1%) 65 68	2, 6, 19, 35	0
2	B	120/123 (97%)	-0.06	1 (0%) 83 86	2, 7, 24, 34	0
2	D	123/123 (100%)	0.11	4 (3%) 44 48	2, 9, 33, 44	0
All	All	468/472 (99%)	-0.04	7 (1%) 70 74	2, 7, 24, 44	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	123	SER	5.0
2	D	122	GLY	4.3
2	D	14	PRO	3.0
1	C	73	SER	3.0
2	B	43	GLN	2.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.