



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 10:49 AM GMT

PDB ID : 2AP2
Title : SINGLE CHAIN FV OF C219 IN COMPLEX WITH SYNTHETIC EPITOPE PEPTIDE
Authors : Van Den Elsen, J.M.H.; Rose, D.R.
Deposited on : 1999-03-22
Resolution : 2.40 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2207 (2.40-2.40)
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)
RSRZ outliers	66119	2210 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	115	
1	C	115	
2	B	122	
2	D	122	
3	E	17	
3	F	17	
4	P	14	
4	Q	14	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3954 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (ANTIBODY (LIGHT CHAIN)).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	115	Total	C	N	O	S	0	0	0
			888	561	144	179	4			
1	C	115	Total	C	N	O	S	0	0	0
			888	561	144	179	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	PHE	-	SEE REMARK 999	GB 196573
A	-1	VAL	-	SEE REMARK 999	GB 196573
A	0	ARG	-	SEE REMARK 999	GB 196573
A	112	PRO	LEU	CONFLICT	GB 196573
C	-2	PHE	-	SEE REMARK 999	GB 196573
C	-1	VAL	-	SEE REMARK 999	GB 196573
C	0	ARG	-	SEE REMARK 999	GB 196573
C	112	PRO	LEU	CONFLICT	GB 196573

- Molecule 2 is a protein called PROTEIN (ANTIBODY (HEAVY CHAIN)).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	122	Total	C	N	O	S	0	0	0
			949	597	159	190	3			
2	D	122	Total	C	N	O	S	0	0	0
			949	597	159	190	3			

- Molecule 3 is a protein called PROTEIN (C-MYC TAG AND HIS TAG).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	2	Total	C	N	O	0	0	1
			7	3	2	2			
3	F	4	Total	C	N	O	0	0	1
			25	13	5	7			

- Molecule 4 is a protein called PROTEIN (P-GLYCOPROTEIN).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	P	12	Total	C	N	O	0	0	1
			74	45	13	16			
4	Q	14	Total	C	N	O	0	0	0
			110	65	22	23			


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	19	Total	O	0	0
			19	19		
5	B	10	Total	O	0	0
			10	10		
5	C	12	Total	O	0	0
			12	12		
5	D	15	Total	O	0	0
			15	15		
5	E	1	Total	O	0	0
			1	1		
5	P	2	Total	O	0	0
			2	2		
5	Q	5	Total	O	0	0
			5	5		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTEIN (ANTIBODY (LIGHT CHAIN))




- Molecule 1: PROTEIN (ANTIBODY (LIGHT CHAIN))



- Molecule 2: PROTEIN (ANTIBODY (HEAVY CHAIN))



- Molecule 2: PROTEIN (ANTIBODY (HEAVY CHAIN))



- Molecule 3: PROTEIN (C-MYC TAG AND HIS TAG)

Chain E: 

SER	GLU	GLN	LYS	LEU	ILE	S123	E124	GLU	ASP	LEU	ASN	HIS	HIS	HIS	HIS

- Molecule 3: PROTEIN (C-MYC TAG AND HIS TAG)

Chain F: 

S123	K126	LEU	ILE	SER	GLU	GLU	ASP	LEU	ASN	HIS	HIS	HIS	HIS

- Molecule 4: PROTEIN (P-GLYCOPROTEIN)

Chain P: 

V1	V2	L6	R10	E11	G12	ARG	THR

- Molecule 4: PROTEIN (P-GLYCOPROTEIN)

Chain Q: 

V1	V2	Q3	E4	A5	L6	D7	R10	R13	T14

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.79Å 82.66Å 93.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.40 38.80 – 2.40	Depositor EDS
% Data completeness (in resolution range)	95.6 (50.00-2.40) 95.6 (38.80-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.39Å)	Xtriage
Refinement program	CNS 0.5	Depositor
R, R_{free}	0.221 , 0.274 0.230 , 0.274	Depositor DCC
R_{free} test set	2047 reflections (11.02%)	DCC
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.473	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 28.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 21301 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3954	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/909	0.59	0/1237
1	C	0.33	0/909	0.56	0/1237
2	B	0.31	0/971	0.60	0/1323
2	D	0.32	0/971	0.62	0/1323
3	E	1.65	0/6	0.72	0/7
3	F	0.80	0/24	0.43	0/31
4	P	0.62	0/73	0.52	0/100
4	Q	0.38	0/109	0.59	0/143
All	All	0.35	0/3972	0.59	0/5401

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	888	0	862	53	0
1	C	888	0	862	48	0
2	B	949	0	912	52	0
2	D	949	0	912	44	0
3	E	7	0	4	1	0
3	F	25	0	18	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	P	74	0	65	3	0
4	Q	110	0	114	12	0
5	A	19	0	0	3	0
5	B	10	0	0	2	0
5	C	12	0	0	2	0
5	D	15	0	0	1	0
5	E	1	0	0	0	0
5	P	2	0	0	0	0
5	Q	5	0	0	2	0
All	All	3954	0	3749	199	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 26.

The worst 5 of 199 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:35:HIS:HD2	2:D:47:TRP:HE1	1.11	0.96
2:D:81:LEU:HD13	2:D:83:LEU:HD11	1.49	0.94
2:B:35:HIS:HD2	2:B:47:TRP:HE1	1.17	0.93
1:C:86:ALA:O	1:C:89:LEU:HD23	1.80	0.81
2:B:6:GLN:HE21	2:B:115:THR:HG23	1.45	0.81

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	113/115 (98%)	99 (88%)	8 (7%)	6 (5%)	3	1
1	C	113/115 (98%)	100 (88%)	9 (8%)	4 (4%)	6	4
2	B	120/122 (98%)	110 (92%)	7 (6%)	3 (2%)	9	8
2	D	120/122 (98%)	111 (92%)	7 (6%)	2 (2%)	14	17
3	F	2/17 (12%)	2 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	P	10/14 (71%)	9 (90%)	0	1 (10%)	1	0
4	Q	12/14 (86%)	11 (92%)	0	1 (8%)	1	0
All	All	490/519 (94%)	442 (90%)	31 (6%)	17 (4%)	6	4

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	0	ARG
1	A	89	LEU
1	C	11	LEU
1	C	90	ALA
2	D	43	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	101/101 (100%)	89 (88%)	12 (12%)	8	10
1	C	101/101 (100%)	95 (94%)	6 (6%)	28	42
2	B	103/103 (100%)	96 (93%)	7 (7%)	22	34
2	D	103/103 (100%)	95 (92%)	8 (8%)	18	27
3	E	1/17 (6%)	1 (100%)	0	100	100
3	F	3/17 (18%)	3 (100%)	0	100	100
4	P	6/11 (54%)	6 (100%)	0	100	100
4	Q	11/11 (100%)	10 (91%)	1 (9%)	14	20
All	All	429/464 (92%)	395 (92%)	34 (8%)	18	26

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	71	ILE
1	C	29	LEU
2	D	86	LEU
2	B	86	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	60	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	31	ASN
1	C	34	ASN
2	D	43	GLN
2	B	43	GLN
2	B	65	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	115/115 (100%)	-0.28	1 (0%) 81 81	14, 31, 51, 57	0
1	C	115/115 (100%)	-0.01	2 (1%) 67 65	20, 39, 61, 75	0
2	B	122/122 (100%)	-0.15	4 (3%) 44 42	20, 32, 60, 73	0
2	D	122/122 (100%)	-0.33	1 (0%) 83 82	19, 31, 52, 75	0
3	E	2/17 (11%)	0.47	0 100 100	36, 36, 36, 38	0
3	F	4/17 (23%)	1.08	1 (25%) 1 1	42, 52, 62, 70	0
4	P	12/14 (85%)	-0.02	0 100 100	28, 43, 66, 68	0
4	Q	14/14 (100%)	-0.34	0 100 100	15, 27, 34, 47	0
All	All	506/536 (94%)	-0.18	9 (1%) 65 63	14, 33, 60, 75	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	43	GLN	4.1
2	B	43	GLN	4.1
1	A	12	THR	3.1
3	F	126	LYS	2.7
1	C	10	SER	2.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.