



wwPDB X-ray Structure Validation Summary Report

Mar 1, 2014 – 01:24 AM GMT

PDB ID : 4AWL
Title : The NF-Y transcription factor is structurally and functionally a sequence specific histone
Authors : Nardini, M.; Gnesutta, N.; Donati, G.; Gatta, R.; Forni, C.; Fossati, A.; Vonrhein, C.; Moras, D.; Romier, C.; Mantovani, R.; Bolognesi, M.
Deposited on : 2012-06-04
Resolution : 3.08 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2204 (3.16-3.00)
Clashscore	79885	1061 (3.14-3.02)
Ramachandran outliers	78287	1025 (3.14-3.02)
Sidechain outliers	78261	1025 (3.14-3.02)
RSRZ outliers	66119	2206 (3.16-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	78	
2	B	94	
3	C	94	
4	I	25	
5	J	25	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2949 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NUCLEAR TRANSCRIPTION FACTOR Y SUBUNIT ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	62	Total	C	N	O	S	0	0	0
			530	330	114	84	2			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	MET	-	EXPRESSION TAG	UNP P23511
A	304	GLY	-	EXPRESSION TAG	UNP P23511
A	305	SER	-	EXPRESSION TAG	UNP P23511
A	306	LEU	-	EXPRESSION TAG	UNP P23511
A	307	VAL	-	EXPRESSION TAG	UNP P23511
A	308	PRO	-	EXPRESSION TAG	UNP P23511
A	309	ARG	-	EXPRESSION TAG	UNP P23511

- Molecule 2 is a protein called NUCLEAR TRANSCRIPTION FACTOR Y SUBUNIT BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	92	Total	C	N	O	S	0	0	0
			745	473	125	142	5			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	48	MET	-	EXPRESSION TAG	UNP P25208

- Molecule 3 is a protein called NUCLEAR TRANSCRIPTION FACTOR Y SUBUNIT GAMMA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	80	Total	C	N	O	S	0	0	0
			653	423	111	116	3			

- Molecule 4 is a DNA chain called HSP70 PROMOTER FRAGMENT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	I	25	Total	C	N	O	P	0	0	0
			505	242	91	148	24			

- Molecule 5 is a DNA chain called HSP70 PROMOTER FRAGMENT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	J	25	Total	C	N	O	P	0	0	0
			514	245	97	148	24			

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	O	0	0
			1	1		
6	I	1	Total	O	0	0
			1	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NUCLEAR TRANSCRIPTION FACTOR Y SUBUNIT ALPHA

Chain A: 




- Molecule 2: NUCLEAR TRANSCRIPTION FACTOR Y SUBUNIT BETA

Chain B: 



- Molecule 3: NUCLEAR TRANSCRIPTION FACTOR Y SUBUNIT GAMMA

Chain C: 




- Molecule 4: HSP70 PROMOTER FRAGMENT

Chain I: 



- Molecule 5: HSP70 PROMOTER FRAGMENT

Chain J: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.70Å 62.56Å 139.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.46 – 3.08 35.46 – 3.08	Depositor EDS
% Data completeness (in resolution range)	100.0 (35.46-3.08) 94.9 (35.46-3.08)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.88 (at 3.06Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.191 , 0.250 0.199 , 0.250	Depositor DCC
R_{free} test set	418 reflections (4.95%)	DCC
Wilson B-factor (Å ²)	76.6	Xtriage
Anisotropy	0.218	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 31.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 8866 reflections (0.011%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2949	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/541	1.02	3/714 (0.4%)
2	B	0.71	0/757	0.72	1/1014 (0.1%)
3	C	0.64	0/663	0.83	2/894 (0.2%)
4	I	1.04	0/565	1.84	20/869 (2.3%)
5	J	1.10	1/577 (0.2%)	2.01	25/890 (2.8%)
All	All	0.83	1/3103 (0.0%)	1.39	51/4381 (1.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	J	-4	DA	C3'-O3'	-5.44	1.36	1.44

The worst 5 of 51 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	290	PHE	CB-CA-C	14.63	139.67	110.40
5	J	-12	DA	O4'-C1'-N9	-13.21	98.75	108.00
5	J	-21	DA	O4'-C1'-N9	-10.76	100.47	108.00
1	A	289	PHE	CB-CA-C	9.56	129.53	110.40
4	I	23	DG	P-O3'-C3'	9.20	130.74	119.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	530	0	0	11	0
2	B	745	0	0	4	0
3	C	653	0	0	2	0
4	I	505	0	0	1	0
5	J	514	0	170	5	0
6	B	1	0	0	0	0
6	I	1	0	0	0	0
All	All	2949	0	170	21	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

The worst 5 of 21 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:259:GLU:CG	1:A:260:GLY:N	2.37	0.85
3:C:42:GLU:OE1	3:C:42:GLU:CA	2.30	0.77
1:A:243:TYR:C	1:A:243:TYR:CD1	2.67	0.67
1:A:289:PHE:O	1:A:290:PHE:CB	2.43	0.65
5:J:-23:DC:OP2	5:J:-23:DC:H2'	2.01	0.60

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	60/78 (77%)	47 (78%)	11 (18%)	2 (3%)	6	32
2	B	90/94 (96%)	83 (92%)	6 (7%)	1 (1%)	21	64
3	C	78/94 (83%)	74 (95%)	4 (5%)	0	100	100
All	All	228/266 (86%)	204 (90%)	21 (9%)	3 (1%)	18	58

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	260	GLY
1	A	235	PRO
2	B	139	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	52/67 (78%)	35 (67%)	17 (33%)	0	0
2	B	82/84 (98%)	70 (85%)	12 (15%)	5	18
3	C	71/85 (84%)	62 (87%)	9 (13%)	6	24
All	All	205/236 (87%)	167 (82%)	38 (18%)	2	9

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	51	ARG
2	B	67	LYS
3	C	96	LEU
2	B	55	ILE
2	B	79	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	62/78 (79%)	-0.08	0 100 100	55, 85, 123, 132	0
2	B	92/94 (97%)	-0.20	0 100 100	39, 78, 132, 171	0
3	C	80/94 (85%)	-0.25	0 100 100	40, 74, 105, 123	0
4	I	25/25 (100%)	-0.60	0 100 100	54, 72, 108, 116	0
5	J	25/25 (100%)	-0.53	0 100 100	48, 68, 90, 91	0
All	All	284/316 (89%)	-0.25	0 100 100	39, 77, 123, 171	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.