



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 04:26 PM GMT

PDB ID : 4AX6  
Title : HYPOCREA JECORINA CEL6A D221A MUTANT SOAKED WITH 6-CHLORO-4-PHENYLUMBELLIFERYL-BETA-CELLOBIOSIDE  
Authors : Wu, M.; Nerinckx, W.; Piens, K.; Ishida, T.; Hansson, H.; Stahlberg, J.; Sandgren, M.  
Deposited on : 2012-06-10  
Resolution : 2.30 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	363	
1	B	363	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	NAG	A	501	-	X
2	NAG	B	501	-	X
3	MAN	A	503	-	X
3	MAN	A	505	-	X
3	MAN	A	507	-	X
3	MAN	A	508	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Geometry	Electron density
3	MAN	A	509	-	X
3	MAN	B	507	-	X
3	MAN	B	508	-	X

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 6308 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called EXOGLUCANASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	363	Total	C	N	O	S	0	2	0
			2750	1746	463	531	10			
1	B	363	Total	C	N	O	S	0	0	0
			2744	1742	463	529	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	221	ALA	ASP	ENGINEERED MUTATION	UNP P07987
B	221	ALA	ASP	ENGINEERED MUTATION	UNP P07987

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is SUGAR (ALPHA-D-MANNOSE) (three-letter code: MAN) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is 6-CHLORANYL-7-OXIDANYL-4-PHENYL-CHROMEN-2-ONE (three-letter code: UWU) (formula: C<sub>15</sub>H<sub>9</sub>ClO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	Cl	O	0	0
			19	15	1	3		
4	B	1	Total	C	Cl	O	0	0
			19	15	1	3		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	2	Total	C	O	0	0
			22	12	10		
5	B	2	Total	C	O	0	0
			22	12	10		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	221	ALA	ASP	ENGINEERED MUTATION	UNP P07987
B	221	ALA	ASP	ENGINEERED MUTATION	UNP P07987

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	298	Total 298	O 298	0	0
6	B	246	Total 246	O 246	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: EXOGLUCANASE 2

Chain A: 



- Molecule 1: EXOGLUCANASE 2

Chain B: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.54Å 52.96Å 66.54Å 77.33° 78.60° 75.66°	Depositor
Resolution (Å)	64.17 – 2.30 29.99 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.4 (64.17-2.30) 98.4 (29.99-2.30)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.43 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.204 , 0.254 0.204 , 0.254	Depositor DCC
$R_{free}$ test set	1379 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	11.2	Xtriage
Anisotropy	0.661	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 10.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 27611 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	6308	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, NAG, UWU, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.50	3/2832 (0.1%)	0.49	0/3882
1	B	0.50	7/2820 (0.2%)	0.49	0/3866
All	All	0.50	10/5652 (0.2%)	0.49	0/7748

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	342	TRP	CD2-CE2	5.12	1.47	1.41
1	B	430	TRP	CD2-CE2	5.11	1.47	1.41
1	A	272	TRP	CD2-CE2	5.10	1.47	1.41
1	B	367	TRP	CD2-CE2	5.08	1.47	1.41
1	B	269	TRP	CD2-CE2	5.06	1.47	1.41
1	B	309	TRP	CD2-CE2	5.03	1.47	1.41
1	A	309	TRP	CD2-CE2	5.03	1.47	1.41
1	B	99	TRP	CD2-CE2	5.02	1.47	1.41
1	A	99	TRP	CD2-CE2	5.01	1.47	1.41
1	B	135	TRP	CD2-CE2	5.00	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the

chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2750	0	2617	14	0
1	B	2744	0	2609	10	0
2	A	28	0	26	0	0
2	B	28	0	26	0	0
3	A	77	0	70	0	0
3	B	55	0	50	0	0
4	A	19	0	8	1	0
4	B	19	0	8	3	0
5	A	22	0	19	0	0
5	B	22	0	19	1	0
6	A	298	0	0	0	0
6	B	246	0	0	2	0
All	All	6308	0	5452	28	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (28) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:B:600:UWU:H8	6:B:2074:HOH:O	1.93	0.66
1:B:410:ARG:HH21	1:B:426:GLN:HG3	1.62	0.61
4:B:600:UWU:C16	4:B:600:UWU:H5	2.30	0.61
1:B:264:ALA:HB2	1:B:299:LEU:HG	1.88	0.56
1:A:112:ALA:HA	1:A:432:GLN:HE22	1.73	0.53
4:B:600:UWU:H16	4:B:600:UWU:H5	1.94	0.49
1:A:277:ASP:OD1	1:A:340:HIS:HE1	1.96	0.48
1:A:87:THR:HG22	1:A:256:PRO:HB2	1.97	0.47
1:B:304:ALA:HB1	5:B:601:BGC:H2	1.96	0.47
1:A:109:SER:HA	1:A:113:ILE:HD12	1.97	0.47
1:A:262:LEU:HG	1:A:296:LEU:HD11	1.98	0.46
1:B:305:ASN:ND2	1:B:306:TYR:H	2.13	0.46
1:B:220:PRO:HA	1:B:263:ASP:CG	2.36	0.46
4:A:600:UWU:H5	4:A:600:UWU:C16	2.44	0.46
1:B:109:SER:HA	1:B:113:ILE:HD12	1.96	0.46
1:A:86:ALA:O	1:A:213:ARG:HD3	2.17	0.44
1:A:361:GLN:HG3	1:A:370:VAL:HG11	2.00	0.44
1:B:180:ALA:HB3	6:B:2076:HOH:O	2.17	0.44
1:A:410:ARG:HB2	1:A:426:GLN:HE22	1.82	0.44
1:B:302:ASN:OD1	1:B:305:ASN:HB3	2.18	0.44
1:A:374:GLY:HA2	1:A:438:LEU:HA	2.01	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:374:GLY:HA2	1:B:438:LEU:HA	2.02	0.42
1:B:137:ASP:HA	1:B:169:TYR:O	2.20	0.41
1:A:305:ASN:ND2	1:A:306:TYR:H	2.17	0.41
1:A:277:ASP:HB3	1:A:278:PRO:HD3	2.03	0.41
1:A:255:LEU:HB3	1:A:256:PRO:HD2	2.03	0.41
1:A:287:TYR:HD1	1:A:296:LEU:HD23	1.85	0.41
1:A:133:PHE:HB2	1:A:394:VAL:HA	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/363 (100%)	351 (97%)	12 (3%)	0	100	100
1	B	361/363 (99%)	347 (96%)	14 (4%)	0	100	100
All	All	724/726 (100%)	698 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/283 (101%)	284 (100%)	1 (0%)	95	98
1	B	283/283 (100%)	281 (99%)	2 (1%)	91	97
All	All	568/566 (100%)	565 (100%)	3 (0%)	94	98

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	431	PHE
1	B	299	LEU
1	B	431	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	182	ASN
1	A	237	ASN
1	A	285	ASN
1	A	305	ASN
1	A	362	GLN
1	A	426	GLN
1	A	432	GLN
1	A	441	ASN
1	B	182	ASN
1	B	237	ASN
1	B	285	ASN
1	B	305	ASN
1	B	319	GLN
1	B	351	GLN
1	B	382	ASN
1	B	441	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	BGC	A	601	5,4	10,11,12	0.75	0	11,15,17	2.15	3 (27%)
5	BGC	A	602	5	10,11,12	0.49	0	11,15,17	1.10	1 (9%)
5	BGC	B	601	5,4	10,11,12	0.89	0	11,15,17	2.67	3 (27%)
5	BGC	B	602	5	10,11,12	0.51	0	11,15,17	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BGC	A	601	5,4	-	0/2/19/22	0/1/1/1
5	BGC	A	602	5	-	0/2/19/22	0/1/1/1
5	BGC	B	601	5,4	-	0/2/19/22	0/1/1/1
5	BGC	B	602	5	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	601	BGC	O5-C5-C6	6.68	113.99	106.98
5	A	601	BGC	O5-C5-C6	5.25	112.50	106.98
5	B	601	BGC	O5-C5-C4	4.13	115.89	110.65
5	A	601	BGC	O5-C5-C4	3.26	114.78	110.65
5	A	602	BGC	O5-C5-C6	2.91	110.03	106.98
5	B	601	BGC	C6-C5-C4	-2.43	107.12	113.00
5	A	601	BGC	O4-C4-C3	-2.09	105.66	110.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.6 Ligand geometry ⓘ

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	501	1	12,14,15	0.57	0	15,19,21	0.82	1 (6%)
2	NAG	A	502	1	12,14,15	0.60	0	15,19,21	0.79	0
3	MAN	A	503	1	10,11,12	0.64	0	11,15,17	1.02	0
3	MAN	A	504	1	10,11,12	0.68	0	11,15,17	1.02	1 (9%)
3	MAN	A	505	1	10,11,12	0.68	0	11,15,17	0.65	0
3	MAN	A	506	1	10,11,12	0.73	0	11,15,17	1.04	1 (9%)
3	MAN	A	507	1	10,11,12	0.70	0	11,15,17	0.80	0
3	MAN	A	508	1	10,11,12	0.73	0	11,15,17	0.77	0
3	MAN	A	509	1	10,11,12	0.74	0	11,15,17	0.91	1 (9%)
4	UWU	A	600	5	21,21,21	2.37	4 (19%)	28,30,30	1.20	3 (10%)
2	NAG	B	501	1	12,14,15	0.60	0	15,19,21	1.15	2 (13%)
2	NAG	B	502	1	12,14,15	0.63	0	15,19,21	0.78	1 (6%)
3	MAN	B	504	1	10,11,12	0.71	0	11,15,17	0.94	1 (9%)
3	MAN	B	506	1	10,11,12	0.73	0	11,15,17	0.74	0
3	MAN	B	507	1	10,11,12	0.71	0	11,15,17	0.79	1 (9%)
3	MAN	B	508	1	10,11,12	0.72	0	11,15,17	0.98	0
3	MAN	B	509	1	10,11,12	0.74	0	11,15,17	1.02	1 (9%)
4	UWU	B	600	5	21,21,21	2.33	4 (19%)	28,30,30	1.21	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	501	1	-	0/6/23/26	0/1/1/1
2	NAG	A	502	1	-	0/6/23/26	0/1/1/1
3	MAN	A	503	1	-	0/2/19/22	0/1/1/1
3	MAN	A	504	1	-	0/2/19/22	0/1/1/1
3	MAN	A	505	1	-	0/2/19/22	0/1/1/1
3	MAN	A	506	1	-	0/2/19/22	0/1/1/1
3	MAN	A	507	1	-	0/2/19/22	0/1/1/1
3	MAN	A	508	1	-	0/2/19/22	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	A	509	1	-	0/2/19/22	0/1/1/1
4	UWU	A	600	5	-	0/4/4/4	0/1/3/3
2	NAG	B	501	1	-	0/6/23/26	0/1/1/1
2	NAG	B	502	1	-	0/6/23/26	0/1/1/1
3	MAN	B	504	1	-	0/2/19/22	0/1/1/1
3	MAN	B	506	1	-	0/2/19/22	0/1/1/1
3	MAN	B	507	1	-	0/2/19/22	0/1/1/1
3	MAN	B	508	1	-	0/2/19/22	0/1/1/1
3	MAN	B	509	1	-	0/2/19/22	0/1/1/1
4	UWU	B	600	5	-	0/4/4/4	0/1/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	600	UWU	O8-C2	8.66	1.37	1.33
4	B	600	UWU	O8-C2	8.48	1.37	1.33
4	A	600	UWU	C4-C11	-4.68	1.40	1.49
4	B	600	UWU	C4-C11	-4.63	1.41	1.49
4	A	600	UWU	C8-C7	2.91	1.40	1.36
4	B	600	UWU	C8-C7	2.80	1.40	1.36
4	A	600	UWU	C5-C6	2.50	1.39	1.36
4	B	600	UWU	C5-C6	2.28	1.39	1.36

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	600	UWU	C2-O8-C8A	-3.59	117.43	122.33
4	B	600	UWU	C2-O8-C8A	-3.50	117.56	122.33
3	A	506	MAN	O5-C5-C6	2.73	109.85	106.98
3	B	504	MAN	O5-C5-C6	2.72	109.83	106.98
2	B	501	NAG	C3-C2-N2	-2.60	107.80	111.76
4	B	600	UWU	C8-C8A-C4A	-2.53	120.00	123.10
4	A	600	UWU	O8-C2-C3	2.49	122.33	119.41
2	B	501	NAG	O5-C5-C6	2.36	109.46	106.98
2	B	502	NAG	O5-C5-C6	2.34	109.44	106.98
4	A	600	UWU	C8-C8A-C4A	-2.34	120.24	123.10
4	B	600	UWU	O8-C2-C3	2.20	121.99	119.41
3	A	504	MAN	O5-C5-C6	2.19	109.28	106.98
3	B	509	MAN	C4-C3-C2	2.15	113.39	110.50
2	A	501	NAG	O5-C5-C6	2.10	109.19	106.98
3	B	507	MAN	O5-C5-C6	2.09	109.17	106.98
3	A	509	MAN	O5-C5-C6	2.07	109.15	106.98



There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	363/363 (100%)	0.09	3 (0%)	83 90	8, 11, 17, 21	1 (0%)
1	B	363/363 (100%)	0.27	3 (0%)	83 90	10, 14, 22, 28	3 (0%)
All	All	726/726 (100%)	0.18	6 (0%)	83 90	8, 12, 19, 28	4 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	319	GLN	3.3
1	A	158	ASN	3.2
1	A	159	GLY	2.9
1	B	119	ALA	2.4
1	A	161	ASN	2.1
1	B	85	THR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	BGC	B	601	11/12	0.23	1.74	15,15,15,15	0
5	BGC	B	602	11/12	0.16	0.46	14,15,15,15	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	BGC	A	601	11/12	0.15	0.12	11,11,11,11	0
5	BGC	A	602	11/12	0.13	-0.45	10,10,11,11	0

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MAN	A	508	11/12	0.24	6.81	22,23,23,23	0
3	MAN	A	503	11/12	0.31	5.42	19,20,21,21	0
2	NAG	B	501	14/15	0.22	4.40	18,19,19,19	0
3	MAN	B	507	11/12	0.31	3.66	29,30,30,31	0
2	NAG	A	501	14/15	0.24	3.20	16,16,16,17	0
3	MAN	A	505	11/12	0.19	3.17	16,16,16,16	0
3	MAN	A	509	11/12	0.26	2.82	19,20,20,20	0
3	MAN	B	508	11/12	0.19	2.12	29,30,30,30	0
3	MAN	A	507	11/12	0.24	2.05	18,18,18,18	0
3	MAN	B	504	11/12	0.23	1.93	19,19,19,19	0
4	UWU	B	600	19/19	0.21	1.86	15,15,16,16	0
2	NAG	B	502	14/15	0.17	1.24	14,14,15,15	0
3	MAN	A	506	11/12	0.18	1.04	16,16,16,16	0
4	UWU	A	600	19/19	0.16	0.82	12,12,12,13	0
3	MAN	A	504	11/12	0.22	0.79	16,16,16,16	0
3	MAN	B	509	11/12	0.23	0.61	28,28,28,28	0
3	MAN	B	506	11/12	0.17	0.17	25,25,26,26	0
2	NAG	A	502	14/15	0.12	-0.83	10,10,10,10	0

## 6.5 Other polymers

There are no such residues in this entry.