



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 08:51 AM GMT

PDB ID : 3B29  
Title : Human leukotriene C4 synthase in complex with dodecyl-beta-D-selenomalto  
side  
Authors : Saino, H.; Ago, H.; Miyano, M.  
Deposited on : 2011-07-22  
Resolution : 3.20 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance


The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1824 (3.30-3.10)
Clashscore	79885	1078 (3.26-3.14)
Ramachandran outliers	78287	1059 (3.26-3.14)
Sidechain outliers	78261	1058 (3.26-3.14)
RSRZ outliers	66119	1825 (3.30-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	156	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	LSM	A	203	-	X
4	LSM	A	207	-	X
4	LSM	A	210	-	X

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 1187 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Leukotriene C4 synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	142	1072	724	174	172	2	0	1	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	151	HIS	-	EXPRESSION TAG	UNP Q16873
A	152	HIS	-	EXPRESSION TAG	UNP Q16873
A	153	HIS	-	EXPRESSION TAG	UNP Q16873
A	154	HIS	-	EXPRESSION TAG	UNP Q16873
A	155	HIS	-	EXPRESSION TAG	UNP Q16873
A	156	HIS	-	EXPRESSION TAG	UNP Q16873

- Molecule 2 is GLUTATHIONE (three-letter code: GSH) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>3</sub>O<sub>6</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			20	10	3	6	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is SUGAR (DODECYL 4-O-ALPHA-D-GLUCOPYRANOSYL-1-SELENO-BETA-D-GLUCOPYRANOSIDE) (three-letter code: LSM) (formula: C<sub>24</sub>H<sub>46</sub>O<sub>10</sub>Se).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C 7 7	0	0
4	A	1	Total C Se 12 11 1	0	0
4	A	1	Total C 12 12	0	0
4	A	1	Total C O Se 32 21 10 1	0	0
4	A	1	Total C O Se 24 18 5 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total O 3 3	0	0

i

- Molecule 1: Leukotriene C4 synthase

HIS	MET
	K2
	V5
	L8
	L14
	E58
	Y59
	F60
	P61
	L64
	W68
	G71
	H75
	A78
	G83
	L87
	L91
	H92
	Y93
	Y97
	Q102
	L105
	A106
	P107
	A112
	L131
	P132
	R136
	L143
	ARG
	THR
	LEU
	LEU
	PRO
	TRP
	ALA
	HIS
	HIS
	HIS
	HIS

## 4 Data and refinement statistics

Property	Value	Source
Space group	F 2 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	168.59Å 168.59Å 168.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.42 – 3.20 25.42 – 3.20	Depositor EDS
% Data completeness (in resolution range)	90.2 (25.42-3.20) 90.1 (25.42-3.20)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.83 (at 3.24Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.203 , 0.209 0.204 , 0.214	Depositor DCC
$R_{free}$ test set	277 reflections (4.83%)	DCC
Wilson B-factor (Å <sup>2</sup> )	47.0	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 16.2	EDS
Estimated twinning fraction	0.060 for k,h,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 6015 reflections	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	1187	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GSH, LSM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.23	0/1104	0.33	0/1509

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1072	0	1085	18	0
2	A	20	0	15	1	0
3	A	5	0	0	0	0
4	A	87	0	118	2	0
5	A	3	0	0	0	0
All	All	1187	0	1218	19	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (19) close contacts within the same asymmetric unit are listed below.



Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:75:HIS:HD2	1:A:78:ALA:H	1.14	0.93
1:A:106:ALA:HB3	1:A:107:PRO:HD3	1.50	0.90
1:A:131:LEU:HB3	1:A:132:PRO:HD3	1.72	0.71
1:A:8:LEU:HD11	1:A:71:GLY:HA3	1.76	0.66
1:A:60:PHE:HB3	1:A:61:PRO:HD3	1.81	0.63
4:A:209:LSM:O3'	4:A:209:LSM:H1B	2.01	0.61
1:A:75:HIS:HD2	1:A:78:ALA:N	1.93	0.60
1:A:75:HIS:CD2	1:A:78:ALA:H	2.06	0.58
1:A:106:ALA:HB3	1:A:107:PRO:CD	2.34	0.49
1:A:91:LEU:O	1:A:91:LEU:HD23	2.12	0.48
1:A:58:GLU:OE1	2:A:201:GSH:N1	2.41	0.48
1:A:83:GLY:O	1:A:87:LEU:HG	2.16	0.45
1:A:106:ALA:CB	1:A:107:PRO:HD3	2.32	0.45
1:A:5:VAL:HB	1:A:68:TRP:CZ3	2.53	0.44
1:A:64:LEU:O	1:A:68:TRP:HD1	2.01	0.43
1:A:112:ALA:HB2	4:A:210:LSM:H5A	2.01	0.42
1:A:102[B]:GLN:NE2	1:A:105:LEU:HD12	2.35	0.42
1:A:93:TYR:HA	1:A:107:PRO:HB2	2.02	0.41
1:A:91:LEU:C	1:A:91:LEU:HD23	2.40	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/156 (90%)	138 (98%)	3 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/120 (79%)	90 (95%)	5 (5%)	32	75

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LEU
1	A	97	TYR
1	A	102[A]	GLN
1	A	102[B]	GLN
1	A	136	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GSH	A	201	-	19,19,19	0.79	0	24,24,24	0.63	0
3	SO4	A	202	-	4,4,4	0.14	0	6,6,6	0.08	0
4	LSM	A	203	-	6,6,36	1.53	1 (16%)	5,5,47	0.81	0
4	LSM	A	207	-	10,11,36	1.28	1 (10%)	9,10,47	0.98	0
4	LSM	A	208	-	11,11,36	1.27	1 (9%)	10,10,47	0.74	0
4	LSM	A	209	-	33,33,36	0.88	2 (6%)	44,44,47	0.98	2 (4%)
4	LSM	A	210	-	24,24,36	0.70	1 (4%)	29,29,47	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GSH	A	201	-	-	0/24/24/24	0/0/0/0
3	SO4	A	202	-	-	0/0/0/0	0/0/0/0
4	LSM	A	203	-	-	0/4/4/61	0/0/0/2
4	LSM	A	207	-	-	0/9/9/61	0/0/0/2
4	LSM	A	208	-	-	0/9/9/61	0/0/0/2
4	LSM	A	209	-	-	0/16/58/61	0/2/2/2
4	LSM	A	210	-	-	0/13/35/61	0/1/1/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	208	LSM	C1-C2	-4.03	1.53	1.55
4	A	207	LSM	C10-C9	-3.80	1.53	1.55
4	A	209	LSM	C9-C8	-3.71	1.53	1.55
4	A	203	LSM	C6-C7	-3.67	1.53	1.55
4	A	210	LSM	O5'-C1'	2.21	1.44	1.41
4	A	209	LSM	O5'-C1'	2.09	1.44	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	209	LSM	C1B-O1B-C4'	-3.06	110.18	117.99
4	A	209	LSM	SE-C1-C2	-2.38	108.71	113.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	142/156 (91%)	-0.43	0 100 100	20, 29, 42, 51	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	LSM	A	203	7/35	0.23	3.45	37,39,41,41	0
4	LSM	A	207	12/35	0.25	2.88	63,71,91,94	0
4	LSM	A	210	24/35	0.34	2.10	49,59,65,68	1
4	LSM	A	208	12/35	0.17	1.45	45,50,60,61	0
4	LSM	A	209	32/35	0.24	1.00	46,71,80,81	0
2	GSH	A	201	20/20	0.12	-0.58	28,30,35,36	0
3	SO4	A	202	5/5	0.12	-0.83	32,33,33,33	5

## 6.5 Other polymers ⓘ

There are no such residues in this entry.