



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 02:24 PM GMT

PDB ID : 1B3B
Title : THERMOTOGA MARITIMA GLUTAMATE DEHYDROGENASE MUTANT N97D, G376K
Authors : Knapp, S.; Lebbink, J.H.G.; Van Der Oost, J.; Devos, W.M.; Rice, D.; Ladenstein, R.
Deposited on : 1998-12-07
Resolution : 3.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1078 (3.16-3.04)
Ramachandran outliers	78287	1044 (3.16-3.04)
Sidechain outliers	78261	1044 (3.16-3.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	415	
1	B	415	
1	C	415	
1	D	415	
1	E	415	
1	F	415	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 19008 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (GLUTAMATE DEHYDROGENASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			
1	B	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			
1	C	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			
1	D	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			
1	E	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			
1	F	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	97	ASP	ASN	ENGINEERED	UNP P96110
A	376	LYS	GLY	ENGINEERED	UNP P96110
B	97	ASP	ASN	ENGINEERED	UNP P96110
B	376	LYS	GLY	ENGINEERED	UNP P96110
C	97	ASP	ASN	ENGINEERED	UNP P96110
C	376	LYS	GLY	ENGINEERED	UNP P96110
D	97	ASP	ASN	ENGINEERED	UNP P96110
D	376	LYS	GLY	ENGINEERED	UNP P96110
E	97	ASP	ASN	ENGINEERED	UNP P96110
E	376	LYS	GLY	ENGINEERED	UNP P96110
F	97	ASP	ASN	ENGINEERED	UNP P96110
F	376	LYS	GLY	ENGINEERED	UNP P96110


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)

Chain A:




• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)

Chain B:




• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)

Chain C:




• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)


Chain D:



• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)


Chain E:





• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)

Chain F:



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	145.10Å 145.10Å 272.50Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 3.10	Depositor
% Data completeness (in resolution range)	94.0 (8.00-3.10)	Depositor
R_{merge}	0.10	Depositor
R_{sym}	0.05	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.225 , 0.298	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	19008	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/3227	0.53	0/4367
1	B	0.33	0/3227	0.52	0/4367
1	C	0.33	0/3227	0.51	0/4367
1	D	0.31	0/3227	0.51	0/4367
1	E	0.35	0/3227	0.53	0/4367
1	F	0.33	0/3227	0.51	1/4367 (0.0%)
All	All	0.33	0/19362	0.52	1/26202 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	168	LEU	CA-CB-CG	5.21	127.29	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3168	0	3192	118	18
1	B	3168	0	3192	117	20
1	C	3168	0	3192	114	2
1	D	3168	0	3192	124	2

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	3168	0	3192	139	18
1	F	3168	0	3192	123	20
All	All	19008	0	19152	722	40

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 19.

The worst 5 of 722 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:E:248:ASN:HD21	1:E:272:LYS:HE2	1.22	1.03
1:F:248:ASN:HD21	1:F:272:LYS:HE2	1.32	0.95
1:A:248:ASN:HD21	1:A:272:LYS:HE2	1.38	0.88
1:E:224:ALA:O	1:E:228:ILE:HG13	1.74	0.88
1:C:354:GLN:HG2	1:C:360:PHE:HA	1.55	0.87

The worst 5 of 40 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:299:HIS:CG	1:E:304:GLU:OE1[6_655]	0.79	1.41
1:B:279:GLU:CG	1:F:330:ARG:O[6_665]	1.00	1.20
1:A:279:GLU:OE1	1:E:330:ARG:NE[6_655]	1.11	1.09
1:B:296:GLY:O	1:F:329:ARG:NH2[6_665]	1.20	1.00
1:A:299:HIS:ND1	1:E:304:GLU:OE1[6_655]	1.28	0.92

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	407/415 (98%)	365 (90%)	39 (10%)	3 (1%)	30	76
1	B	407/415 (98%)	366 (90%)	37 (9%)	4 (1%)	22	68
1	C	407/415 (98%)	369 (91%)	35 (9%)	3 (1%)	30	76
1	D	407/415 (98%)	370 (91%)	33 (8%)	4 (1%)	22	68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	407/415 (98%)	373 (92%)	32 (8%)	2 (0%)	38	81
1	F	407/415 (98%)	366 (90%)	37 (9%)	4 (1%)	22	68
All	All	2442/2490 (98%)	2209 (90%)	213 (9%)	20 (1%)	27	74

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	178	LEU
1	A	178	LEU
1	A	303	ALA
1	B	178	LEU
1	B	303	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/339 (98%)	303 (91%)	30 (9%)	14	47
1	B	333/339 (98%)	303 (91%)	30 (9%)	14	47
1	C	333/339 (98%)	301 (90%)	32 (10%)	12	42
1	D	333/339 (98%)	303 (91%)	30 (9%)	14	47
1	E	333/339 (98%)	305 (92%)	28 (8%)	16	52
1	F	333/339 (98%)	304 (91%)	29 (9%)	15	49
All	All	1998/2034 (98%)	1819 (91%)	179 (9%)	14	47

5 of 179 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	283	GLU
1	D	166	THR
1	F	269	THR
1	C	348	SER
1	C	408	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such

sidechains are listed below:

Mol	Chain	Res	Type
1	C	365	GLN
1	D	61	ASN
1	E	215	GLN
1	C	357	GLN
1	E	357	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.