



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 01:17 PM GMT

PDB ID : 1B4A
Title : STRUCTURE OF THE ARGININE REPRESSOR FROM BACILLUS
STEAROTHERMOPHILUS
Authors : Ni, J.; Sakanyan, V.; Charlier, D.; Glansdorff, N.; Van Duyne, G.D.
Deposited on : 1998-12-18
Resolution : 2.50 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	149	
1	B	149	
1	C	149	
1	D	149	
1	E	149	
1	F	149	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7487 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ARGININE REPRESSOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	146	Total	C	N	O	S	0	0	0
			1150	721	206	216	7			
1	B	146	Total	C	N	O	S	0	0	0
			1150	721	206	216	7			
1	C	146	Total	C	N	O	S	0	0	0
			1150	721	206	216	7			
1	D	146	Total	C	N	O	S	0	0	0
			1150	721	206	216	7			
1	E	146	Total	C	N	O	S	0	0	0
			1150	721	206	216	7			
1	F	146	Total	C	N	O	S	0	0	0
			1150	721	206	216	7			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	87	Total	O	0	0
			87	87		
2	B	120	Total	O	0	0
			120	120		
2	C	74	Total	O	0	0
			74	74		
2	D	109	Total	O	0	0
			109	109		
2	E	114	Total	O	0	0
			114	114		
2	F	83	Total	O	0	0
			83	83		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: ARGININE REPRESSOR

Chain A: 




• Molecule 1: ARGININE REPRESSOR

Chain B: 




• Molecule 1: ARGININE REPRESSOR

Chain C: 



• Molecule 1: ARGININE REPRESSOR



Chain D: 






• Molecule 1: ARGININE REPRESSOR

Chain E: A horizontal bar chart representing the sequence of Chain E. The bar is color-coded: green for good, yellow for medium, and orange for bad. The bar is divided into segments representing different regions of the protein.



• Molecule 1: ARGININE REPRESSOR

Chain F: A horizontal bar chart representing the sequence of Chain F. The bar is color-coded: green for good, yellow for medium, and orange for bad. The bar is divided into segments representing different regions of the protein.



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	72.80Å 121.90Å 227.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.50	Depositor
% Data completeness (in resolution range)	93.0 (8.00-2.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.224 , 0.279	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7487	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/1163	0.64	0/1568
1	B	0.43	0/1163	0.66	1/1568 (0.1%)
1	C	0.46	0/1163	0.69	0/1568
1	D	0.45	0/1163	0.71	3/1568 (0.2%)
1	E	0.41	0/1163	0.62	0/1568
1	F	0.38	0/1163	0.60	0/1568
All	All	0.43	0/6978	0.65	4/9408 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	69	ARG	NE-CZ-NH2	6.89	123.74	120.30
1	D	69	ARG	NE-CZ-NH2	6.15	123.38	120.30
1	D	48	MET	CG-SD-CE	6.11	109.98	100.20
1	D	55	MET	CG-SD-CE	5.92	109.67	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1150	0	1194	66	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1150	0	1194	57	0
1	C	1150	0	1194	107	0
1	D	1150	0	1194	60	0
1	E	1150	0	1194	60	0
1	F	1150	0	1194	49	0
2	A	87	0	0	18	1
2	B	120	0	0	9	2
2	C	74	0	0	24	0
2	D	109	0	0	7	0
2	E	114	0	0	16	1
2	F	83	0	0	9	0
All	All	7487	0	7164	371	3

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 26.

The worst 5 of 371 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:11:ARG:CZ	2:C:220:HOH:O	1.65	1.33
1:C:68:GLN:HB3	2:C:197:HOH:O	1.32	1.24
1:B:133:ARG:HB3	2:B:266:HOH:O	1.48	1.11
1:C:9:LYS:HB3	2:C:216:HOH:O	1.52	1.09
1:B:90:THR:HG21	1:C:133:ARG:HA	1.38	1.04

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:A:192:HOH:O	2:A:210:HOH:O[8_555]	0.18	2.02
2:B:168:HOH:O	2:B:168:HOH:O[3_555]	0.33	1.87
2:B:249:HOH:O	2:E:209:HOH:O[3_655]	1.21	0.99

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/149 (97%)	138 (96%)	6 (4%)	0	100	100
1	B	144/149 (97%)	141 (98%)	3 (2%)	0	100	100
1	C	144/149 (97%)	135 (94%)	8 (6%)	1 (1%)	30	50
1	D	144/149 (97%)	141 (98%)	2 (1%)	1 (1%)	30	50
1	E	144/149 (97%)	139 (96%)	5 (4%)	0	100	100
1	F	144/149 (97%)	133 (92%)	7 (5%)	4 (3%)	8	10
All	All	864/894 (97%)	827 (96%)	31 (4%)	6 (1%)	30	50

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	68	GLN
1	F	68	GLN
1	F	67	ASP
1	F	70	PHE
1	F	66	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	130/133 (98%)	93 (72%)	37 (28%)	0	0
1	B	130/133 (98%)	105 (81%)	25 (19%)	2	3
1	C	130/133 (98%)	99 (76%)	31 (24%)	1	1
1	D	130/133 (98%)	106 (82%)	24 (18%)	2	4
1	E	130/133 (98%)	103 (79%)	27 (21%)	2	3
1	F	130/133 (98%)	99 (76%)	31 (24%)	1	1
All	All	780/798 (98%)	605 (78%)	175 (22%)	1	2

5 of 175 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	78	ARG
1	D	46	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	82	ASP
1	C	94	LEU
1	C	133	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	74	GLN
1	C	143	ASN
1	F	71	ASN
1	C	104	HIS
1	C	112	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.