



# wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 12:21 AM GMT

PDB ID : 2B7M  
Title : Crystal Structure of the *S. cerevisiae* Exocyst Component Exo70p  
Authors : Hamburger, Z.A.; Hamburger, A.E.; West, A.P.; Weis, W.I.  
Deposited on : 2005-10-04  
Resolution : 3.50 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1243 (3.70-3.30)
Clashscore	79885	1039 (3.66-3.34)
Ramachandran outliers	78287	1000 (3.66-3.34)
Sidechain outliers	78261	1000 (3.66-3.34)
RSRZ outliers	66119	1243 (3.70-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	566	
1	B	566	
1	C	566	
1	D	566	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 16823 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exocyst complex component EXO70.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	513	Total	C	N	O	S	Se	0	0	0
			4153	2637	703	795	5	13			
1	B	523	Total	C	N	O	S	Se	0	0	0
			4247	2699	721	809	5	13			
1	C	518	Total	C	N	O	S	Se	0	0	0
			4205	2676	712	799	5	13			
1	D	520	Total	C	N	O	S	Se	0	0	0
			4218	2683	715	802	5	13			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	58	GLY	-	CLONING ARTIFACT	UNP P19658
A	59	ALA	-	CLONING ARTIFACT	UNP P19658
A	60	MSE	-	CLONING ARTIFACT	UNP P19658
A	61	GLY	-	CLONING ARTIFACT	UNP P19658
A	62	SER	-	CLONING ARTIFACT	UNP P19658
A	110	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	169	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	214	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	239	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	376	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	393	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	451	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	478	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	495	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	515	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	527	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	586	MSE	MET	MODIFIED RESIDUE	UNP P19658
A	588	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	58	GLY	-	CLONING ARTIFACT	UNP P19658
B	59	ALA	-	CLONING ARTIFACT	UNP P19658
B	60	MSE	-	CLONING ARTIFACT	UNP P19658

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	61	GLY	-	CLONING ARTIFACT	UNP P19658
B	62	SER	-	CLONING ARTIFACT	UNP P19658
B	110	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	169	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	214	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	239	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	376	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	393	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	451	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	478	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	495	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	515	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	527	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	586	MSE	MET	MODIFIED RESIDUE	UNP P19658
B	588	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	58	GLY	-	CLONING ARTIFACT	UNP P19658
C	59	ALA	-	CLONING ARTIFACT	UNP P19658
C	60	MSE	-	CLONING ARTIFACT	UNP P19658
C	61	GLY	-	CLONING ARTIFACT	UNP P19658
C	62	SER	-	CLONING ARTIFACT	UNP P19658
C	110	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	169	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	214	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	239	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	376	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	393	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	451	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	478	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	495	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	515	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	527	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	586	MSE	MET	MODIFIED RESIDUE	UNP P19658
C	588	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	58	GLY	-	CLONING ARTIFACT	UNP P19658
D	59	ALA	-	CLONING ARTIFACT	UNP P19658
D	60	MSE	-	CLONING ARTIFACT	UNP P19658
D	61	GLY	-	CLONING ARTIFACT	UNP P19658
D	62	SER	-	CLONING ARTIFACT	UNP P19658
D	110	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	169	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	214	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	239	MSE	MET	MODIFIED RESIDUE	UNP P19658

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
D	376	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	393	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	451	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	478	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	495	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	515	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	527	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	586	MSE	MET	MODIFIED RESIDUE	UNP P19658
D	588	MSE	MET	MODIFIED RESIDUE	UNP P19658





● Molecule 1: Exocyst complex component EXO70

Chain D: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	415.73Å 55.13Å 132.93Å 90.00° 104.05° 90.00°	Depositor
Resolution (Å)	20.00 – 3.50 29.93 – 3.50	Depositor EDS
% Data completeness (in resolution range)	95.5 (20.00-3.50) 95.3 (29.93-3.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.98 (at 3.47Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.254 , 0.305 0.243 , 0.299	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	102.1	Xtriage
Anisotropy	0.363	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 79.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 36077 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	16823	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	129.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/4212	0.62	0/5654
1	B	0.33	0/4308	0.58	1/5779 (0.0%)
1	C	0.34	0/4265	0.57	0/5720
1	D	0.37	0/4278	0.59	0/5738
All	All	0.37	0/17063	0.59	1/22891 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	125	SER	N-CA-C	5.53	125.94	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4153	0	4177	295	0
1	B	4247	0	4278	287	0
1	C	4205	0	4241	272	0
1	D	4218	0	4252	295	0
All	All	16823	0	16948	1130	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 33.

The worst 5 of 1130 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:398:ARG:HB2	1:B:401:TRP:CZ2	1.58	1.37
1:A:120:ASN:HB3	1:D:320:ARG:HH22	1.01	1.10
1:C:526:LEU:HD12	1:C:552:PHE:HB2	1.28	1.10
1:A:476:ILE:HG21	1:A:511:TYR:CZ	1.86	1.10
1:D:398:ARG:HB2	1:D:401:TRP:CZ2	1.89	1.08

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	507/566 (90%)	403 (80%)	84 (17%)	20 (4%)	5	45
1	B	517/566 (91%)	413 (80%)	82 (16%)	22 (4%)	4	42
1	C	510/566 (90%)	410 (80%)	77 (15%)	23 (4%)	4	40
1	D	512/566 (90%)	418 (82%)	79 (15%)	15 (3%)	7	54
All	All	2046/2264 (90%)	1644 (80%)	322 (16%)	80 (4%)	5	45

5 of 80 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	96	GLY
1	A	97	LEU
1	A	416	GLU
1	A	420	TRP
1	A	497	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	470/501 (94%)	446 (95%)	24 (5%)	33	79
1	B	480/501 (96%)	457 (95%)	23 (5%)	35	81
1	C	476/501 (95%)	454 (95%)	22 (5%)	37	82
1	D	477/501 (95%)	453 (95%)	24 (5%)	34	79
All	All	1903/2004 (95%)	1810 (95%)	93 (5%)	35	80

5 of 93 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	567	LYS
1	C	319	LEU
1	D	453	ASN
1	B	577	LEU
1	C	199	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 64 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	493	ASN
1	C	297	ASN
1	D	468	HIS
1	B	501	HIS
1	B	606	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	513/566 (90%)	-0.21	7 (1%)	72 38	11, 86, 199, 202	0
1	B	523/566 (92%)	-0.10	7 (1%)	74 40	44, 139, 202, 202	0
1	C	518/566 (91%)	-0.03	5 (0%)	79 47	36, 152, 202, 202	0
1	D	520/566 (91%)	-0.09	4 (0%)	83 53	11, 135, 202, 202	0
All	All	2074/2264 (91%)	-0.11	23 (1%)	77 44	11, 132, 202, 202	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	121	ARG	3.2
1	B	623	ARG	3.2
1	B	223	GLU	3.1
1	B	606	HIS	3.1
1	B	236	SER	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.