



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 04:32 AM GMT

PDB ID : 4B7U  
Title : PLASMODIUM FALCIPARUM L-LACTATE DEHYDROGENASE COM-  
PLEXED WITH BICINE  
Authors : Birkinshaw, R.W.; Brady, R.L.  
Deposited on : 2012-08-22  
Resolution : 1.88 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	5260 (1.90-1.86)
Clashscore	79885	6268 (1.90-1.86)
Ramachandran outliers	78287	6195 (1.90-1.86)
Sidechain outliers	78261	6196 (1.90-1.86)
RSRZ outliers	66119	5262 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	322	
1	B	322	
1	C	322	
1	D	322	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10481 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-LACTATE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	17	6	0
			2445	1557	418	457	13			
1	B	305	Total	C	N	O	S	9	4	0
			2340	1490	399	437	14			
1	C	305	Total	C	N	O	S	10	7	0
			2361	1505	404	439	13			
1	D	306	Total	C	N	O	S	10	8	0
			2379	1514	407	444	14			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	330	HIS	-	EXPRESSION TAG	UNP Q71T02
A	331	HIS	-	EXPRESSION TAG	UNP Q71T02
A	332	HIS	-	EXPRESSION TAG	UNP Q71T02
A	333	HIS	-	EXPRESSION TAG	UNP Q71T02
A	334	HIS	-	EXPRESSION TAG	UNP Q71T02
A	335	HIS	-	EXPRESSION TAG	UNP Q71T02
B	330	HIS	-	EXPRESSION TAG	UNP Q71T02
B	331	HIS	-	EXPRESSION TAG	UNP Q71T02
B	332	HIS	-	EXPRESSION TAG	UNP Q71T02
B	333	HIS	-	EXPRESSION TAG	UNP Q71T02
B	334	HIS	-	EXPRESSION TAG	UNP Q71T02
B	335	HIS	-	EXPRESSION TAG	UNP Q71T02
C	330	HIS	-	EXPRESSION TAG	UNP Q71T02
C	331	HIS	-	EXPRESSION TAG	UNP Q71T02
C	332	HIS	-	EXPRESSION TAG	UNP Q71T02
C	333	HIS	-	EXPRESSION TAG	UNP Q71T02
C	334	HIS	-	EXPRESSION TAG	UNP Q71T02
C	335	HIS	-	EXPRESSION TAG	UNP Q71T02
D	330	HIS	-	EXPRESSION TAG	UNP Q71T02
D	331	HIS	-	EXPRESSION TAG	UNP Q71T02
D	332	HIS	-	EXPRESSION TAG	UNP Q71T02

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	333	HIS	-	EXPRESSION TAG	UNP Q71T02
D	334	HIS	-	EXPRESSION TAG	UNP Q71T02
D	335	HIS	-	EXPRESSION TAG	UNP Q71T02

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		
2	D	1	Total	Ca	0	0
			1	1		
2	C	1	Total	Ca	0	0
			1	1		

- Molecule 3 is BICINE (three-letter code: BCN) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			11	6	1	4		
3	B	1	Total	C	N	O	0	0
			11	6	1	4		
3	B	1	Total	C	N	O	0	0
			11	6	1	4		
3	B	1	Total	C	N	O	0	0
			11	6	1	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	N	O	0	0
			11	6	1	4		
3	D	1	Total	C	N	O	0	0
			11	6	1	4		
3	D	1	Total	C	N	O	0	0
			11	6	1	4		
3	D	1	Total	C	N	O	0	0
			11	6	1	4		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	266	Total	O	0	0
			266	266		
6	B	211	Total	O	0	0
			211	211		
6	C	196	Total	O	0	0
			196	196		
6	D	171	Total	O	0	0
			171	171		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: L-LACTATE DEHYDROGENASE

Chain A:




#### • Molecule 1: L-LACTATE DEHYDROGENASE

Chain B:




#### • Molecule 1: L-LACTATE DEHYDROGENASE

Chain C:



#### • Molecule 1: L-LACTATE DEHYDROGENASE

Chain D:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	155.29Å 154.78Å 58.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.96 – 1.88 48.96 – 1.88	Depositor EDS
% Data completeness (in resolution range)	99.4 (48.96-1.88) 99.5 (48.96-1.88)	Depositor EDS
$R_{merge}$	0.01	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.17 (at 1.88Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.176 , 0.201 0.174 , 0.199	Depositor DCC
$R_{free}$ test set	5761 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.5	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 42.6	EDS
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	2 of 114851 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10481	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, CA, BCN, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.62	3/2483 (0.1%)	0.59	3/3364 (0.1%)
1	B	0.55	1/2375 (0.0%)	0.53	0/3217
1	C	0.56	0/2398	0.53	0/3250
1	D	0.57	2/2415 (0.1%)	0.58	2/3272 (0.1%)
All	All	0.58	6/9671 (0.1%)	0.56	5/13103 (0.0%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	109	ARG	CG-CD	-10.29	1.26	1.51
1	A	310	GLU	CG-CD	9.48	1.66	1.51
1	A	314	LYS	CD-CE	-7.08	1.33	1.51
1	B	109	ARG	CG-CD	-6.97	1.34	1.51
1	A	105(B)	LYS	CB-CG	-6.03	1.36	1.52
1	D	157	LYS	CG-CD	-5.14	1.34	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	109	ARG	CG-CD-NE	-8.76	93.41	111.80
1	A	310	GLU	CG-CD-OE1	-8.64	101.01	118.30
1	A	310	GLU	CG-CD-OE2	8.34	134.98	118.30
1	D	109	ARG	CB-CG-CD	7.75	131.74	111.60
1	A	314	LYS	CG-CD-CE	5.04	127.01	111.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2445	0	2516	13	0
1	B	2340	0	2411	6	0
1	C	2361	0	2436	9	0
1	D	2379	0	2445	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	11	0	11	0	0
3	B	33	0	35	0	0
3	C	11	0	11	0	0
3	D	33	0	35	0	0
4	A	8	0	14	5	0
5	A	8	0	12	0	0
5	D	4	0	6	0	0
6	A	266	0	0	1	0
6	B	211	0	0	0	0
6	C	196	0	0	0	0
6	D	171	0	0	0	0
All	All	10481	0	9932	31	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (31) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:103(E):SER:C	1:A:105(A):ASP:N	2.01	1.14
1:A:118:LYS:HZ1	4:A:1332:MPD:HM1	1.73	0.53
1:A:30:MET:HE3	1:A:247:TYR:CZ	2.47	0.50
1:A:118:LYS:NZ	4:A:1332:MPD:HM1	2.27	0.50
1:C:69:THR:HG21	1:D:248:VAL:HG12	1.94	0.49
1:A:132(B):ASN:O	1:A:159:LYS:NZ	2.43	0.47
1:D:271[B]:CYS:SG	1:D:291:VAL:HG21	2.55	0.47
1:B:271[B]:CYS:SG	1:B:291:VAL:HG21	2.55	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:142:VAL:CG1	1:B:195:HIS:HB3	2.45	0.46
1:A:248:VAL:HG12	1:B:69:THR:HG21	1.98	0.46
1:C:30:MET:CE	1:D:30:MET:CE	2.95	0.45
1:A:72:MET:CE	1:B:253:ALA:HB2	2.47	0.45
1:D:142[A]:VAL:HG12	1:D:195:HIS:HB3	2.00	0.44
1:B:30:MET:HE2	1:B:247:TYR:CZ	2.53	0.44
1:A:118:LYS:NZ	4:A:1332:MPD:C5	2.82	0.43
1:A:118:LYS:HZ1	4:A:1332:MPD:C5	2.33	0.42
1:C:30:MET:HE2	1:C:247:TYR:CZ	2.54	0.42
1:C:69:THR:HG22	1:D:249:ALA:HA	2.01	0.42
1:C:30:MET:HE3	1:D:30:MET:HE1	2.00	0.42
1:D:142[A]:VAL:HG11	1:D:163:LEU:O	2.19	0.42
1:C:285:ILE:HD13	1:C:323:LYS:HG2	2.02	0.42
1:D:40:ILE:HG21	1:D:49:VAL:HG22	2.00	0.42
1:A:69:THR:HG21	1:B:248:VAL:HG12	2.02	0.42
1:A:103(E):SER:C	1:A:105(A):ASP:CA	2.84	0.42
1:C:30:MET:HE3	1:D:30:MET:CE	2.50	0.42
1:A:310:GLU:HG3	6:A:2251:HOH:O	2.19	0.41
1:C:40:ILE:HG21	1:C:49:VAL:CG2	2.50	0.41
1:A:118:LYS:NZ	4:A:1332:MPD:H53	2.36	0.40
1:D:43:LYS:HB2	1:D:45[B]:LEU:HG	2.02	0.40
1:D:203:LYS:NZ	1:D:221:ASP:OD1	2.52	0.40
1:C:191:ILE:HG12	1:C:201:LEU:HG	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	318/322 (99%)	310 (98%)	7 (2%)	1 (0%)	50 36
1	B	305/322 (95%)	298 (98%)	6 (2%)	1 (0%)	50 36
1	C	308/322 (96%)	299 (97%)	8 (3%)	1 (0%)	50 36
1	D	310/322 (96%)	305 (98%)	4 (1%)	1 (0%)	50 36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1241/1288 (96%)	1212 (98%)	25 (2%)	4 (0%)	50 36

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	164	GLY
1	D	164	GLY
1	A	164	GLY
1	C	164	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	269/269 (100%)	265 (98%)	4 (2%)	76 70
1	B	258/269 (96%)	254 (98%)	4 (2%)	75 68
1	C	260/269 (97%)	257 (99%)	3 (1%)	82 78
1	D	263/269 (98%)	261 (99%)	2 (1%)	89 88
All	All	1050/1076 (98%)	1037 (99%)	13 (1%)	87 78

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73(B)[A]	TYR
1	A	73(B)[B]	TYR
1	A	140	ASN
1	A	245	SER
1	B	73(B)[A]	TYR
1	B	73(B)[B]	TYR
1	B	140	ASN
1	B	181	ASN
1	C	72	MET
1	C	73(B)[A]	TYR
1	C	73(B)[B]	TYR
1	D	73(B)[A]	TYR
1	D	73(B)[B]	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	243	HIS
1	C	178	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	BCN	A	1331	2	10,10,10	0.68	0	11,11,11	0.82	0
4	MPD	A	1332	-	7,7,7	0.28	0	10,10,10	0.36	0
5	EDO	A	1333	-	3,3,3	0.56	0	2,2,2	0.32	0
5	EDO	A	1334	-	3,3,3	0.51	0	2,2,2	0.35	0
3	BCN	B	1331	2	10,10,10	0.66	0	11,11,11	0.93	0
3	BCN	B	1332	-	10,10,10	0.71	0	11,11,11	0.75	0
3	BCN	B	1333	-	10,10,10	0.65	0	11,11,11	0.82	0
3	BCN	C	1331	2	10,10,10	0.62	0	11,11,11	0.95	0
3	BCN	D	1332	2	10,10,10	0.73	0	11,11,11	0.70	0
3	BCN	D	1333	-	10,10,10	0.75	0	11,11,11	0.78	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	BCN	D	1334	-	10,10,10	0.62	0	11,11,11	0.88	0
5	EDO	D	1335	-	3,3,3	0.53	0	2,2,2	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BCN	A	1331	2	-	0/10/10/10	0/0/0/0
4	MPD	A	1332	-	-	0/5/5/5	0/0/0/0
5	EDO	A	1333	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1334	-	-	0/1/1/1	0/0/0/0
3	BCN	B	1331	2	-	0/10/10/10	0/0/0/0
3	BCN	B	1332	-	-	0/10/10/10	0/0/0/0
3	BCN	B	1333	-	-	0/10/10/10	0/0/0/0
3	BCN	C	1331	2	-	0/10/10/10	0/0/0/0
3	BCN	D	1332	2	-	0/10/10/10	0/0/0/0
3	BCN	D	1333	-	-	0/10/10/10	0/0/0/0
3	BCN	D	1334	-	-	0/10/10/10	0/0/0/0
5	EDO	D	1335	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	316/322 (98%)	-0.17	6 (1%) 64 64	11, 18, 27, 44	7 (2%)
1	B	305/322 (94%)	-0.09	9 (2%) 48 48	12, 19, 29, 38	3 (0%)
1	C	305/322 (94%)	-0.04	12 (3%) 37 37	12, 19, 29, 40	3 (0%)
1	D	306/322 (95%)	-0.14	4 (1%) 74 75	12, 19, 28, 38	4 (1%)
All	All	1232/1288 (95%)	-0.11	31 (2%) 52 55	11, 19, 29, 44	17 (1%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	18	ALA	6.9
1	B	18	ALA	6.1
1	A	73(B)[A]	TYR	5.4
1	A	18	ALA	4.8
1	D	73(B)[A]	TYR	4.7
1	D	18	ALA	4.2
1	D	331	HIS	3.5
1	C	100	PHE	3.5
1	B	100	PHE	3.4
1	B	115	LEU	3.2
1	D	330	HIS	3.1
1	A	245	SER	3.1
1	C	73(B)[A]	TYR	3.1
1	B	111	ASP	3.1
1	C	328	LEU	2.9
1	C	115	LEU	2.9
1	C	111	ASP	2.8
1	C	112	LEU	2.7
1	C	19	PRO	2.5
1	B	73(B)[A]	TYR	2.4
1	A	243	HIS	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	19	PRO	2.2
1	B	329	ALA	2.2
1	C	118	LYS	2.2
1	C	101	THR	2.2
1	A	105(A)	ASP	2.2
1	B	112	LEU	2.1
1	C	309	SER	2.1
1	A	330	HIS	2.1
1	C	54	ILE	2.0
1	B	110	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	EDO	A	1334	4/4	0.14	-	49,50,50,50	0
3	BCN	D	1332	11/11	0.23	-	32,36,37,39	0
3	BCN	B	1332	11/11	0.15	-	31,35,38,40	0
3	BCN	C	1331	11/11	0.22	-	38,39,40,41	0
3	BCN	B	1333	11/11	0.25	-	41,48,50,50	0
2	CA	C	401	1/1	0.33	-	61,61,61,61	0
5	EDO	D	1335	4/4	0.14	-	43,43,43,43	0
3	BCN	D	1333	11/11	0.27	-	62,64,65,65	0
5	EDO	A	1333	4/4	0.17	-	54,54,54,54	0
4	MPD	A	1332	8/8	0.23	-	50,52,53,53	0
3	BCN	D	1334	11/11	0.26	-	36,42,46,46	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	A	401	1/1	0.15	-	48,48,48,48	0
2	CA	B	401	1/1	0.28	-	66,66,66,66	0
3	BCN	B	1331	11/11	0.20	-	33,35,36,38	0
3	BCN	A	1331	11/11	0.13	-	23,25,26,27	0
2	CA	D	401	1/1	0.29	-	78,78,78,78	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.