



# wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 04:01 PM GMT

PDB ID : 2B92  
Title : Crystal-structure of the N-terminal Large GTPase Domain of human Guanylate Binding protein 1 (hGBP1) in complex with GDP/AlF3  
Authors : Ghosh, A.; Praefcke, G.J.K.; Renault, L.; Wittinghofer, A.; Herrmann, C.  
Deposited on : 2005-10-10  
Resolution : 3.20 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1824 (3.30-3.10)
Clashscore	79885	1078 (3.26-3.14)
Ramachandran outliers	78287	1059 (3.26-3.14)
Sidechain outliers	78261	1058 (3.26-3.14)
RSRZ outliers	66119	1825 (3.30-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	328	
1	B	328	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	MG	B	595	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4666 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interferon-induced guanylate-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	288	Total	C	N	O	S	0	0	0
			2284	1466	385	421	12			
1	B	290	Total	C	N	O	S	0	0	0
			2301	1478	387	424	12			


There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	CLONING ARTIFACT	UNP P32455
A	-9	HIS	-	EXPRESSION TAG	UNP P32455
A	-8	HIS	-	EXPRESSION TAG	UNP P32455
A	-7	HIS	-	EXPRESSION TAG	UNP P32455
A	-6	HIS	-	EXPRESSION TAG	UNP P32455
A	-5	HIS	-	EXPRESSION TAG	UNP P32455
A	-4	HIS	-	EXPRESSION TAG	UNP P32455
A	-3	MET	-	CLONING ARTIFACT	UNP P32455
A	-2	ARG	-	CLONING ARTIFACT	UNP P32455
A	-1	GLY	-	CLONING ARTIFACT	UNP P32455
A	0	SER	-	CLONING ARTIFACT	UNP P32455
B	-10	MET	-	CLONING ARTIFACT	UNP P32455
B	-9	HIS	-	EXPRESSION TAG	UNP P32455
B	-8	HIS	-	EXPRESSION TAG	UNP P32455
B	-7	HIS	-	EXPRESSION TAG	UNP P32455
B	-6	HIS	-	EXPRESSION TAG	UNP P32455
B	-5	HIS	-	EXPRESSION TAG	UNP P32455
B	-4	HIS	-	EXPRESSION TAG	UNP P32455
B	-3	MET	-	CLONING ARTIFACT	UNP P32455
B	-2	ARG	-	CLONING ARTIFACT	UNP P32455
B	-1	GLY	-	CLONING ARTIFACT	UNP P32455
B	0	SER	-	CLONING ARTIFACT	UNP P32455

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula:  $C_{10}H_{15}N_5O_{11}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
3	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 4 is ALUMINUM FLUORIDE (three-letter code: AF3) (formula:  $AlF_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Al	F	0	0
			4	1	3		
4	B	1	Total	Al	F	0	0
			4	1	3		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	9	Total	O	0	0
			9	9		
5	B	6	Total	O	0	0
			6	6		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interferon-induced guanylate-binding protein 1

Chain A:



- Molecule 1: Interferon-induced guanylate-binding protein 1

Chain B:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	148.82Å 103.92Å 55.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.20 29.62 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.3 (30.00-3.20) 99.3 (29.62-3.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.18 (at 3.18Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.204 , 0.250 0.206 , 0.252	Depositor DCC
$R_{free}$ test set	989 reflections (7.22%)	DCC
Wilson B-factor (Å <sup>2</sup> )	79.0	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 30.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 14695 reflections (0.007%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4666	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 65.95 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 6.5306e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MG, AF3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.26	13/2334 (0.6%)	0.40	0/3156
1	B	1.31	19/2352 (0.8%)	0.44	2/3180 (0.1%)
All	All	1.28	32/4686 (0.7%)	0.42	2/6336 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	5
All	All	0	8

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	85	HIS	N-CA	-10.08	1.26	1.46
1	B	84	PRO	N-CA	-8.45	1.32	1.47
1	A	26	GLU	CB-CG	8.39	1.68	1.52
1	B	147	GLU	CB-CG	8.05	1.67	1.52
1	A	251	GLU	CG-CD	7.92	1.63	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	210	LYS	C-N-CA	-6.33	109.02	122.30
1	B	219	PHE	O-C-N	-5.59	113.75	122.70

There are no chirality outliers.



5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	212	THR	Peptide
1	A	243	HIS	Peptide
1	A	245	ARG	Peptide
1	B	154	SER	Mainchain
1	B	66	SER	Peptide

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2284	0	2308	69	0
1	B	2301	0	2323	77	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	28	0	12	1	0
3	B	28	0	12	1	0
4	A	4	0	0	0	0
4	B	4	0	0	0	0
5	A	9	0	0	1	0
5	B	6	0	0	1	0
All	All	4666	0	4655	144	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 15.

The worst 5 of 144 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:219:PHE:O	1:A:222:PRO:HD2	1.39	1.19
1:B:171:PHE:O	1:B:172:VAL:HG12	1.42	1.16
1:A:49:THR:HG23	1:A:129:ASN:ND2	1.63	1.12
1:B:244:ARG:HG3	1:B:244:ARG:HH11	1.11	1.11
1:B:49:THR:HG23	1:B:129:ASN:ND2	1.76	1.01

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/328 (87%)	252 (89%)	27 (10%)	5 (2%)	13	60
1	B	286/328 (87%)	250 (87%)	24 (8%)	12 (4%)	4	31
All	All	570/656 (87%)	502 (88%)	51 (9%)	17 (3%)	7	42

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	LYS
1	A	219	PHE
1	A	220	ASN
1	B	155	LYS
1	B	219	PHE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/293 (88%)	229 (89%)	28 (11%)	9	37
1	B	259/293 (88%)	219 (85%)	40 (15%)	4	18
All	All	516/586 (88%)	448 (87%)	68 (13%)	6	27

5 of 68 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	14	ILE
1	B	66	SER
1	B	252	LYS
1	B	17	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	47	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	111	ASN
1	B	214	GLN
1	B	129	ASN
1	B	18	ASN
1	B	194	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GDP	A	593	2,4	30,30,30	2.07	9 (30%)	44,47,47	2.37	8 (18%)
4	AF3	A	594	3,2,5	0,3,3	0.00	-	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GDP	B	593	2,4	30,30,30	2.15	11 (36%)	44,47,47	2.27	8 (18%)
4	AF3	B	594	3,2,5	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	593	2,4	-	0/16/32/32	0/1/3/3
4	AF3	A	594	3,2,5	-	0/0/0/0	0/0/0/0
3	GDP	B	593	2,4	-	0/16/32/32	0/1/3/3
4	AF3	B	594	3,2,5	-	0/0/0/0	0/0/0/0

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	593	GDP	O4'-C1'	5.24	1.49	1.41
3	B	593	GDP	O4'-C1'	5.02	1.49	1.41
3	B	593	GDP	PB-O2B	-4.12	1.39	1.54
3	A	593	GDP	PB-O2B	-4.12	1.39	1.54
3	B	593	GDP	PB-O3B	-4.00	1.40	1.54

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	593	GDP	C6-C5-N7	-12.02	132.52	134.14
3	B	593	GDP	C6-C5-N7	-11.10	132.65	134.14
3	B	593	GDP	C2-N3-C4	4.03	120.75	115.09
3	A	593	GDP	C2-N3-C4	3.98	120.68	115.09
3	A	593	GDP	PA-O3A-PB	-3.89	120.26	131.68

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	288/328 (87%)	-0.18	0	100 100	33, 58, 90, 98	0
1	B	290/328 (88%)	-0.16	2 (0%)	84 38	32, 58, 90, 99	0
All	All	578/656 (88%)	-0.17	2 (0%)	91 58	32, 58, 90, 99	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	307	GLY	2.3
1	B	308	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	B	595	1/1	0.27	2.68	43,43,43,43	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	A	595	1/1	0.21	0.70	40,40,40,40	0
4	AF3	A	594	4/4	0.21	0.48	32,37,38,38	0
4	AF3	B	594	4/4	0.22	0.34	30,34,36,37	0
3	GDP	A	593	28/28	0.19	0.03	46,49,55,59	0
3	GDP	B	593	28/28	0.16	-1.11	47,51,55,56	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.