



wwPDB X-ray Structure Validation Summary Report

Oct 21, 2014 – 05:31 PM EDT

PDB ID : 2BCC
Title : STIGMATELLIN-BOUND CYTOCHROME BC1 COMPLEX FROM CHICKEN
Authors : Zhang, Z.; Huang, L.; Shulmeister, V.M.; Chi, Y.I.; Kim, K.K.; Hung, L.W.; Crofts, A.R.; Berry, E.A.; Kim, S.H.
Deposited on : 1998-09-18
Resolution : 3.50 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable24103
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable24103

1 Overall quality at a glance

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1039 (3.66-3.34)
Ramachandran outliers	78287	1000 (3.66-3.34)
Sidechain outliers	78261	1000 (3.66-3.34)
RSRZ outliers	66119	1243 (3.70-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	446	
2	B	422	
3	C	380	
4	D	241	
5	E	196	
6	F	109	
7	G	81	
8	H	78	
9	I	33	
10	J	62	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
11	BOG	D	242	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
12	HEM	C	381	-	X
12	HEM	C	382	-	X
12	HEM	D	243	-	X
13	FES	E	197	-	X
14	U10	C	383	-	X
15	PEE	C	384	-	X
15	PEE	E	198	-	X
16	SIG	C	385	-	X

2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 15754 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UBIQUINOL CYTOCHROME C OXIDOREDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	442	Total	C	N	O	S	0	0	0
			3423	2147	601	657	18			

- Molecule 2 is a protein called UBIQUINOL CYTOCHROME C OXIDOREDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	406	Total	C	N	O	S	0	0	0
			2994	1878	518	591	7			

- Molecule 3 is a protein called UBIQUINOL CYTOCHROME C OXIDOREDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	379	Total	C	N	O	S	0	0	0
			3002	2013	473	504	12			

- Molecule 4 is a protein called UBIQUINOL CYTOCHROME C OXIDOREDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	241	Total	C	N	O	S	0	0	0
			1899	1214	326	345	14			

- Molecule 5 is a protein called UBIQUINOL CYTOCHROME C OXIDOREDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	196	Total	C	N	O	S	0	0	0
			1512	953	266	285	8			

- Molecule 6 is a protein called UBIQUINOL CYTOCHROME C OXIDOREDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	100	Total	C	N	O	S	0	0	0
			875	557	153	162	3			

- Molecule 7 is a protein called UBIQUINOL CYTOCHROME C OXIDOREDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	78	Total	C	N	O	S	0	0	0
			626	411	114	100	1			

- Molecule 8 is a protein called UBIQUINOL CYTOCHROME C OXIDOREDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	66	Total	C	N	O	S	0	0	0
			490	301	88	96	5			


- Molecule 9 is a protein called UBIQUINOL CYTOCHROME C OXIDOREDUCTASE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
9	I	33	Total	C	N	O	0	0	0
			159	92	33	34			

- Molecule 10 is a protein called UBIQUINOL CYTOCHROME C OXIDOREDUCTASE.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
10	J	59	Total	C	N	O	0	0	0
			459	299	78	82			

- Molecule 11 is SUGAR (B-OCTYLGLUCOSIDE) (three-letter code: BOG) (formula: C₁₄H₂₈O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	D	1	Total	C	O	0	0
			20	14	6		

- Molecule 12 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
12	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
12	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 13 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
13	E	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 14 is UBIQUINONE-10 (three-letter code: U10) (formula: $C_{59}H_{90}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
14	C	1	Total	C	O	0	0
			29	25	4		

- Molecule 15 is 1,2-DIOLEOYL-SN-GLYCERO-3-PHOSPHOETHANOLAMINE (three-letter code: PEE) (formula: $C_{41}H_{83}NO_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
15	E	1	Total	C	N	O	P	0	0
			49	39	1	8	1		
15	C	1	Total	C	N	O	P	0	0
			49	39	1	8	1		

- Molecule 16 is STIGMATELLIN (three-letter code: SIG) (formula: C₃₀H₄₂O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
16	C	1	Total	C	O	0	0
			35	30	5		


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: UBIQUINOL CYTOCHROME C OXIDOREDUCTASE




• Molecule 2: UBIQUINOL CYTOCHROME C OXIDOREDUCTASE






• Molecule 3: UBIQUINOL CYTOCHROME C OXIDOREDUCTASE

Chain C:




• Molecule 4: UBIQUINOL CYTOCHROME C OXIDOREDUCTASE

Chain D:




• Molecule 5: UBIQUINOL CYTOCHROME C OXIDOREDUCTASE

Chain E:




• Molecule 6: UBIQUINOL CYTOCHROME C OXIDOREDUCTASE

Chain F:





• Molecule 7: UBIQUINOL CYTOCHROME C OXIDOREDUCTASE

Chain G:




• Molecule 8: UBIQUINOL CYTOCHROME C OXIDOREDUCTASE

Chain H:




• Molecule 9: UBIQUINOL CYTOCHROME C OXIDOREDUCTASE

Chain I:



• Molecule 10: UBIQUINOL CYTOCHROME C OXIDOREDUCTASE

Chain J:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	173.46Å 182.45Å 241.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	12.00 – 3.50 87.05 – 3.02	Depositor EDS
% Data completeness (in resolution range)	85.6 (12.00-3.50) 62.8 (87.05-3.02)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 3.01Å)	Xtriage
Refinement program	CNS 0.1	Depositor
R, R_{free}	0.284 , 0.317 0.244 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	56.4	Xtriage
Anisotropy	0.283	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 19.3	EDS
Estimated twinning fraction	0.239 for k,h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.31$, $\langle L^2 \rangle = 0.17$	Xtriage
Outliers	0 of 118814 reflections	Xtriage
F_o, F_c correlation	0.61	EDS
Total number of atoms	15754	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: U10, SIG, FES, HEM, PEE, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/3495	0.78	1/4742 (0.0%)
2	B	0.43	0/3046	0.73	0/4132
3	C	0.52	0/3104	0.85	5/4252 (0.1%)
4	D	0.50	0/1960	0.81	1/2665 (0.0%)
5	E	0.46	0/1548	0.78	1/2095 (0.0%)
6	F	0.49	0/896	0.76	0/1206
7	G	0.53	0/648	1.17	3/882 (0.3%)
8	H	0.44	0/495	0.69	0/669
10	J	0.52	0/470	0.80	1/635 (0.2%)
All	All	0.48	0/15662	0.81	12/21278 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1
10	J	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	G	26	PHE	C-N-CD	-18.95	78.91	120.60
7	G	26	PHE	C-N-CA	13.72	179.63	122.00
7	G	27	PRO	CA-N-CD	-7.67	100.76	111.50
10	J	61	ASN	N-CA-C	6.56	128.72	111.00
3	C	267	PRO	N-CA-C	-6.18	96.03	112.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	76	TYR	Sidechain
10	J	59	TYR	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3423	0	3286	359	0
2	B	2994	0	2906	345	0
3	C	3002	0	3036	423	0
4	D	1899	0	1822	216	0
5	E	1512	0	1483	177	0
6	F	875	0	839	70	0
7	G	626	0	591	83	0
8	H	490	0	445	57	0
9	I	159	0	46	20	0
10	J	459	0	424	53	0
11	D	20	0	28	1	0
12	C	86	0	60	19	0
12	D	43	0	30	2	0
13	E	4	0	0	1	0
14	C	29	0	33	9	0
15	C	49	0	70	7	0
15	E	49	0	70	4	0
16	C	35	0	42	12	0
All	All	15754	0	15211	1654	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 53.

The worst 5 of 1654 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:C:166:TRP:HB2	3:C:175:THR:HB	1.22	1.17

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:36:THR:HG22	1:A:100:LYS:HB3	1.22	1.17
2:B:280:GLY:H	2:B:283:PRO:HD2	1.06	1.16
2:B:168:TYR:HB2	2:B:173:ALA:HB2	1.36	1.05
7:G:60:THR:HG22	7:G:64:GLN:HE21	1.19	1.01

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	440/446 (99%)	337 (77%)	85 (19%)	18 (4%)	4	44
2	B	404/422 (96%)	295 (73%)	78 (19%)	31 (8%)	1	22
3	C	377/380 (99%)	278 (74%)	68 (18%)	31 (8%)	1	21
4	D	239/241 (99%)	194 (81%)	34 (14%)	11 (5%)	4	39
5	E	194/196 (99%)	151 (78%)	28 (14%)	15 (8%)	1	22
6	F	98/109 (90%)	84 (86%)	12 (12%)	2 (2%)	11	63
7	G	76/81 (94%)	52 (68%)	18 (24%)	6 (8%)	1	22
8	H	64/78 (82%)	56 (88%)	7 (11%)	1 (2%)	14	68
10	J	57/62 (92%)	32 (56%)	20 (35%)	5 (9%)	1	19
All	All	1949/2015 (97%)	1479 (76%)	350 (18%)	120 (6%)	2	30

5 of 120 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	282	ARG
1	A	284	TYR
2	B	19	PRO
2	B	20	HIS
2	B	23	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/376 (96%)	333 (93%)	26 (7%)	21	67
2	B	307/336 (91%)	286 (93%)	21 (7%)	22	70
3	C	326/329 (99%)	298 (91%)	28 (9%)	15	58
4	D	201/207 (97%)	193 (96%)	8 (4%)	42	84
5	E	165/169 (98%)	153 (93%)	12 (7%)	20	66
6	F	90/98 (92%)	80 (89%)	10 (11%)	9	42
7	G	60/72 (83%)	53 (88%)	7 (12%)	8	38
8	H	51/74 (69%)	51 (100%)	0	100	100
10	J	41/52 (79%)	40 (98%)	1 (2%)	61	92
All	All	1600/1713 (93%)	1487 (93%)	113 (7%)	21	67

5 of 113 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	104	TYR
3	C	208	ASN
6	F	107	TRP
3	C	129	PHE
3	C	152	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 47 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	429	ASN
3	C	86	ASN
7	G	64	GLN
3	C	16	ASN
3	C	138	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
12	HEM	C	381	3	42,50,50	3.11	9 (21%)	27,82,82	2.56	10 (37%)
12	HEM	C	382	3	42,50,50	3.32	11 (26%)	27,82,82	2.65	10 (37%)
14	U10	C	383	-	29,29,63	3.58	7 (24%)	37,38,79	2.14	9 (24%)
15	PEE	C	384	-	48,48,50	2.51	12 (25%)	53,53,55	4.23	25 (47%)
16	SIG	C	385	-	36,36,36	2.80	13 (36%)	48,50,50	2.38	14 (29%)
11	BOG	D	242	-	20,20,20	1.12	2 (10%)	25,25,25	0.84	1 (4%)
12	HEM	D	243	4	42,50,50	3.37	13 (30%)	27,82,82	3.11	9 (33%)
13	FES	E	197	5	0,4,4	0.00	-	0,4,4	0.00	-
15	PEE	E	198	-	48,48,50	2.58	11 (22%)	53,53,55	4.23	23 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	HEM	C	381	3	-	0/14/114/114	0/0/8/8
12	HEM	C	382	3	-	0/14/114/114	0/0/8/8
14	U10	C	383	-	-	0/23/47/87	0/1/1/1
15	PEE	C	384	-	1/1/4/4	0/52/52/54	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
16	SIG	C	385	-	-	0/29/30/30	0/2/2/2
11	BOG	D	242	-	-	0/11/31/31	0/1/1/1
12	HEM	D	243	4	-	0/14/114/114	0/0/8/8
13	FES	E	197	5	-	0/0/4/4	0/1/1/1
15	PEE	E	198	-	1/1/4/4	0/52/52/54	0/0/0/0

The worst 5 of 78 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	C	382	HEM	C3C-C2C	-14.88	1.33	1.45
14	C	383	U10	C13-C14	14.73	1.62	1.32
12	D	243	HEM	C3B-C2B	-12.32	1.33	1.45
12	C	381	HEM	C3B-C2B	-12.29	1.33	1.45
12	D	243	HEM	C3C-C2C	-11.95	1.36	1.45

The worst 5 of 101 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	C	384	PEE	O4-C10-C11	-17.93	51.07	123.78
15	E	198	PEE	O4-C10-C11	-17.89	51.22	123.78
15	E	198	PEE	O3-C30-C31	12.96	151.56	111.90
15	C	384	PEE	O3-C30-C31	12.89	151.35	111.90
15	E	198	PEE	O3-C30-O5	-10.86	95.11	123.48

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
15	E	198	PEE	C2
15	C	384	PEE	C2

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/446 (99%)	7.66	432 (97%) 0 0	12, 58, 89, 100	0
2	B	406/422 (96%)	7.68	396 (97%) 0 0	36, 71, 100, 100	0
3	C	379/380 (99%)	7.39	377 (99%) 0 0	4, 32, 71, 91	0
4	D	241/241 (100%)	7.51	238 (98%) 0 0	10, 41, 77, 100	0
5	E	196/196 (100%)	6.74	181 (92%) 0 0	17, 76, 100, 100	0
6	F	100/109 (91%)	7.69	100 (100%) 0 0	18, 42, 74, 99	0
7	G	78/81 (96%)	7.61	78 (100%) 0 0	26, 54, 89, 100	0
8	H	66/78 (84%)	8.55	66 (100%) 0 0	27, 67, 86, 88	0
9	I	0/33	-	-	-	-
10	J	59/62 (95%)	7.77	56 (94%) 0 0	37, 53, 86, 100	0
All	All	1967/2048 (96%)	7.53	1924 (97%) 0 0	4, 55, 98, 100	0

The worst 5 of 1924 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	30	PRO	23.0
2	B	44	GLY	19.7
2	B	222	GLN	19.6
7	G	44	CYS	19.5
1	A	269	ALA	19.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
13	FES	E	197	4/4	2.66	2.15	59,60,64,65	0
12	HEM	D	243	43/43	1.86	0.54	10,21,32,35	0
12	HEM	C	381	43/43	1.58	0.24	6,24,32,43	0
12	HEM	C	382	43/43	1.54	-0.01	8,20,32,41	0
15	PEE	C	384	49/51	1.59	-0.11	42,58,76,87	0
11	BOG	D	242	20/20	1.58	-0.12	35,68,81,81	0
16	SIG	C	385	35/35	1.39	-0.49	2,17,26,29	0
15	PEE	E	198	49/51	1.41	-0.57	43,80,98,100	0
14	U10	C	383	29/63	1.13	-1.22	72,88,100,100	0

6.5 Other polymers ⓘ

There are no such residues in this entry.