



Full wwPDB X-ray Structure Validation Report i

Feb 28, 2014 – 11:00 PM GMT

PDB ID : 2BCJ
Title : Crystal Structure of G Protein-Coupled Receptor Kinase 2 in Complex with
Galpha-q and Gbetagamma Subunits
Authors : Tesmer, J.J.G.
Deposited on : 2005-10-19
Resolution : 3.06 Å (reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2629 (3.12-3.00)
Ramachandran outliers	78287	2536 (3.12-3.00)
Sidechain outliers	78261	2539 (3.12-3.00)
RSRZ outliers	66119	2081 (3.12-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	689	
2	B	340	
3	G	71	
4	Q	353	

2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 10878 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called G-protein coupled receptor kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	624	5113	3263	889	925	36	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	670	ALA	SER	ENGINEERED	UNP P21146

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) beta subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	340	2610	1609	468	512	21	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	ACE	-	ACETYLATION	UNP P62871

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) gamma-2 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	G	64	501	316	88	94	3	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	68	SER	CYS	ENGINEERED	UNP P63212

- Molecule 4 is a protein called G alpha I1, Guanine nucleotide-binding protein G(q), alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	Q	317	2615	1671	440	492	12	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Q	35	ARG	-	CLONING ARTIFACT	UNP P21279
Q	36	SER	-	CLONING ARTIFACT	UNP P21279

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	Q	1	Total	Mg	0	0
			1	1		

- Molecule 6 is TETRAFLUOROALUMINATE ION (three-letter code: ALF) (formula: AlF_4^-).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Al	F		
6	Q	1	5	1	4	0	0

- Molecule 7 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $\text{C}_{10}\text{H}_{15}\text{N}_5\text{O}_{11}\text{P}_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
7	Q	1	28	10	5	11	2	0	0

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
8	Q	5	5	5	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: G-protein coupled receptor kinase 2

Chain A:




- Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(T) beta subunit 1

Chain B:



- Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(O) gamma-2 subunit

Chain G:



- Molecule 4: G alpha I1, Guanine nucleotide-binding protein G(q), alpha subunit

Chain Q:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.92Å 129.95Å 122.76Å 90.00° 95.81° 90.00°	Depositor
Resolution (Å)	46.78 – 3.06 46.80 – 3.06	Depositor EDS
% Data completeness (in resolution range)	98.8 (46.78-3.06) 98.8 (46.80-3.06)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 3.06Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.243 , 0.292 0.246 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	82.5	Xtrriage
Anisotropy	0.325	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 59.2	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Outliers	1 of 37811 reflections (0.003%)	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10878	wwPDB-VP
Average B, all atoms (Å ²)	103.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, ALF, MG, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/5228	0.49	0/7025
2	B	0.33	0/2655	0.53	0/3599
3	G	0.36	0/509	0.44	0/685
4	Q	0.33	0/2671	0.49	0/3610
All	All	0.34	0/11063	0.50	0/14919

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5113	0	5116	17	0
2	B	2610	0	2514	30	0
3	G	501	0	510	7	0
4	Q	2615	0	2582	11	0
5	Q	1	0	0	0	0
6	Q	5	0	0	0	0
7	Q	28	0	12	0	0
8	Q	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	10878	0	10734	58	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (58) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:Q:270:ILE:HD13	4:Q:344:VAL:HG13	1.80	0.62
4:Q:231:ALA:HB3	4:Q:234:GLU:HG3	1.82	0.61
1:A:350:SER:HA	1:A:361:VAL:HG12	1.83	0.61
2:B:295:ASN:HD22	2:B:295:ASN:N	1.99	0.60
2:B:25:CYS:HB2	2:B:259:GLN:HE22	1.67	0.59
4:Q:329:THR:HG22	4:Q:340:VAL:HG21	1.87	0.56
2:B:152:LEU:HD22	2:B:196:THR:HB	1.88	0.55
4:Q:196:LEU:CD2	4:Q:345:LYS:HD2	2.39	0.53
1:A:53:VAL:O	1:A:53:VAL:HG12	2.10	0.52
2:B:210:LEU:HD22	2:B:255:LEU:HD22	1.91	0.51
2:B:40:VAL:HG21	2:B:284:LEU:HD21	1.92	0.51
2:B:237:ASN:OD1	2:B:238:GLY:N	2.44	0.50
3:G:56:ALA:O	3:G:63:GLU:HB2	2.12	0.49
1:A:48:GLU:HG3	1:A:53:VAL:HG21	1.93	0.49
2:B:7:LEU:HD21	3:G:13:ARG:HG3	1.94	0.49
4:Q:195:ASP:HB3	4:Q:200:ILE:HG12	1.94	0.49
2:B:54:HIS:CE1	2:B:80:ILE:HD12	2.48	0.49
4:Q:336:ASN:O	4:Q:340:VAL:HG23	2.12	0.47
2:B:254:ASP:HB2	2:B:261:LEU:HD11	1.97	0.47
4:Q:275:LYS:HD3	4:Q:278:LEU:HD12	1.96	0.47
2:B:254:ASP:OD2	3:G:33:ALA:HB1	2.15	0.46
4:Q:125:GLU:O	4:Q:129:VAL:HG23	2.16	0.46
2:B:45:MET:HB3	2:B:308:LEU:HD22	1.96	0.46
2:B:124:TYR:CE2	2:B:135:VAL:HG22	2.51	0.46
2:B:33:ILE:HD13	3:G:34:ALA:HB3	1.97	0.46
2:B:71:VAL:HG12	2:B:81:ILE:HG12	1.98	0.45
1:A:154:CYS:O	1:A:158:ARG:N	2.49	0.45
1:A:246:VAL:HG11	1:A:254:ILE:HG21	1.98	0.45
2:B:201:SER:O	2:B:208:ALA:HA	2.17	0.44
2:B:308:LEU:HD23	2:B:308:LEU:N	2.32	0.44
2:B:63:TRP:CZ2	2:B:328:ALA:HB2	2.53	0.44
1:A:135:HIS:CD2	1:A:140:GLN:HE21	2.36	0.44
2:B:231:ALA:CB	2:B:275:SER:HA	2.48	0.44
1:A:35:PRO:O	1:A:176:GLN:NE2	2.46	0.44
2:B:273:ILE:HD12	2:B:273:ILE:N	2.33	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:250:CYS:HB2	2:B:264:TYR:HB2	2.01	0.43
2:B:292:PHE:N	2:B:292:PHE:CD1	2.86	0.43
2:B:83:ASP:O	2:B:87:THR:N	2.52	0.43
4:Q:345:LYS:HE3	4:Q:349:LEU:HD12	2.02	0.42
2:B:51:LEU:HB3	2:B:82:TRP:CE3	2.54	0.42
1:A:600:LEU:C	1:A:600:LEU:HD12	2.39	0.42
2:B:191:SER:CB	2:B:233:CYS:HA	2.49	0.42
1:A:552:ASP:OD1	1:A:552:ASP:N	2.52	0.42
2:B:30:LEU:HD13	3:G:34:ALA:HB1	2.02	0.42
4:Q:232:LEU:HD21	4:Q:273:LEU:HD13	2.02	0.42
1:A:263:THR:HB	1:A:264:PRO:CD	2.50	0.41
2:B:338:ILE:HG21	3:G:61:PHE:CE2	2.55	0.41
2:B:55:LEU:HD11	2:B:78:LYS:NZ	2.35	0.41
2:B:40:VAL:HG23	3:G:51:LEU:HD11	2.02	0.41
1:A:137:VAL:HG22	4:Q:262:PRO:HA	2.02	0.41
2:B:183:HIS:HE2	2:B:201:SER:HG	1.68	0.41
2:B:87:THR:O	2:B:87:THR:HG22	2.20	0.41
1:A:648:ARG:O	1:A:652:ARG:HG3	2.20	0.41
1:A:197:ILE:HG22	1:A:197:ILE:O	2.20	0.41
1:A:212:ASP:OD1	1:A:212:ASP:N	2.54	0.41
1:A:262:HIS:HB3	1:A:267:LEU:HD12	2.04	0.40
1:A:42:VAL:HG11	1:A:639:GLU:HA	2.03	0.40
1:A:360:GLU:HB2	1:A:368:TYR:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	620/689 (90%)	581 (94%)	35 (6%)	4 (1%)	33 79
2	B	338/340 (99%)	315 (93%)	22 (6%)	1 (0%)	50 88
3	G	62/71 (87%)	59 (95%)	3 (5%)	0	100 100
4	Q	315/353 (89%)	300 (95%)	14 (4%)	1 (0%)	50 88

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1335/1453 (92%)	1255 (94%)	74 (6%)	6 (0%)	43 85

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	2	SER
1	A	335	ASP
4	Q	221	GLU
1	A	231	GLN
1	A	249	GLY
1	A	569	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	561/611 (92%)	546 (97%)	15 (3%)	57 90
2	B	282/282 (100%)	271 (96%)	11 (4%)	43 84
3	G	53/58 (91%)	53 (100%)	0	100 100
4	Q	290/321 (90%)	287 (99%)	3 (1%)	85 97
All	All	1186/1272 (93%)	1157 (98%)	29 (2%)	61 92

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	ARG
1	A	85	PHE
1	A	91	LYS
1	A	204	GLU
1	A	212	ASP
1	A	235	LEU
1	A	278	ASP
1	A	310	ASN
1	A	360	GLU
1	A	431	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	493	THR
1	A	568	MET
1	A	591	ARG
1	A	640	LEU
1	A	651	TYR
2	B	14	LEU
2	B	70	LEU
2	B	71	VAL
2	B	105	TYR
2	B	108	SER
2	B	154	ASP
2	B	172	GLU
2	B	194	PRO
2	B	292	PHE
2	B	295	ASN
2	B	322	ASP
4	Q	95	ASP
4	Q	148	ARG
4	Q	345	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	HIS
1	A	135	HIS
1	A	310	ASN
1	A	508	ASN
1	A	544	ASN
1	A	607	GLN
2	B	155	ASN
2	B	259	GLN
2	B	295	ASN
3	G	4	ASN
3	G	59	ASN
4	Q	109	HIS
4	Q	137	ASN
4	Q	266	ASN
4	Q	299	GLN

5.3.3 RNA i

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

There are no carbohydrates in this entry.

5.6 Ligand geometry i

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GDP	Q	360	5,6	30,30,30	1.46	4 (13%)	44,47,47	3.24	9 (20%)
6	ALF	Q	362	8,5,7	4,4,4	1.86	0	0,6,6	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GDP	Q	360	5,6	-	0/16/32/32	0/1/3/3
6	ALF	Q	362	8,5,7	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	Q	360	GDP	C6-C5	4.55	1.48	1.41
7	Q	360	GDP	C4-N9	-3.11	1.33	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	Q	360	GDP	C5-C4	3.01	1.47	1.40
7	Q	360	GDP	C2-N2	2.17	1.35	1.32

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	Q	360	GDP	C6-C5-N7	18.33	136.61	134.14
7	Q	360	GDP	N3-C4-N9	5.17	134.49	126.91
7	Q	360	GDP	C5-C4-N3	-4.82	118.96	125.94
7	Q	360	GDP	C2-N3-C4	4.20	120.99	115.09
7	Q	360	GDP	C4-C5-N7	-3.64	106.41	109.52
7	Q	360	GDP	C8-N9-C4	2.41	108.74	106.90
7	Q	360	GDP	N2-C2-N3	-2.20	117.32	120.30
7	Q	360	GDP	O4'-C1'-N9	2.15	110.44	108.44
7	Q	360	GDP	C3'-C2'-C1'	2.11	104.21	100.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	624/689 (90%)	0.03	0 100 100	83, 106, 131, 139	0
2	B	340/340 (100%)	-0.01	0 100 100	78, 90, 119, 130	0
3	G	64/71 (90%)	0.08	0 100 100	91, 105, 126, 128	0
4	Q	317/353 (89%)	0.02	0 100 100	90, 106, 123, 128	0
All	All	1345/1453 (92%)	0.02	0 100 100	78, 103, 126, 139	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	MG	Q	361	1/1	0.20	-0.03	87,87,87,87	0
6	ALF	Q	362	5/5	0.19	-0.45	79,81,83,83	0
7	GDP	Q	360	28/28	0.19	-0.60	71,77,78,78	0

6.5 Other polymers

There are no such residues in this entry.