



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 03:58 AM GMT

PDB ID : 4BCM
Title : Structure of CDK2 in complex with cyclin A and a 2-amino-4-heteroaryl
-pyrimidine inhibitor
Authors : Hole, A.J.; Baumli, S.; Wang, S.; Endicott, J.A.; Noble, M.E.M.
Deposited on : 2012-10-02
Resolution : 2.45 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	4471 (2.50-2.42)
Ramachandran outliers	78287	4383 (2.50-2.42)
Sidechain outliers	78261	4385 (2.50-2.42)
RSRZ outliers	66119	3568 (2.50-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	301	
1	C	301	
2	B	262	
3	D	262	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
5	SGM	B	1433	-	X
5	SGM	D	1429	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9111 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLIN-DEPENDENT KINASE 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	296	Total	C	N	O	P	S	0	0	0
			2375	1544	403	419	1	8			
1	C	295	Total	C	N	O	P	S	0	0	0
			2370	1538	400	423	1	8			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	PRO	-	EXPRESSION TAG	UNP P24941
A	-1	GLY	-	EXPRESSION TAG	UNP P24941
A	0	SER	-	EXPRESSION TAG	UNP P24941
C	-2	PRO	-	EXPRESSION TAG	UNP P24941
C	-1	GLY	-	EXPRESSION TAG	UNP P24941
C	0	SER	-	EXPRESSION TAG	UNP P24941

- Molecule 2 is a protein called CYCLIN-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	257	Total	C	N	O	S	0	1	0
			2083	1350	340	382	11			


- Molecule 3 is a protein called CYCLIN-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	236	Total	C	N	O	S	0	0	0
			1918	1249	311	348	10			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	331	LEU	SER	CONFLICT	UNP P20248

- Molecule 4 is 4-(4-METHYL-2-METHYLMINO-3H-1,3-THIAZOL-5-YL)-2-[(4-METHYL-3-MORPHOLIN-4-YLSULFONYL-PHENYL)AMINO]PYRIMIDINE-5-CARBONITRILE (three-letter code: T7Z) (formula: $C_{21}H_{23}N_7O_3S_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			33	21	7	3	2		
4	C	1	Total	C	N	O	S	0	0
			33	21	7	3	2		

- Molecule 5 is MONOTHIOGLYCEROL (three-letter code: SGM) (formula: $C_3H_8O_2S$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	O	S	0	0
			6	3	2	1		
5	D	1	Total	C	O	S	0	0
			6	3	2	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	128	Total	O	0	0
			128	128		
6	B	106	Total	O	0	0
			106	106		
6	C	36	Total	O	0	0
			36	36		
6	D	17	Total	O	0	0
			17	17		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CYCLIN-DEPENDENT KINASE 2

Chain A: 



- Molecule 1: CYCLIN-DEPENDENT KINASE 2

Chain C: 




- Molecule 2: CYCLIN-A2

Chain B: 



- Molecule 3: CYCLIN-A2

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.07Å 135.41Å 148.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.46 – 2.45 52.46 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.0 (52.46-2.45) 99.0 (52.46-2.45)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 2.45Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.196 , 0.258 0.188 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	41.1	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 40.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 55153 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9111	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, T7Z, SGM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2425	0.56	0/3288
1	C	0.29	0/2419	0.46	0/3282
2	B	0.37	0/2137	0.51	0/2901
3	D	0.29	0/1964	0.45	0/2661
All	All	0.34	0/8945	0.50	0/12132

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2375	0	17	16	0
1	C	2370	0	7	12	0
2	B	2083	0	0	16	0
3	D	1918	0	0	11	0
4	A	33	0	23	5	0
4	C	33	0	23	3	0
5	B	6	0	7	1	0
5	D	6	0	7	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	128	0	0	4	0
6	B	106	0	0	4	0
6	C	36	0	0	3	0
6	D	17	0	0	1	0
All	All	9111	0	84	52	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

All (52) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:193:CYS:SG	5:B:1433:SGM:S1	2.29	1.30
3:D:193:CYS:SG	5:D:1429:SGM:S1	2.31	1.28
1:A:-2:PRO:HA	1:A:68:ASP:OD2	1.83	0.77
1:A:89:LYS:NZ	4:A:1297:T7Z:H172	2.01	0.75
4:C:1295:T7Z:H061	4:C:1295:T7Z:C31	2.15	0.75
1:A:150:ARG:NH2	2:B:268:GLU:O	2.20	0.74
2:B:327:CYS:SG	6:B:2073:HOH:O	2.45	0.73
2:B:317:GLN:OE1	6:B:2016:HOH:O	2.06	0.73
4:A:1297:T7Z:H061	4:A:1297:T7Z:C31	2.21	0.68
1:C:204:PRO:O	6:C:2034:HOH:O	2.12	0.68
1:C:157:ARG:O	6:C:2024:HOH:O	2.11	0.68
1:A:41:THR:O	2:B:288:LYS:NZ	2.28	0.67
1:A:246:GLN:NE2	6:A:2107:HOH:O	2.25	0.67
2:B:203:GLN:O	6:B:2025:HOH:O	2.12	0.65
1:A:51:GLU:OE2	6:A:2016:HOH:O	2.16	0.61
3:D:303:THR:O	3:D:303:THR:CG2	2.49	0.60
3:D:203:GLN:O	6:D:2002:HOH:O	2.17	0.58
1:C:116:ALA:O	1:C:120:SER:OG	2.22	0.57
1:A:71:HIS:NE2	2:B:296:HIS:NE2	2.52	0.57
1:C:148:LEU:N	6:C:2019:HOH:O	2.38	0.56
4:A:1297:T7Z:C06	4:A:1297:T7Z:C31	2.85	0.54
4:C:1295:T7Z:C06	4:C:1295:T7Z:C31	2.86	0.54
3:D:305:ASP:CB	5:D:1429:SGM:H12	2.41	0.50
1:C:150:ARG:NH1	1:C:160:TPO:O2P	2.45	0.50
3:D:419:HIS:CD2	3:D:419:HIS:N	2.80	0.49
1:A:10:ILE:CG2	4:A:1297:T7Z:H211	2.43	0.49
1:A:71:HIS:CD2	2:B:296:HIS:NE2	2.81	0.48
3:D:383:THR:N	3:D:386:SER:OG	2.48	0.47
1:A:235:ASP:O	6:A:2099:HOH:O	2.20	0.47
1:C:71:HIS:ND1	3:D:296:HIS:NE2	2.63	0.46
1:A:73:GLU:CG	1:A:74:ASN:N	2.79	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:71:HIS:NE2	2:B:296:HIS:CD2	2.84	0.45
2:B:347:TYR:OH	2:B:397:THR:OG1	2.34	0.45
1:C:181:SER:OG	1:C:182:THR:N	2.50	0.45
1:A:71:HIS:CE1	2:B:296:HIS:CE1	3.05	0.44
1:C:101:LEU:N	1:C:102:PRO:CD	2.80	0.44
1:A:195:GLU:O	1:A:199:ARG:N	2.50	0.44
4:C:1295:T7Z:N32	4:C:1295:T7Z:H061	2.32	0.43
1:A:200:ARG:NH2	6:A:2088:HOH:O	2.51	0.43
2:B:177:ASP:OD1	2:B:177:ASP:O	2.37	0.43
1:C:165:THR:O	1:C:166:LEU:C	2.58	0.42
1:C:288:ASP:OD1	1:C:288:ASP:N	2.51	0.42
3:D:383:THR:O	3:D:386:SER:OG	2.38	0.42
1:C:67:LEU:N	1:C:79:VAL:O	2.53	0.41
2:B:228:GLN:N	2:B:269:GLU:OE2	2.53	0.41
2:B:274:GLU:N	2:B:277:GLU:OE1	2.54	0.41
3:D:415:ASN:OD1	3:D:416:SER:N	2.54	0.41
1:A:83:LEU:O	4:A:1297:T7Z:N11	2.54	0.41
2:B:412:LYS:NZ	6:B:2036:HOH:O	2.53	0.41
3:D:216:ASP:OD2	3:D:408:SER:OG	2.38	0.41
2:B:176:PRO:N	2:B:179[B]:HIS:CD2	2.89	0.41
1:C:150:ARG:NH2	3:D:268:GLU:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	291/301 (97%)	278 (96%)	12 (4%)	1 (0%)	50	70
1	C	292/301 (97%)	269 (92%)	20 (7%)	3 (1%)	22	35
2	B	256/262 (98%)	253 (99%)	3 (1%)	0	100	100
3	D	230/262 (88%)	219 (95%)	10 (4%)	1 (0%)	43	64
All	All	1069/1126 (95%)	1019 (95%)	45 (4%)	5 (0%)	38	57

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	166	LEU
1	C	164	VAL
1	A	164	VAL
1	C	256	ASP
3	D	420	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/264 (98%)	251 (97%)	8 (3%)	52	77
1	C	259/264 (98%)	254 (98%)	5 (2%)	69	89
2	B	232/236 (98%)	230 (99%)	2 (1%)	87	96
3	D	212/236 (90%)	209 (99%)	3 (1%)	78	93
All	All	962/1000 (96%)	944 (98%)	18 (2%)	69	89

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	LEU
1	A	73	GLU
1	A	122	ARG
1	A	150	ARG
1	A	206	ASP
1	A	248	PHE
1	A	295	HIS
1	A	296	LEU
2	B	209	SER
2	B	374	GLU
1	C	36	ARG
1	C	122	ARG
1	C	199	ARG
1	C	218	THR
1	C	248	PHE
3	D	338	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	D	398	TYR
3	D	424	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	TPO	A	160	1	10,10,11	5.23	1 (10%)	12,14,16	1.36	1 (8%)
1	TPO	C	160	1	10,10,11	5.33	1 (10%)	12,14,16	1.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	160	1	-	0/9/11/13	0/0/0/0
1	TPO	C	160	1	-	0/9/11/13	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	160	TPO	O-C	16.70	1.22	1.11
1	A	160	TPO	O-C	16.29	1.22	1.11

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	TPO	CG2-CB-CA	-2.31	108.46	113.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates i

There are no carbohydrates in this entry.

5.6 Ligand geometry i

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	T7Z	A	1297	-	36,36,36	3.82	16 (44%)	48,52,52	3.76	17 (35%)
5	SGM	B	1433	-	5,5,5	0.52	0	5,5,5	0.77	0
4	T7Z	C	1295	-	36,36,36	3.90	16 (44%)	48,52,52	3.24	20 (41%)
5	SGM	D	1429	-	5,5,5	0.39	0	5,5,5	1.48	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	T7Z	A	1297	-	-	0/19/32/32	0/4/4/4
5	SGM	B	1433	-	-	0/4/4/4	0/0/0/0
4	T7Z	C	1295	-	-	0/19/32/32	0/4/4/4
5	SGM	D	1429	-	-	0/4/4/4	0/0/0/0

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	1295	T7Z	C05-N04	11.24	1.52	1.38
4	A	1297	T7Z	C17-N16	-10.85	1.35	1.47
4	C	1295	T7Z	C17-N16	-10.82	1.35	1.47
4	A	1297	T7Z	C05-N04	10.81	1.51	1.38
4	C	1295	T7Z	C21-N16	-10.81	1.35	1.47
4	A	1297	T7Z	C21-N16	-10.64	1.35	1.47
4	C	1295	T7Z	C10-N11	6.00	1.45	1.36
4	A	1297	T7Z	C10-N11	5.21	1.44	1.36
4	C	1295	T7Z	C03-N04	5.13	1.42	1.35
4	A	1297	T7Z	C03-N04	4.73	1.42	1.35
4	C	1295	T7Z	C03-S33	-4.51	1.65	1.73
4	C	1295	T7Z	C03-N02	4.44	1.47	1.26
4	A	1297	T7Z	C03-N02	4.40	1.47	1.26
4	A	1297	T7Z	C03-S33	-4.18	1.66	1.73
4	A	1297	T7Z	C30-C31	3.49	1.50	1.44
4	A	1297	T7Z	C08-C07	3.28	1.54	1.50
4	A	1297	T7Z	C05-C07	3.23	1.45	1.37
4	A	1297	T7Z	C14-S15	3.18	1.82	1.77
4	C	1295	T7Z	C08-C07	3.16	1.54	1.50
4	C	1295	T7Z	C30-C31	3.14	1.49	1.44
4	C	1295	T7Z	C05-C07	3.12	1.45	1.37
4	C	1295	T7Z	C14-S15	2.97	1.81	1.77
4	A	1297	T7Z	C12-N11	2.94	1.47	1.40
4	C	1295	T7Z	C12-N11	2.93	1.47	1.40
4	C	1295	T7Z	C07-S33	-2.57	1.63	1.74
4	A	1297	T7Z	C07-S33	-2.51	1.64	1.74
4	A	1297	T7Z	C13-C14	2.50	1.43	1.39
4	A	1297	T7Z	C29-C30	2.45	1.42	1.38
4	A	1297	T7Z	C06-C05	2.25	1.53	1.49
4	C	1295	T7Z	C06-C05	2.22	1.53	1.49
4	C	1295	T7Z	C13-C14	2.20	1.42	1.39
4	C	1295	T7Z	C29-C30	2.08	1.42	1.38

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1297	T7Z	C03-N04-C05	-15.43	110.95	112.36
4	C	1295	T7Z	C03-N04-C05	-11.25	111.33	112.36
4	A	1297	T7Z	C21-N16-C17	10.07	124.61	112.16
4	C	1295	T7Z	N28-C10-N09	-7.81	119.96	126.68
4	C	1295	T7Z	C21-N16-C17	7.57	121.53	112.16
4	A	1297	T7Z	C20-C21-N16	6.78	112.86	108.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1297	T7Z	C07-C05-N04	6.60	109.95	106.82
4	A	1297	T7Z	N28-C10-N09	-6.32	121.24	126.68
4	A	1297	T7Z	O23-S15-O22	-5.82	108.99	119.38
4	C	1295	T7Z	C07-C05-N04	5.61	109.48	106.82
4	C	1295	T7Z	O23-S15-O22	-5.57	109.44	119.38
4	A	1297	T7Z	O22-S15-N16	5.39	112.20	106.71
4	C	1295	T7Z	C06-C05-C07	-5.07	127.26	131.18
4	A	1297	T7Z	C06-C05-C07	-4.94	127.36	131.18
4	C	1295	T7Z	C01-N02-C03	4.65	120.41	117.89
4	A	1297	T7Z	C17-N16-S15	-4.27	109.24	117.00
4	C	1295	T7Z	C17-N16-S15	-4.07	109.59	117.00
4	C	1295	T7Z	C18-C17-N16	4.00	110.95	108.22
4	C	1295	T7Z	C08-N09-C10	3.86	121.83	116.18
4	A	1297	T7Z	C18-C17-N16	3.84	110.84	108.22
4	C	1295	T7Z	C25-C24-C14	-3.46	120.53	124.10
4	A	1297	T7Z	C08-N09-C10	3.44	121.22	116.18
4	C	1295	T7Z	C26-C24-C14	3.37	120.68	116.23
4	A	1297	T7Z	C01-N02-C03	3.24	119.64	117.89
4	C	1295	T7Z	C30-C29-N28	-3.12	119.44	123.93
4	A	1297	T7Z	C30-C29-N28	-3.09	119.48	123.93
5	D	1429	SGM	C2-C1-S1	-3.00	109.93	114.66
4	A	1297	T7Z	C26-C24-C14	2.97	120.14	116.23
4	C	1295	T7Z	O23-S15-N16	2.89	109.66	106.71
4	C	1295	T7Z	C29-N28-C10	2.87	121.60	116.02
4	C	1295	T7Z	C21-N16-S15	2.72	121.94	117.00
4	C	1295	T7Z	O22-S15-N16	2.60	109.36	106.71
4	A	1297	T7Z	C29-N28-C10	2.53	120.93	116.02
4	C	1295	T7Z	C20-C21-N16	2.52	109.95	108.22
4	A	1297	T7Z	C30-C08-N09	-2.24	118.83	121.30
4	C	1295	T7Z	C13-C14-S15	2.16	120.16	117.50
4	C	1295	T7Z	C27-C26-C24	-2.07	119.08	122.02
4	A	1297	T7Z	C13-C14-S15	2.03	119.99	117.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	296/301 (98%)	-0.09	3 (1%) 79 80	19, 31, 83, 116	0
1	C	295/301 (98%)	0.65	37 (12%) 5 4	30, 68, 103, 118	0
2	B	257/262 (98%)	-0.13	3 (1%) 75 76	18, 33, 58, 93	0
3	D	236/262 (90%)	0.76	32 (13%) 4 3	27, 64, 99, 115	0
All	All	1084/1126 (96%)	0.29	75 (6%) 17 16	18, 47, 96, 118	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	418	TYR	6.0
3	D	372	TRP	5.4
3	D	332	LEU	5.2
3	D	423	LEU	5.0
1	C	215	ILE	5.0
1	C	227	TRP	4.5
1	C	234	PRO	4.4
3	D	333	ALA	4.3
3	D	319	PHE	4.1
1	A	-2	PRO	4.0
3	D	359	ALA	4.0
3	D	413	TYR	4.0
1	C	236	TYR	4.0
1	C	209	ILE	3.9
3	D	336	LEU	3.9
1	C	243	TRP	3.8
3	D	421	VAL	3.7
1	C	287	GLN	3.5
1	C	240	PHE	3.5
1	C	101	LEU	3.5
1	C	226	VAL	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	213	PHE	3.5
3	D	428	GLU	3.4
1	C	170	ALA	3.3
3	D	335	PHE	3.3
1	C	252	VAL	3.3
3	D	399	LEU	3.3
1	C	251	VAL	3.2
3	D	416	SER	3.2
1	C	173	ILE	3.2
1	C	175	LEU	3.2
3	D	411	GLU	3.2
1	C	290	THR	3.1
1	C	179	TYR	3.0
3	D	360	PHE	2.9
3	D	414	LYS	2.8
3	D	331	LEU	2.8
3	D	377	ILE	2.7
3	D	361	HIS	2.7
2	B	284	ASP	2.7
3	D	419	HIS	2.7
1	C	216	PHE	2.7
1	C	284	PRO	2.7
1	C	231	THR	2.6
3	D	318	TYR	2.6
1	C	239	SER	2.6
1	C	282	ALA	2.6
1	C	233	MET	2.6
3	D	176	PRO	2.6
3	D	376	LEU	2.5
3	D	395	HIS	2.5
1	C	273	LYS	2.5
3	D	400	LYS	2.5
1	C	162	GLU	2.4
1	C	286	PHE	2.4
3	D	224	GLU	2.4
1	C	256	ASP	2.3
3	D	179	HIS	2.3
3	D	426	PRO	2.3
1	C	102	PRO	2.3
1	C	255	LEU	2.2
2	B	431	ASN	2.2
1	C	235	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	285	PHE	2.2
1	C	268	HIS	2.2
3	D	427	PRO	2.2
3	D	384	LEU	2.2
2	B	283	ASP	2.1
3	D	280	TYR	2.1
1	A	15	TYR	2.1
1	A	41	THR	2.1
1	C	247	ASP	2.1
1	C	289	VAL	2.0
1	C	15	TYR	2.0
1	C	166	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPO	A	160	11/12	0.12	-0.78	20,30,33,33	0
1	TPO	C	160	11/12	0.14	-1.01	50,58,69,71	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	SGM	D	1429	6/6	0.27	8.89	69,70,73,85	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	SGM	B	1433	6/6	0.20	2.71	50,58,59,61	0
4	T7Z	C	1295	33/33	0.17	0.49	37,65,103,105	0
4	T7Z	A	1297	33/33	0.15	-0.07	33,43,105,106	0

6.5 Other polymers ⓘ

There are no such residues in this entry.