



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 02:54 AM GMT

PDB ID : 2BO2
Title : EGF DOMAINS 1,2,5 OF HUMAN EMR2, A 7-TM IMMUNE SYSTEM MOLECULE, IN COMPLEX WITH CALCIUM.
Authors : Abbott, R.J.M.; Spendlove, I.; Roversi, P.; Teriete, P.; Knott, V.; Handford, P.A.; McDonnell, J.M.; Lea, S.M.
Deposited on : 2005-04-07
Resolution : 2.60 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1718 (2.60-2.60)
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)
RSRZ outliers	66119	1718 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	143	
1	B	143	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	CAC	A	200	X	X

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 2125 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called EGF-LIKE MODULE CONTAINING MUCIN-LIKE HORMONE RECEPTOR-LIKE 2 PRECURSOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	140	Total	C	N	O	S	0	0	0
			1050	627	182	222	19			
1	B	138	Total	C	N	O	S	0	0	0
			1034	616	179	220	19			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Ca	0	0
			2	2		
2	A	2	Total	Ca	0	0
			2	2		

- Molecule 3 is CACODYLATE ION (three-letter code: CAC) (formula: C₂H₆AsO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	As	C	O	0	0
			5	1	2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	24	Total	O	0	0
			24	24		
4	B	8	Total	O	0	0
			8	8		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: EGF-LIKE MODULE CONTAINING MUCIN-LIKE HORMONE RECEPTOR-LIKE 2 PRECURSOR

Chain A: 



- Molecule 1: EGF-LIKE MODULE CONTAINING MUCIN-LIKE HORMONE RECEPTOR-LIKE 2 PRECURSOR

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.84Å 61.38Å 35.58Å 90.00° 84.10° 90.00°	Depositor
Resolution (Å)	29.54 – 2.60 29.54 – 2.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (29.54-2.60) 99.2 (29.54-2.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 2.61Å)	Xtriage
Refinement program	TNT 5F.0.1	Depositor
R, R_{free}	0.238 , 0.260 0.253 , 0.292	Depositor DCC
R_{free} test set	516 reflections (5.06%)	DCC
Wilson B-factor (Å ²)	43.9	Xtriage
Anisotropy	0.658	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 32.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 10714 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	2125	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/1073	0.45	0/1456
1	B	0.30	0/1056	0.45	0/1432
All	All	0.28	0/2129	0.45	0/2888

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1050	0	928	38	0
1	B	1034	0	905	53	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	5	0	0	0	0
4	A	24	0	0	1	0
4	B	8	0	0	0	0
All	All	2125	0	1833	86	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including

hydrogens) of the entry. The overall clashscore for this entry is 22.

All (86) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:36:THR:HG23	1:B:38:MET:H	1.30	0.94
1:A:36:THR:HG23	1:A:38:MET:H	1.33	0.94
1:B:130:ARG:O	1:B:133:ILE:HG12	1.74	0.86
1:A:32:GLU:HG2	1:A:33:ILE:HG12	1.60	0.83
1:A:24:ASN:HB3	1:A:25:PRO:HD2	1.62	0.81
1:B:130:ARG:HB2	1:B:143:GLU:HG2	1.61	0.79
1:A:48:ALA:O	1:A:50:LEU:HD22	1.83	0.78
1:B:54:SER:HB3	1:B:55:CYS:HA	1.67	0.77
1:B:130:ARG:HG3	1:B:143:GLU:CD	2.06	0.76
1:B:48:ALA:HB3	1:B:51:SER:OG	1.87	0.73
1:A:124:ARG:HG3	1:A:124:ARG:HH21	1.56	0.70
1:B:36:THR:HG23	1:B:38:MET:N	2.05	0.70
1:B:119:TYR:CZ	1:B:138:LYS:HD3	2.28	0.68
1:A:124:ARG:O	1:A:127:TRP:HB2	1.93	0.68
1:B:57:LYS:HD3	1:B:58:PHE:CE1	2.29	0.67
1:A:122:ARG:NE	1:A:136:ASN:OD1	2.29	0.66
1:B:132:GLY:O	1:B:134:PRO:HD3	1.96	0.66
1:A:38:MET:HB3	1:B:131:HIS:ND1	2.11	0.65
1:A:52:LYS:HG2	1:A:53:VAL:N	2.10	0.65
1:A:53:VAL:HG13	1:A:54:SER:N	2.13	0.64
1:B:95:ASP:OD1	1:B:117:GLY:N	2.31	0.64
1:B:22:ARG:NH1	1:B:33:ILE:HD11	2.12	0.63
1:A:137:GLN:NE2	1:A:139:ASP:OD1	2.32	0.63
1:A:127:TRP:O	1:A:128:LYS:HE3	1.99	0.62
1:B:87:ASN:ND2	1:B:89:SER:OG	2.32	0.62
1:B:133:ILE:HD13	1:B:133:ILE:N	2.13	0.62
1:B:6:ALA:N	1:B:17:ASN:O	2.29	0.62
1:A:119:TYR:CZ	1:A:138:LYS:HD3	2.36	0.61
1:B:131:HIS:H	1:B:131:HIS:CD2	2.17	0.61
1:A:138:LYS:HB2	1:B:50:LEU:CD2	2.33	0.59
1:B:99:CYS:HA	1:B:104:HIS:HD2	1.67	0.59
1:A:124:ARG:NH2	1:A:124:ARG:HG3	2.15	0.58
1:B:54:SER:CB	1:B:55:CYS:HA	2.32	0.58
1:B:59:SER:HG	1:B:85:PHE:HZ	1.52	0.58
1:A:36:THR:HG23	1:A:38:MET:N	2.12	0.57
1:B:110:THR:OG1	1:B:121:CYS:HB3	2.06	0.56
1:B:99:CYS:HA	1:B:104:HIS:CD2	2.41	0.55
1:A:131:HIS:CE1	1:B:32:GLU:HA	2.41	0.55
1:B:36:THR:HG22	1:B:39:GLU:HG3	1.88	0.55
1:B:36:THR:HG21	1:B:38:MET:HE2	1.89	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:32:GLU:HG2	1:B:33:ILE:HG12	1.90	0.54
1:A:24:ASN:HB3	1:A:25:PRO:CD	2.36	0.54
1:A:55:CYS:HB2	1:A:59:SER:OG	2.09	0.53
1:A:138:LYS:HB2	1:B:50:LEU:HD23	1.90	0.52
1:A:17:ASN:ND2	4:A:2006:HOH:O	2.32	0.52
1:B:11:GLN:O	1:B:12:ASP:HB2	2.09	0.52
1:B:58:PHE:CD2	1:B:76:TYR:CE2	2.98	0.52
1:B:54:SER:HA	1:B:55:CYS:C	2.31	0.51
1:B:124:ARG:NH1	1:B:127:TRP:CZ2	2.79	0.51
1:B:96:VAL:O	1:B:98:GLU:OE2	2.29	0.49
1:A:130:ARG:HG2	1:A:130:ARG:HH11	1.78	0.49
1:B:118:SER:OG	1:B:138:LYS:HE2	2.13	0.48
1:B:127:TRP:N	1:B:127:TRP:CD1	2.80	0.48
1:B:22:ARG:HH11	1:B:33:ILE:HD11	1.77	0.48
1:A:53:VAL:CG1	1:A:54:SER:N	2.77	0.48
1:B:130:ARG:HB2	1:B:143:GLU:CG	2.38	0.47
1:B:130:ARG:HG3	1:B:143:GLU:OE2	2.13	0.47
1:A:130:ARG:NH1	1:A:130:ARG:HG2	2.28	0.47
1:B:16:VAL:HG22	1:B:20:ALA:O	2.15	0.47
1:B:76:TYR:HA	1:B:94:GLN:O	2.15	0.47
1:B:80:SER:C	1:B:82:ALA:H	2.17	0.47
1:B:106:CYS:HB3	1:B:110:THR:OG1	2.15	0.46
1:A:124:ARG:H	1:A:124:ARG:HG2	1.39	0.46
1:A:16:VAL:HG23	1:A:22:ARG:HB2	1.98	0.46
1:A:127:TRP:C	1:A:128:LYS:HE3	2.36	0.45
1:B:85:PHE:HB2	1:B:90:GLU:HB2	1.98	0.45
1:B:36:THR:CG2	1:B:38:MET:HE2	2.47	0.45
1:A:106:CYS:HB3	1:A:110:THR:OG1	2.16	0.45
1:A:87:ASN:HD22	1:A:89:SER:H	1.64	0.44
1:B:48:ALA:HB3	1:B:51:SER:HG	1.81	0.44
1:A:131:HIS:CD2	1:B:32:GLU:HB2	2.53	0.44
1:A:87:ASN:HD22	1:A:89:SER:N	2.16	0.44
1:A:7:ARG:HG2	1:A:7:ARG:H	1.58	0.44
1:A:48:ALA:O	1:A:50:LEU:N	2.51	0.43
1:B:57:LYS:HD3	1:B:58:PHE:HE1	1.79	0.43
1:B:16:VAL:HG23	1:B:17:ASN:ND2	2.33	0.43
1:B:119:TYR:CE2	1:B:138:LYS:HD3	2.53	0.43
1:A:11:GLN:O	1:A:12:ASP:HB2	2.18	0.43
1:A:111:VAL:O	1:A:111:VAL:HG13	2.19	0.42
1:B:97:ASP:HA	1:B:114:ASN:ND2	2.35	0.42
1:A:56:GLY:HA3	1:A:91:ASN:HD22	1.84	0.42
1:B:22:ARG:NH1	1:B:33:ILE:CD1	2.80	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:119:TYR:CE2	1:A:138:LYS:HD3	2.54	0.42
1:A:109:SER:O	1:A:124:ARG:HD3	2.20	0.41
1:B:8:TRP:HZ2	1:B:19:THR:O	2.04	0.41
1:B:27:PHE:CE1	1:B:66:GLY:HA2	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/143 (96%)	126 (91%)	10 (7%)	2 (1%)	16	32
1	B	134/143 (94%)	116 (87%)	18 (13%)	0	100	100
All	All	272/286 (95%)	242 (89%)	28 (10%)	2 (1%)	30	58

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	49	THR
1	A	130	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	125/128 (98%)	103 (82%)	22 (18%)	3	4
1	B	123/128 (96%)	104 (85%)	19 (15%)	4	6
All	All	248/256 (97%)	207 (84%)	41 (16%)	3	5

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ARG
1	A	16	VAL
1	A	22	ARG
1	A	31	SER
1	A	36	THR
1	A	43	ASP
1	A	50	LEU
1	A	52	LYS
1	A	53	VAL
1	A	54	SER
1	A	83	LYS
1	A	87	ASN
1	A	94	GLN
1	A	95	ASP
1	A	108	SER
1	A	116	VAL
1	A	124	ARG
1	A	128	LYS
1	A	130	ARG
1	A	133	ILE
1	A	137	GLN
1	A	143	GLU
1	B	7	ARG
1	B	16	VAL
1	B	19	THR
1	B	28	SER
1	B	36	THR
1	B	43	ASP
1	B	49	THR
1	B	50	LEU
1	B	57	LYS
1	B	58	PHE
1	B	83	LYS
1	B	87	ASN
1	B	108	SER
1	B	110	THR
1	B	111	VAL
1	B	128	LYS
1	B	131	HIS
1	B	133	ILE
1	B	143	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	ASN
1	A	131	HIS
1	A	137	GLN
1	B	11	GLN
1	B	87	ASN
1	B	137	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CAC	A	200	-	4,4,4	8.22	4 (100%)	6,6,6	5.82	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CAC	A	200	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	200	CAC	O2-AS	14.51	1.84	1.68
3	A	200	CAC	AS-C2	5.26	1.99	1.90
3	A	200	CAC	AS-C1	5.19	1.99	1.90
3	A	200	CAC	O1-AS	-2.29	1.58	1.66

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	200	CAC	O2-AS-O1	-14.23	109.47	112.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	140/143 (97%)	0.21	6 (4%) 34 30	17, 30, 58, 71	0
1	B	138/143 (96%)	0.57	6 (4%) 34 30	44, 58, 76, 88	0
All	All	278/286 (97%)	0.39	12 (4%) 31 30	17, 51, 71, 88	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	53	VAL	5.4
1	A	133	ILE	5.3
1	A	132	GLY	4.9
1	B	82	ALA	4.0
1	B	58	PHE	3.7
1	B	79	VAL	3.5
1	A	51	SER	2.8
1	B	83	LYS	2.6
1	B	7	ARG	2.4
1	B	130	ARG	2.2
1	A	130	ARG	2.1
1	A	141	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	CAC	A	200	5/5	0.47	3.88	90,90,91,91	5
2	CA	A	156	1/1	0.14	-0.72	27,27,27,27	0
2	CA	A	155	1/1	0.14	-0.88	22,22,22,22	0
2	CA	B	155	1/1	0.12	-1.15	54,54,54,54	0
2	CA	B	156	1/1	0.11	-2.01	64,64,64,64	0

6.5 Other polymers

There are no such residues in this entry.