



wwPDB X-ray Structure Validation Summary Report (i)

Feb 27, 2014 – 01:56 PM GMT

PDB ID : 3BSB
Title : Crystal Structure of Human Pumilio1 in Complex with CyclinB reverse RNA
Authors : Gupta, Y.K.; Nair, D.T.; Wharton, R.P.; Aggarwal, A.K.
Deposited on : 2007-12-23
Resolution : 2.80 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1799 (2.80-2.80)
Clashscore	79885	2295 (2.80-2.80)
Ramachandran outliers	78287	2252 (2.80-2.80)
Sidechain outliers	78261	2254 (2.80-2.80)
RSRZ outliers	66119	1802 (2.80-2.80)
RNA backbone	1838	1076 (3.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	C	9	
2	A	343	
2	B	343	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5820 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(*UP*UP*UP*AP*AP*UP*GP*UP*U)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	9	Total	C	N	O	P	0	0	0
			184	84	27	65	8			

- Molecule 2 is a protein called Pumilio homolog 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	340	Total	C	N	O	S	0	0	0
			2742	1741	490	494	17			
2	B	341	Total	C	N	O	S	0	0	0
			2749	1743	490	499	17			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	48	Total	O	0	0
			48	48		
3	B	83	Total	O	0	0
			83	83		
3	C	14	Total	O	0	0
			14	14		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-R(*UP*UP*UP*AP*AP*UP*GP*UP*U)-3'

Chain C: 




- Molecule 2: Pumilio homolog 1

Chain A: 



- Molecule 2: Pumilio homolog 1

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	35.87Å 64.29Å 321.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.87 – 2.80 25.30 – 2.80	Depositor EDS
% Data completeness (in resolution range)	86.3 (25.87-2.80) 89.6 (25.30-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.08 (at 2.80Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.244 , 0.268 0.249 , 0.272	Depositor DCC
R_{free} test set	1329 reflections (8.32%)	DCC
Wilson B-factor (Å ²)	39.4	Xtriage
Anisotropy	0.934	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 19.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 17894 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5820	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	1.33	1/204 (0.5%)	2.65	24/315 (7.6%)
2	A	0.42	0/2796	0.61	0/3776
2	B	0.59	0/2803	0.81	9/3786 (0.2%)
All	All	0.56	1/5803 (0.0%)	0.88	33/7877 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	3	U	C3'-O3'	6.25	1.50	1.42

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	8	U	N1-C1'-C2'	-13.22	96.81	114.00
1	C	3	U	P-O3'-C3'	10.59	132.41	119.70
1	C	2	U	O4'-C4'-C3'	-10.28	93.72	104.00
1	C	7	G	N9-C1'-C2'	-9.98	101.02	112.00
2	B	1108	PRO	CA-N-CD	-9.26	98.54	111.50

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	1	U	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	C	4	A	Sidechain
1	C	6	U	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	184	0	95	27	0
2	A	2742	0	2737	132	0
2	B	2749	0	2735	203	0
3	A	48	0	0	20	0
3	B	83	0	0	64	0
3	C	14	0	0	10	0
All	All	5820	0	5567	343	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 31.

The worst 5 of 343 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:A:829:ARG:NE	2:A:837:ARG:HH12	1.27	1.28
2:B:982:GLN:HB3	3:B:141:HOH:O	1.26	1.25
2:A:1138:LYS:HG2	3:A:166:HOH:O	1.37	1.22
2:A:829:ARG:NE	2:A:837:ARG:NH1	1.86	1.21
2:B:1098:ILE:HG12	3:B:145:HOH:O	1.43	1.14

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	338/343 (98%)	315 (93%)	18 (5%)	5 (2%)	15	46
2	B	339/343 (99%)	313 (92%)	21 (6%)	5 (2%)	15	46
All	All	677/686 (99%)	628 (93%)	39 (6%)	10 (2%)	15	46

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	922	ARG
2	A	1134	PRO
2	B	1107	GLY
2	B	1134	PRO
2	B	1166	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	298/306 (97%)	286 (96%)	12 (4%)	42	79
2	B	299/306 (98%)	270 (90%)	29 (10%)	12	32
All	All	597/612 (98%)	556 (93%)	41 (7%)	22	53

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	912	GLU
2	B	984	GLN
2	B	1163	LYS
2	B	940	LYS
2	B	957	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	860	GLN
2	B	891	GLN
2	B	1109	HIS
2	B	883	ASN
2	A	949	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	C	9/9 (100%)	8 (88%)	4 (44%)

5 of 8 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	C	2	U
1	C	3	U
1	C	4	A
1	C	5	A
1	C	6	U

All (4) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	C	1	U
1	C	2	U
1	C	4	A
1	C	6	U

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	9/9 (100%)	1.67	3 (33%) 1 1	58, 70, 86, 90	0
2	A	340/343 (99%)	0.22	9 (2%) 53 54	25, 47, 73, 87	0
2	B	341/343 (99%)	0.02	6 (1%) 65 66	19, 37, 69, 87	0
All	All	690/695 (99%)	0.14	18 (2%) 53 54	19, 42, 73, 90	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	828	GLY	4.3
2	A	1156	TYR	3.8
2	B	840	ARG	3.8
1	C	7	G	3.2
2	A	1154	TYR	2.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.