



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 01:01 PM GMT

PDB ID : 2BT3
Title : AGAO IN COMPLEX WITH RUTHENIUM-C4-WIRE AT 1.73
ANGSTROMS
Authors : Langley, D.B.; Duff, A.P.; Freeman, H.C.; Guss, J.M.; Juda, G.A.; Dooley,
D.M.; Contakes, S.M.; Halpern-Manners, N.W.; Dunn, A.R.; Gray, H.B.
Deposited on : 2005-05-26
Resolution : 1.73 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1657 (1.76-1.72)
Clashscore	79885	1881 (1.76-1.72)
Ramachandran outliers	78287	1859 (1.76-1.72)
Sidechain outliers	78261	1859 (1.76-1.72)
RSRZ outliers	66119	1658 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	646	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	NA	A	1630	-	X
4	R4A	A	1631[A]	-	X
4	R4A	A	1631[B]	-	X
5	SO4	A	1633	-	X
5	SO4	A	1634	-	X
6	GOL	A	1636	-	X
6	GOL	A	1637	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
6	GOL	A	1638	-	X
6	GOL	A	1639	-	X
6	GOL	A	1640	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10157 atoms, of which 4604 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHENYLETHYLAMINE OXIDASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	620	Total	C	H	N	O	S	0	8	0
			9397	3093	4502	856	935	11			


- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cu	0	0
			1	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is BIS[1H,1'H-2,2'-BIPYRIDINATO(2-)-KAPPA 2 N 1 ,N 1']{3-[4-(1,10-DIHYDRO-1,10-PHENANTHROLIN-4-YL-KAPPA 2 N 1 ,N 10)BUTOXY]-N,N-DIMETHYLANILINATO(2-)}RUTHENIUM (three-letter code: R4A) (formula: C₄₄H₆₁N₇ORu).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	Ru		
4	A	1	169	80	72	13	2	2	0	1

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
5	A	1	5	4	1	0	0
5	A	1	5	4	1	0	0
5	A	1	5	4	1	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			11	3	5	3		
6	A	1	Total	C	H	O	0	0
			11	3	5	3		
6	A	1	Total	C	H	O	0	0
			11	3	5	3		
6	A	1	Total	C	H	O	0	0
			11	3	5	3		
6	A	1	Total	C	H	O	0	0
			11	3	5	3		
6	A	1	Total	C	H	O	0	0
			11	3	5	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	508	Total	O	0	0
			508	508		

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	158.06Å 62.91Å 92.10Å 90.00° 112.11° 90.00°	Depositor
Resolution (Å)	15.02 – 1.73 15.02 – 1.73	Depositor EDS
% Data completeness (in resolution range)	94.5 (15.02-1.73) 94.5 (15.02-1.73)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 1.73Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.154 , 0.171 0.164 , 0.178	Depositor DCC
R_{free} test set	4106 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	20.0	Xtriage
Anisotropy	0.668	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.47 , 44.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 82054 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10157	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, R4A, NA, SO4, TPQ, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	3/5037 (0.1%)	0.78	7/6858 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	226	GLU	CD-OE2	22.33	1.50	1.25
1	A	226	GLU	CD-OE1	10.64	1.37	1.25
1	A	226	GLU	C-N	7.70	1.47	1.33

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	316	ASP	CB-CG-OD2	6.05	123.75	118.30
1	A	357	ASP	CB-CG-OD2	5.54	123.28	118.30
1	A	66	ASP	CB-CG-OD2	5.36	123.13	118.30
1	A	246	ASP	CB-CG-OD2	5.28	123.05	118.30
1	A	115	ASP	CB-CG-OD2	5.07	122.86	118.30
1	A	266	ASP	CB-CG-OD2	5.05	122.84	118.30
1	A	612	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the

chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4895	4502	225	20	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	97	72	30	10	0
5	A	15	0	0	0	0
6	A	36	30	18	0	0
7	A	508	0	0	0	0
All	All	5553	4604	273	22	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (22) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:294[A]:GLN:HG3	1:A:296:TYR:CZ	2.29	0.68
1:A:307:TYR:CE1	4:A:1631[B]:R4A:CAM	2.78	0.66
1:A:102:GLU:HA	1:A:105:PHE:CE2	2.34	0.62
1:A:599:TRP:CD2	1:A:600:PRO:HA	2.34	0.61
1:A:307:TYR:CE1	4:A:1631[B]:R4A:HAM	2.39	0.58
1:A:102:GLU:OE1	4:A:1631[B]:R4A:HBI1	2.09	0.52
1:A:102:GLU:OE2	4:A:1631[A]:R4A:HCC1	2.11	0.50
1:A:294[A]:GLN:HG3	1:A:296:TYR:CE2	2.48	0.48
1:A:280[B]:MET:HG3	1:A:435:PHE:CE2	2.50	0.47
4:A:1631[B]:R4A:NAD	4:A:1631[B]:R4A:HBB	2.30	0.46
1:A:105:PHE:CE2	1:A:307:TYR:OH	2.68	0.46
1:A:105:PHE:HE2	1:A:307:TYR:OH	1.99	0.46
1:A:280[B]:MET:HG3	1:A:435:PHE:CD2	2.52	0.45
1:A:595:ARG:HG3	1:A:597:GLU:HB2	1.98	0.45
1:A:105:PHE:CD2	4:A:1631[A]:R4A:HL21	2.53	0.44
1:A:592:HIS:HE1	1:A:602[A]:MET:SD	2.41	0.44
1:A:105:PHE:CE2	4:A:1631[A]:R4A:HL21	2.53	0.44
1:A:294[A]:GLN:CG	1:A:296:TYR:CE2	3.02	0.42
1:A:307:TYR:CZ	4:A:1631[B]:R4A:HAM	2.55	0.41
4:A:1631[B]:R4A:CAC	4:A:1631[B]:R4A:HBB	2.51	0.41
1:A:282:VAL:HA	1:A:432:GLN:O	2.21	0.41
1:A:102:GLU:OE1	4:A:1631[B]:R4A:HBH1	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	625/646 (97%)	606 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	521/537 (97%)	516 (99%)	5 (1%)	85	73

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	267	ARG
1	A	336	ARG
1	A	369	ARG
1	A	376	PHE
1	A	444	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	309	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPQ	A	382	1	14,14,15	4.57	4 (28%)	17,19,21	1.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPQ	A	382	1	-	0/4/22/24	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	382	TPQ	O-C	16.22	1.22	1.11
1	A	382	TPQ	C3-C4	3.24	1.40	1.35
1	A	382	TPQ	O4-C4	-2.46	1.27	1.34
1	A	382	TPQ	CA-C	2.24	1.52	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	R4A	A	1631[A]	-	53,53,63	7.79	25 (47%)	58,87,101	3.35	38 (65%)
4	R4A	A	1631[B]	-	53,53,63	7.55	24 (45%)	58,87,101	3.34	36 (62%)
5	SO4	A	1632	-	4,4,4	0.23	0	6,6,6	0.54	0
5	SO4	A	1633	-	4,4,4	0.20	0	6,6,6	0.18	0
5	SO4	A	1634	-	4,4,4	0.13	0	6,6,6	0.09	0
6	GOL	A	1635	-	5,5,5	0.23	0	5,5,5	0.38	0
6	GOL	A	1636	-	5,5,5	0.29	0	5,5,5	0.28	0
6	GOL	A	1637	-	5,5,5	0.28	0	5,5,5	0.43	0
6	GOL	A	1638	-	5,5,5	0.30	0	5,5,5	0.25	0
6	GOL	A	1639	-	5,5,5	0.40	0	5,5,5	0.42	0
6	GOL	A	1640	-	5,5,5	0.21	0	5,5,5	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	R4A	A	1631[A]	-	5/5/18/19	0/5/143/150	0/0/10/11
4	R4A	A	1631[B]	-	5/5/18/19	0/5/143/150	0/0/10/11
5	SO4	A	1632	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1633	-	-	0/0/0/0	0/0/0/0
5	SO4	A	1634	-	-	0/0/0/0	0/0/0/0
6	GOL	A	1635	-	-	0/4/4/4	0/0/0/0
6	GOL	A	1636	-	-	0/4/4/4	0/0/0/0
6	GOL	A	1637	-	-	0/4/4/4	0/0/0/0
6	GOL	A	1638	-	-	0/4/4/4	0/0/0/0
6	GOL	A	1639	-	-	0/4/4/4	0/0/0/0
6	GOL	A	1640	-	-	0/4/4/4	0/0/0/0

All (49) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1631[A]	R4A	CCB-NCA	-31.25	1.27	1.45
4	A	1631[A]	R4A	CBB-NBA	-29.54	1.28	1.45
4	A	1631[B]	R4A	CCB-NCA	-29.40	1.28	1.45
4	A	1631[B]	R4A	CBB-NBA	-28.56	1.28	1.45
4	A	1631[A]	R4A	CCK-NCL	-19.31	1.34	1.45
4	A	1631[B]	R4A	CCK-NCL	-18.75	1.34	1.45
4	A	1631[A]	R4A	CBK-NBL	-18.68	1.34	1.45
4	A	1631[B]	R4A	CBK-NBL	-17.36	1.35	1.45
4	A	1631[B]	R4A	CAN-NAA	-15.49	1.34	1.47
4	A	1631[A]	R4A	CAN-NAA	-14.39	1.35	1.47
4	A	1631[A]	R4A	CCK-CCJ	-7.76	1.40	1.51
4	A	1631[B]	R4A	CCK-CCJ	-7.63	1.40	1.51
4	A	1631[B]	R4A	CBD-CBE	-7.13	1.32	1.53
4	A	1631[A]	R4A	CBD-CBE	-7.01	1.33	1.53
4	A	1631[B]	R4A	CCD-CCE	-7.00	1.33	1.53
4	A	1631[A]	R4A	CCD-CCE	-6.95	1.33	1.53
4	A	1631[A]	R4A	CAI-CAJ	5.58	1.41	1.33
4	A	1631[A]	R4A	CBH-CBG	-5.55	1.39	1.52
4	A	1631[B]	R4A	CBH-CBG	-5.47	1.39	1.52
4	A	1631[B]	R4A	CAI-CAJ	5.42	1.41	1.33
4	A	1631[A]	R4A	CCG-CCH	-5.38	1.39	1.50
4	A	1631[B]	R4A	CCG-CCH	-5.19	1.39	1.50
4	A	1631[B]	R4A	CAK-CAL	-4.77	1.48	1.54
4	A	1631[B]	R4A	CBI-CBH	-4.70	1.39	1.53
4	A	1631[A]	R4A	CBI-CBH	-4.70	1.39	1.53
4	A	1631[A]	R4A	CCC-CCB	-4.49	1.33	1.51
4	A	1631[B]	R4A	CBC-CBB	-4.38	1.34	1.51
4	A	1631[A]	R4A	CBC-CBB	-4.37	1.34	1.51
4	A	1631[B]	R4A	CCC-CCB	-4.30	1.34	1.51
4	A	1631[B]	R4A	CCG-CCF	-3.84	1.49	1.55
4	A	1631[A]	R4A	CAK-CAL	-3.71	1.50	1.54
4	A	1631[B]	R4A	CAK-CAB	-3.70	1.46	1.52
4	A	1631[A]	R4A	CAK-CAB	-3.37	1.47	1.52
4	A	1631[A]	R4A	CCG-CCF	-3.35	1.50	1.55
4	A	1631[B]	R4A	CAK-CAJ	-3.34	1.40	1.52
4	A	1631[A]	R4A	CAK-CAJ	-3.25	1.41	1.52
4	A	1631[B]	R4A	CAM-CAL	3.20	1.40	1.33
4	A	1631[A]	R4A	CAM-CAL	3.16	1.40	1.33
4	A	1631[B]	R4A	CBJ-CBK	-2.92	1.39	1.51
4	A	1631[B]	R4A	CBI-CBJ	-2.77	1.39	1.51
4	A	1631[A]	R4A	CBJ-CBK	-2.73	1.40	1.51
4	A	1631[A]	R4A	CCH-CCI	2.68	1.39	1.32
4	A	1631[A]	R4A	CAC-CAH	2.66	1.46	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1631[B]	R4A	CCH-CCI	2.61	1.39	1.32
4	A	1631[B]	R4A	CAC-CAH	2.59	1.46	1.41
4	A	1631[A]	R4A	CBI-CBJ	-2.57	1.40	1.51
4	A	1631[B]	R4A	CCJ-CCI	-2.57	1.39	1.48
4	A	1631[A]	R4A	CCJ-CCI	-2.53	1.40	1.48
4	A	1631[A]	R4A	CAN-CAM	-2.04	1.40	1.48

All (74) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1631[A]	R4A	CCG-CCF-NCA	8.08	116.41	102.28
4	A	1631[B]	R4A	CCG-CCF-NCA	7.95	116.18	102.28
4	A	1631[B]	R4A	CCJ-CCK-NCL	7.81	119.84	110.37
4	A	1631[A]	R4A	CCJ-CCK-NCL	7.50	119.47	110.37
4	A	1631[B]	R4A	CBC-CBB-NBA	6.63	123.05	110.23
4	A	1631[A]	R4A	CCK-NCL-CCG	6.30	121.72	111.86
4	A	1631[B]	R4A	CCC-CCB-NCA	6.25	122.31	110.23
4	A	1631[A]	R4A	CBC-CBB-NBA	6.15	122.12	110.23
4	A	1631[A]	R4A	CCC-CCB-NCA	6.03	121.89	110.23
4	A	1631[B]	R4A	CCK-NCL-CCG	5.61	120.64	111.86
4	A	1631[B]	R4A	CBJ-CBK-NBL	5.25	120.38	110.23
4	A	1631[B]	R4A	CCF-CCG-NCL	5.12	113.35	102.15
4	A	1631[A]	R4A	CCF-CCG-NCL	5.09	113.29	102.15
4	A	1631[A]	R4A	CBJ-CBK-NBL	4.88	119.67	110.23
4	A	1631[B]	R4A	CCH-CCG-NCL	4.88	120.72	110.65
4	A	1631[A]	R4A	CAJ-CAK-CAL	4.82	127.99	112.87
4	A	1631[B]	R4A	CAJ-CAK-CAL	4.56	127.19	112.87
4	A	1631[A]	R4A	CCH-CCG-NCL	4.51	119.97	110.65
4	A	1631[A]	R4A	CCE-CCF-CCG	4.51	124.29	116.61
4	A	1631[B]	R4A	CAJ-CAK-CAB	4.42	118.49	108.93
4	A	1631[B]	R4A	CBI-CBJ-CBK	4.28	120.33	111.26
4	A	1631[A]	R4A	CBI-CBJ-CBK	4.20	120.15	111.26
4	A	1631[A]	R4A	CAJ-CAK-CAB	4.08	117.75	108.93
4	A	1631[B]	R4A	CCE-CCF-CCG	4.03	123.46	116.61
4	A	1631[A]	R4A	CAF-CAE-NAD	-4.01	119.95	122.72
4	A	1631[B]	R4A	CCD-CCC-CCB	3.99	119.72	111.26
4	A	1631[A]	R4A	CBJ-CBI-CBH	3.98	120.28	111.45
4	A	1631[B]	R4A	CBD-CBC-CBB	3.95	119.62	111.26
4	A	1631[A]	R4A	CCD-CCC-CCB	3.93	119.58	111.26
4	A	1631[A]	R4A	CBD-CBC-CBB	3.80	119.30	111.26
4	A	1631[B]	R4A	CBJ-CBI-CBH	3.80	119.88	111.45
4	A	1631[B]	R4A	CAL-CAK-CAB	3.58	114.32	108.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1631[A]	R4A	CCF-CCG-CCH	3.56	126.74	115.10
4	A	1631[A]	R4A	CAL-CAK-CAB	3.55	114.27	108.54
4	A	1631[A]	R4A	CCB-NCA-CCF	3.48	124.05	112.75
4	A	1631[B]	R4A	CBI-CBH-CBG	3.48	118.98	110.53
4	A	1631[B]	R4A	CAF-CAE-NAD	-3.47	120.32	122.72
4	A	1631[B]	R4A	CCF-CCG-CCH	3.31	125.92	115.10
4	A	1631[A]	R4A	CBB-NBA-CBF	3.27	123.36	112.75
4	A	1631[B]	R4A	CBD-CBE-CBF	3.23	118.39	110.53
4	A	1631[A]	R4A	CBI-CBH-CBG	3.20	118.30	110.53
4	A	1631[A]	R4A	CCD-CCE-CCF	3.13	118.13	110.53
4	A	1631[B]	R4A	CCB-NCA-CCF	3.02	122.56	112.75
4	A	1631[A]	R4A	CBD-CBE-CBF	3.01	117.85	110.53
4	A	1631[B]	R4A	CCD-CCE-CCF	3.00	117.83	110.53
4	A	1631[B]	R4A	CAC-CAB-CAK	-2.92	119.18	125.18
4	A	1631[A]	R4A	CAK-CAL-CAM	2.85	118.12	112.02
4	A	1631[A]	R4A	CBC-CBD-CBE	2.76	117.57	111.45
4	A	1631[B]	R4A	CBB-NBA-CBF	2.75	121.69	112.75
4	A	1631[A]	R4A	CAG-CAH-CAC	-2.73	115.73	118.71
4	A	1631[B]	R4A	CCK-CCJ-CCI	2.70	120.48	112.02
4	A	1631[A]	R4A	CBH-CBG-NBL	2.59	120.58	111.07
4	A	1631[A]	R4A	CCK-CCJ-CCI	2.59	120.11	112.02
4	A	1631[B]	R4A	CCC-CCD-CCE	2.57	117.16	111.45
4	A	1631[A]	R4A	CAC-CAB-CAK	-2.55	119.94	125.18
4	A	1631[A]	R4A	CBK-NBL-CBG	2.54	121.01	112.75
4	A	1631[B]	R4A	CAG-CAH-CAC	-2.53	115.94	118.71
4	A	1631[B]	R4A	CCE-CCF-NCA	2.53	120.36	111.07
4	A	1631[B]	R4A	CBE-CBF-NBA	2.52	120.34	111.07
4	A	1631[A]	R4A	CCC-CCD-CCE	2.51	117.02	111.45
4	A	1631[B]	R4A	CCJ-CCI-CCH	-2.50	119.81	123.67
4	A	1631[B]	R4A	CBH-CBG-NBL	2.50	120.24	111.07
4	A	1631[B]	R4A	CBC-CBD-CBE	2.46	116.91	111.45
4	A	1631[B]	R4A	CAK-CAL-CAM	2.46	117.28	112.02
4	A	1631[A]	R4A	CBE-CBF-NBA	2.37	119.79	111.07
4	A	1631[B]	R4A	CAC-CAH-CAI	-2.30	118.07	123.34
4	A	1631[B]	R4A	CBK-NBL-CBG	2.29	120.18	112.75
4	A	1631[A]	R4A	CCJ-CCI-CCH	-2.28	120.16	123.67
4	A	1631[A]	R4A	CCE-CCF-NCA	2.24	119.30	111.07
4	A	1631[A]	R4A	CAC-CAH-CAI	-2.19	118.32	123.34
4	A	1631[A]	R4A	CL1-CAL-CAM	-2.11	118.19	125.74
4	A	1631[A]	R4A	CAE-CAF-CAG	2.08	122.73	119.44
4	A	1631[A]	R4A	CAM-CAN-NAA	2.06	121.06	111.00
4	A	1631[B]	R4A	OL5-CL4-CL3	-2.03	107.75	114.71

All (10) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	1631[A]	R4A	CAK
4	A	1631[A]	R4A	CBG
4	A	1631[A]	R4A	CCF
4	A	1631[A]	R4A	CBF
4	A	1631[A]	R4A	CCG
4	A	1631[B]	R4A	CAK
4	A	1631[B]	R4A	CBG
4	A	1631[B]	R4A	CCF
4	A	1631[B]	R4A	CBF
4	A	1631[B]	R4A	CCG

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	620/646 (95%)	0.04	36 (5%)	22 27	5, 12, 28, 62	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	53	SER	10.5
1	A	52	GLY	8.2
1	A	54	GLU	6.5
1	A	9	ALA	5.4
1	A	51	ALA	5.0
1	A	628	ASN	4.9
1	A	314	GLY	4.4
1	A	266	ASP	4.3
1	A	315	CYS	4.3
1	A	55	ALA	4.0
1	A	212	PRO	3.4
1	A	225	PRO	3.4
1	A	105	PHE	3.3
1	A	358	LEU	3.1
1	A	313	LEU	3.0
1	A	462	PRO	3.0
1	A	88	VAL	2.9
1	A	68	SER	2.8
1	A	215	THR	2.6
1	A	223	THR	2.5
1	A	336	ARG	2.5
1	A	207	PRO	2.4
1	A	318	LEU	2.3
1	A	208	GLU	2.3
1	A	265	GLY	2.3
1	A	228	PRO	2.3
1	A	50	GLY	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	116	GLU	2.2
1	A	267	ARG	2.2
1	A	463	GLY	2.2
1	A	70	ALA	2.2
1	A	145	ALA	2.1
1	A	459	THR	2.1
1	A	332	PHE	2.1
1	A	414	GLY	2.0
1	A	339	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPQ	A	382	14/15	0.11	1.63	7,14,33,46	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	SO4	A	1634	5/5	0.38	17.35	74,80,84,86	0
6	GOL	A	1639	6/6	0.21	14.07	38,41,43,47	0
6	GOL	A	1637	6/6	0.19	11.15	55,57,62,65	0
5	SO4	A	1633	5/5	0.33	9.66	107,107,108,110	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	GOL	A	1636	6/6	0.17	6.97	29,35,41,43	0
6	GOL	A	1640	6/6	0.16	5.91	28,40,42,50	0
4	R4A	A	1631[A]	44/53	0.32	5.84	22,31,38,41	75
6	GOL	A	1638	6/6	0.18	3.83	27,48,58,58	0
4	R4A	A	1631[B]	44/53	0.32	2.80	40,42,44,46	75
3	NA	A	1630	1/1	0.22	2.53	21,21,21,21	0
6	GOL	A	1635	6/6	0.14	1.68	23,26,28,28	0
5	SO4	A	1632	5/5	0.12	1.31	47,49,52,53	0
2	CU	A	1629	1/1	0.11	1.13	19,19,19,19	0

6.5 Other polymers ⓘ

There are no such residues in this entry.