



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 01:19 AM GMT

PDB ID : 2BZS  
Title : BINDING OF ANTI-CANCER PRODRUG CB1954 TO THE ACTIVATING ENZYME NQO2 REVEALED BY THE CRYSTAL STRUCTURE OF THEIR COMPLEX.  
Authors : Abu Khader, M.M.; Heap, J.T.; De Matteis, C.; Kellam, B.; Doughty, S.W.; Minton, N.; Paoli, M.  
Deposited on : 2005-08-22  
Resolution : 2.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	4888 (2.00-2.00)
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	236	
1	B	236	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	CB1	A	235	-	X

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4091 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NRH DEHYDROGENASE [QUINONE] 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	S	0	0	0
			1824	1174	304	338	8			
1	B	230	Total	C	N	O	S	11	0	0
			1824	1174	304	338	8			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is 5-(AZIRIDIN-1-YL)-2,4-DINITROBENZAMIDE (three-letter code: CB1) (formula: C<sub>9</sub>H<sub>8</sub>N<sub>4</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			18	9	4	5		
4	B	1	Total	C	N	O	0	0
			18	9	4	5		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	167	Total	O	0	0
			167	167		
5	B	132	Total	O	0	0
			132	132		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: NRH DEHYDROGENASE [QUINONE] 2

Chain A: 



#### • Molecule 1: NRH DEHYDROGENASE [QUINONE] 2

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.83Å 106.11Å 61.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 29.60 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (30.00-2.00) 99.4 (29.60-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.41 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.193 , 0.247 0.198 , 0.240	Depositor DCC
$R_{free}$ test set	1813 reflections (5.25%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.6	Xtriage
Anisotropy	0.121	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 40.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 36325 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4091	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CB1, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.83	0/1874	0.80	2/2542 (0.1%)
1	B	0.79	0/1874	0.80	2/2542 (0.1%)
All	All	0.81	0/3748	0.80	4/5084 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	118	ARG	NE-CZ-NH2	-9.39	115.61	120.30
1	B	118	ARG	NE-CZ-NH1	8.22	124.41	120.30
1	A	200	ARG	NE-CZ-NH2	-5.58	117.51	120.30
1	A	127	ASP	CB-CG-OD2	-5.46	113.39	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	229	GLY	Peptide

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1824	0	1779	20	0
1	B	1824	0	1779	39	3
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	53	0	31	3	0
3	B	53	0	31	2	0
4	A	18	0	8	10	0
4	B	18	0	8	5	0
5	A	167	0	0	4	0
5	B	132	0	0	9	0
All	All	4091	0	3636	62	3

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (62) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:216:LYS:HD2	5:B:2121:HOH:O	1.58	1.02
1:B:56:THR:O	1:B:56:THR:HG23	1.61	0.98
4:A:235:CB1:H91	5:A:2166:HOH:O	1.66	0.94
4:A:235:CB1:HN11	1:B:150:GLY:N	1.66	0.93
4:A:235:CB1:N1	1:B:150:GLY:N	2.26	0.83
4:A:235:CB1:HN12	1:B:150:GLY:HA2	1.46	0.80
4:A:235:CB1:HN11	1:B:149:GLY:C	1.83	0.80
4:A:235:CB1:N1	1:B:150:GLY:CA	2.46	0.79
1:B:92:ARG:HD3	5:B:2073:HOH:O	1.83	0.78
1:A:149:GLY:C	4:B:236:CB1:HN11	1.89	0.76
4:A:235:CB1:N1	1:B:150:GLY:HA2	2.03	0.74
3:A:233:FAD:O4B	1:B:200:ARG:HD3	1.88	0.74
1:B:56:THR:O	1:B:56:THR:CG2	2.34	0.73
4:A:235:CB1:N1	1:B:149:GLY:C	2.43	0.72
1:A:89:LYS:HG3	5:A:2090:HOH:O	1.92	0.69
1:A:229:GLY:HA3	1:A:230:GLN:HB2	1.75	0.67
1:B:51:THR:OG1	1:B:53:LYS:HE3	1.95	0.65

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:26:VAL:O	1:A:30:SER:HB2	1.97	0.64
1:A:229:GLY:CA	1:A:230:GLN:HB2	2.29	0.62
1:A:54:ASP:OD2	1:A:118:ARG:HD2	1.98	0.62
1:A:200:ARG:HD3	3:B:234:FAD:O4B	2.03	0.59
1:A:150:GLY:HA2	4:B:236:CB1:N1	2.19	0.58
1:B:218:GLU:HG3	5:B:2125:HOH:O	2.05	0.56
1:B:216:LYS:CE	5:B:2121:HOH:O	2.56	0.54
1:A:92:ARG:NH2	5:A:2090:HOH:O	2.40	0.54
1:B:168:LEU:HD13	1:B:186:PRO:HG3	1.89	0.54
1:A:149:GLY:C	4:B:236:CB1:N1	2.60	0.53
3:A:233:FAD:H51A	1:B:17:PHE:HB2	1.90	0.53
1:B:177:HIS:HD2	1:B:177:HIS:O	1.92	0.52
4:B:236:CB1:C7	4:B:236:CB1:O3	2.58	0.51
1:B:225:HIS:CD2	1:B:229:GLY:O	2.63	0.51
1:B:92:ARG:NH2	5:B:2074:HOH:O	2.42	0.51
1:B:187:GLN:HE21	1:B:210:ARG:HH11	1.58	0.51
1:B:49:ARG:O	1:B:118:ARG:HD3	2.10	0.51
1:A:187:GLN:HE21	1:A:210:ARG:HH11	1.59	0.50
1:B:218:GLU:CD	5:B:2125:HOH:O	2.49	0.50
1:B:54:ASP:OD2	1:B:118:ARG:HD2	2.12	0.49
1:A:17:PHE:HB2	3:B:234:FAD:H51A	1.94	0.48
1:A:2:GLY:O	1:A:3:LYS:HG2	2.13	0.48
1:B:218:GLU:CG	5:B:2125:HOH:O	2.61	0.48
1:A:157:LYS:HZ2	1:A:230:GLN:H	1.61	0.47
1:B:85:THR:HG21	5:B:2063:HOH:O	2.14	0.47
1:B:60:SER:O	1:B:62:PRO:HD3	2.15	0.46
1:A:83:ASP:OD2	1:A:118:ARG:NH2	2.31	0.46
1:A:67:TYR:CZ	1:A:71:THR:HG21	2.52	0.44
1:A:4:LYS:HG2	1:A:35:THR:HB	2.00	0.44
4:A:235:CB1:H1	5:A:2167:HOH:O	2.18	0.44
3:A:233:FAD:O4B	1:B:200:ARG:CD	2.64	0.44
1:B:177:HIS:CD2	1:B:177:HIS:O	2.71	0.43
1:B:195:ALA:HB1	1:B:199:GLU:HB2	2.00	0.43
1:B:205:ALA:O	1:B:209:GLN:HG3	2.19	0.43
1:A:150:GLY:CA	4:B:236:CB1:N1	2.82	0.43
1:B:216:LYS:CD	5:B:2121:HOH:O	2.33	0.43
1:B:70:GLU:OE1	1:B:70:GLU:HA	2.19	0.42
4:A:235:CB1:C3	4:A:235:CB1:O2	2.67	0.42
1:B:138:GLN:HA	1:B:180:GLY:O	2.20	0.42
1:A:138:GLN:HA	1:A:180:GLY:O	2.20	0.42
1:A:166:TYR:CD2	1:B:228:PHE:CE2	3.08	0.42
1:B:22:LYS:HE3	1:B:22:LYS:HB3	1.92	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:67:TYR:CZ	1:B:71:THR:HG21	2.55	0.41
1:B:73:GLU:O	1:B:77:GLN:HG2	2.21	0.41
1:B:225:HIS:HD2	1:B:229:GLY:O	2.03	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:58:THR:CG2	1:B:218:GLU:CG[4_556]	1.67	0.53
1:B:58:THR:OG1	1:B:218:GLU:CB[4_556]	1.81	0.39
1:B:58:THR:OG1	1:B:218:GLU:CG[4_556]	2.14	0.06

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/236 (97%)	223 (98%)	5 (2%)	0	100	100
1	B	228/236 (97%)	219 (96%)	9 (4%)	0	100	100
All	All	456/472 (97%)	442 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/198 (98%)	186 (96%)	8 (4%)	41	35
1	B	194/198 (98%)	188 (97%)	6 (3%)	52	49
All	All	388/396 (98%)	374 (96%)	14 (4%)	47	42

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	VAL
1	A	53	LYS
1	A	58	THR
1	A	60	SER
1	A	79	SER
1	A	160	VAL
1	A	204	VAL
1	A	230	GLN
1	B	24	VAL
1	B	63	GLU
1	B	127	ASP
1	B	160	VAL
1	B	182	LYS
1	B	194	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	172	GLN
1	A	187	GLN
1	A	212	GLN
1	B	172	GLN
1	B	177	HIS
1	B	187	GLN
1	B	225	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	FAD	A	233	-	58,58,58	1.10	2 (3%)	85,89,89	1.94	18 (21%)
4	CB1	A	235	-	19,19,19	2.44	6 (31%)	28,28,28	3.93	12 (42%)
3	FAD	B	234	-	58,58,58	1.24	8 (13%)	85,89,89	1.86	17 (20%)
4	CB1	B	236	-	19,19,19	2.24	5 (26%)	28,28,28	4.03	14 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	233	-	-	0/34/50/50	0/1/6/6
4	CB1	A	235	-	-	0/16/18/18	0/1/2/2
3	FAD	B	234	-	-	0/34/50/50	0/1/6/6
4	CB1	B	236	-	-	0/16/18/18	0/1/2/2

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	236	CB1	C8-C5	6.16	1.48	1.40
4	A	235	CB1	C8-C5	6.13	1.48	1.40
4	A	235	CB1	C9-C7	5.43	1.62	1.46
4	B	236	CB1	C8-N3	-4.11	1.40	1.46
3	B	234	FAD	C5X-N5	3.79	1.41	1.35
4	B	236	CB1	C9-C7	3.77	1.57	1.46
3	B	234	FAD	C2A-N3A	3.64	1.39	1.32
3	A	233	FAD	C1'-C2'	3.60	1.55	1.51
4	A	235	CB1	C8-N3	-3.40	1.41	1.46
4	A	235	CB1	C1-N	-3.23	1.41	1.46
3	A	233	FAD	C2A-N3A	3.12	1.38	1.32

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	236	CB1	C1-N	-3.11	1.41	1.46
3	B	234	FAD	C1'-C2'	2.91	1.54	1.51
4	A	235	CB1	O4-N3	-2.79	1.20	1.25
3	B	234	FAD	C2A-N1A	2.49	1.38	1.33
4	A	235	CB1	C7-N6	2.25	1.49	1.46
4	B	236	CB1	C9-N6	-2.19	1.43	1.46
3	B	234	FAD	C10-N1	2.14	1.39	1.35
3	B	234	FAD	C9A-C5X	-2.12	1.38	1.42
3	B	234	FAD	C1'-N10	2.12	1.50	1.48
3	B	234	FAD	PA-O3P	2.00	1.63	1.59

All (61) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	236	CB1	C7-N6-C5	13.04	148.46	121.80
3	A	233	FAD	N3A-C2A-N1A	-12.02	118.66	128.71
4	A	235	CB1	C9-N6-C5	10.70	143.66	121.80
3	B	234	FAD	N3A-C2A-N1A	-10.43	119.99	128.71
4	B	236	CB1	C9-N6-C5	10.03	142.30	121.80
4	A	235	CB1	C7-N6-C5	9.44	141.11	121.80
4	A	235	CB1	C9-N6-C7	8.44	66.73	60.76
4	B	236	CB1	C9-N6-C7	7.90	66.35	60.76
4	A	235	CB1	C9-C7-N6	-6.25	55.07	59.61
4	A	235	CB1	C4-C5-N6	-5.70	114.99	122.72
4	B	236	CB1	C2-C3-N1	4.13	124.14	118.11
3	B	234	FAD	O5B-PA-O1A	-4.08	93.40	109.37
4	B	236	CB1	C9-C7-N6	-4.08	56.65	59.61
4	A	235	CB1	C-C8-N3	4.00	119.83	115.82
3	B	234	FAD	N3A-C4A-N9A	3.87	132.41	125.43
3	B	234	FAD	C4A-C5A-N7A	-3.81	106.25	109.52
3	B	234	FAD	C4X-N5-C5X	3.62	120.76	116.69
3	A	233	FAD	C5X-C9A-N10	3.60	120.35	116.80
4	B	236	CB1	C7-C9-N6	-3.60	57.00	59.61
4	B	236	CB1	C4-C5-N6	-3.58	117.87	122.72
4	A	235	CB1	C4-C2-C3	-3.58	113.54	121.23
4	A	235	CB1	C4-C2-C1	3.55	121.33	118.24
3	A	233	FAD	N3A-C4A-N9A	3.52	131.79	125.43
3	B	234	FAD	C2-N1-C10	3.37	118.37	114.98
3	A	233	FAD	C4A-C5A-N7A	-3.23	106.75	109.52
3	A	233	FAD	P-O3P-PA	-3.13	122.50	131.68
4	A	235	CB1	O4-N3-O3	-2.84	115.60	121.35
4	B	236	CB1	C5-C8-N3	2.81	123.87	120.45

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	236	CB1	C2-C1-N	2.79	124.88	120.19
4	B	236	CB1	O-C3-N1	-2.73	118.65	122.59
3	A	233	FAD	O5B-PA-O1A	-2.72	98.70	109.37
3	A	233	FAD	C4X-N5-C5X	2.69	119.72	116.69
3	B	234	FAD	O2A-PA-O3P	2.68	117.85	105.14
4	B	236	CB1	O-C3-C2	-2.65	117.07	120.33
3	A	233	FAD	C2A-N3A-C4A	2.61	121.45	114.01
3	B	234	FAD	C8A-N9A-C4A	2.58	108.86	106.90
3	B	234	FAD	N7A-C8A-N9A	-2.51	107.25	114.36
3	A	233	FAD	C5A-C4A-N3A	-2.50	120.27	125.70
3	A	233	FAD	O5B-C5B-C4B	-2.43	100.03	108.94
3	B	234	FAD	C5A-C4A-N3A	-2.43	120.42	125.70
3	A	233	FAD	C2-N1-C10	2.41	117.41	114.98
3	A	233	FAD	O2A-PA-O3P	2.41	116.58	105.14
4	B	236	CB1	C4-C2-C1	2.40	120.32	118.24
3	B	234	FAD	C1B-N9A-C4A	-2.39	122.50	126.64
4	B	236	CB1	C4-C2-C3	-2.30	116.30	121.23
3	A	233	FAD	O4'-C4'-C5'	-2.29	105.42	110.12
4	A	235	CB1	C-C1-N	2.28	118.11	115.82
3	B	234	FAD	O4B-C1B-N9A	-2.23	106.37	108.44
3	B	234	FAD	C8A-N7A-C5A	2.22	110.46	103.58
3	B	234	FAD	C2A-N3A-C4A	2.20	120.28	114.01
3	A	233	FAD	N7A-C8A-N9A	-2.20	108.15	114.36
4	B	236	CB1	C-C8-N3	2.18	118.01	115.82
3	A	233	FAD	C4X-C10-N10	-2.17	119.43	120.51
3	A	233	FAD	C9A-N10-C10	-2.11	119.69	121.77
4	A	235	CB1	C5-C8-N3	2.09	122.99	120.45
3	B	234	FAD	C2B-C1B-N9A	2.08	118.62	113.27
3	B	234	FAD	C4X-C10-N1	-2.06	120.67	122.73
3	B	234	FAD	C4-C4X-C10	2.06	120.27	116.95
4	A	235	CB1	C2-C3-N1	2.04	121.09	118.11
3	A	233	FAD	O2P-P-O3P	2.02	114.72	105.14
3	A	233	FAD	C8A-N7A-C5A	2.01	109.81	103.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	230/236 (97%)	0.20	10 (4%) 34 32	17, 27, 45, 61	0
1	B	228/236 (96%)	0.49	20 (8%) 10 9	18, 31, 52, 69	0
All	All	458/472 (97%)	0.34	30 (6%) 18 17	17, 28, 49, 69	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	GLY	7.7
1	B	1	ALA	7.6
1	B	230	GLN	6.3
1	B	224	ALA	5.7
1	A	1	ALA	5.3
1	B	229	GLY	4.8
1	A	230	GLN	4.8
1	B	56	THR	4.7
1	B	63	GLU	4.4
1	A	229	GLY	4.0
1	B	62	PRO	3.7
1	A	225	HIS	3.6
1	B	128	ILE	3.6
1	B	59	LEU	3.5
1	B	197	GLU	3.4
1	A	57	GLY	3.2
1	A	63	GLU	3.1
1	A	58	THR	2.9
1	B	129	PRO	2.8
1	B	201	LYS	2.7
1	B	225	HIS	2.6
1	B	61	ASN	2.5
1	B	60	SER	2.5
1	B	216	LYS	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	218	GLU	2.4
1	B	198	GLU	2.2
1	A	223	THR	2.2
1	B	194	ILE	2.1
1	B	110	ALA	2.0
1	A	62	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	CB1	A	235	18/18	0.21	3.10	51,52,56,58	0
4	CB1	B	236	18/18	0.25	1.97	50,52,56,57	0
3	FAD	B	234	53/53	0.16	0.56	19,26,48,53	0
3	FAD	A	233	53/53	0.18	0.42	17,22,48,49	0
2	ZN	A	231	1/1	0.05	-1.70	30,30,30,30	0
2	ZN	B	232	1/1	0.03	-10.92	54,54,54,54	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.