



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 05:52 AM GMT

PDB ID : 4BZ5
Title : Crystal structure of Schistosoma mansoni HDAC8
Authors : Marek, M.; Romier, C.
Deposited on : 2013-07-24
Resolution : 1.78 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	4987 (1.80-1.76)
Clashscore	79885	6152 (1.80-1.76)
Ramachandran outliers	78287	6074 (1.80-1.76)
Sidechain outliers	78261	6073 (1.80-1.76)
RSRZ outliers	66119	4990 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	446	
1	B	446	
1	C	446	
1	D	446	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 14813 atoms, of which 16 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HISTONE DEACETYLASE 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	404	Total	C	N	O	S	0	1	0
			3239	2088	539	596	16			
1	B	419	Total	C	N	O	S	0	4	0
			3362	2166	560	619	17			
1	C	418	Total	C	N	O	S	0	6	0
			3366	2169	561	619	17			
1	D	404	Total	C	N	O	S	0	3	0
			3246	2096	538	596	16			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	441	GLY	-	EXPRESSION TAG	UNP A5H660
A	442	SER	-	EXPRESSION TAG	UNP A5H660
A	443	LEU	-	EXPRESSION TAG	UNP A5H660
A	444	VAL	-	EXPRESSION TAG	UNP A5H660
A	445	PRO	-	EXPRESSION TAG	UNP A5H660
A	446	ARG	-	EXPRESSION TAG	UNP A5H660
B	441	GLY	-	EXPRESSION TAG	UNP A5H660
B	442	SER	-	EXPRESSION TAG	UNP A5H660
B	443	LEU	-	EXPRESSION TAG	UNP A5H660
B	444	VAL	-	EXPRESSION TAG	UNP A5H660
B	445	PRO	-	EXPRESSION TAG	UNP A5H660
B	446	ARG	-	EXPRESSION TAG	UNP A5H660
C	441	GLY	-	EXPRESSION TAG	UNP A5H660
C	442	SER	-	EXPRESSION TAG	UNP A5H660
C	443	LEU	-	EXPRESSION TAG	UNP A5H660
C	444	VAL	-	EXPRESSION TAG	UNP A5H660
C	445	PRO	-	EXPRESSION TAG	UNP A5H660
C	446	ARG	-	EXPRESSION TAG	UNP A5H660
D	441	GLY	-	EXPRESSION TAG	UNP A5H660
D	442	SER	-	EXPRESSION TAG	UNP A5H660
D	443	LEU	-	EXPRESSION TAG	UNP A5H660

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	444	VAL	-	EXPRESSION TAG	UNP A5H660
D	445	PRO	-	EXPRESSION TAG	UNP A5H660
D	446	ARG	-	EXPRESSION TAG	UNP A5H660


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total K 2 2	0	0
3	A	2	Total K 2 2	0	0
3	D	2	Total K 2 2	0	0
3	C	2	Total K 2 2	0	0

- Molecule 4 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			14	4	4	6		
4	B	1	Total	C	H	O	0	0
			14	4	4	6		
4	C	1	Total	C	H	O	0	0
			14	4	4	6		
4	D	1	Total	C	H	O	0	0
			14	4	4	6		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	357	Total	O	0	0
			357	357		
5	B	416	Total	O	0	0
			416	416		
5	C	434	Total	O	0	0
			434	434		
5	D	325	Total	O	0	0
			325	325		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: HISTONE DEACETYLASE 8

Chain A:




• Molecule 1: HISTONE DEACETYLASE 8

Chain B:




• Molecule 1: HISTONE DEACETYLASE 8

Chain C:



• Molecule 1: HISTONE DEACETYLASE 8

Chain D:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	70.65Å 70.73Å 98.30Å 75.90° 78.32° 85.59°	Depositor
Resolution (Å)	31.37 – 1.78 31.37 – 1.79	Depositor EDS
% Data completeness (in resolution range)	96.8 (31.37-1.78) 96.8 (31.37-1.79)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 1.78Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.154 , 0.188 0.152 , 0.185	Depositor DCC
R_{free} test set	8327 reflections (5.01%)	DCC
Wilson B-factor (Å ²)	17.9	Xtriage
Anisotropy	0.321	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 45.4	EDS
Estimated twinning fraction	0.064 for -k,-h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 166222 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	14813	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K, ZN, TLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.28	0/3333	0.45	0/4534
1	B	0.30	0/3467	0.47	0/4717
1	C	0.30	0/3477	0.47	0/4731
1	D	0.28	0/3346	0.45	0/4552
All	All	0.29	0/13623	0.46	0/18534

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3239	0	0	10	0
1	B	3362	0	0	18	0
1	C	3366	0	0	10	0
1	D	3246	0	0	11	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	10	4	0	1	0
4	B	10	4	0	0	0
4	C	10	4	0	1	0
4	D	10	4	0	0	0
5	A	357	0	0	1	1
5	B	416	0	0	6	0
5	C	434	0	0	2	1
5	D	325	0	0	2	0
All	All	14797	16	0	46	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (46) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:72:LYS:NZ	5:B:2119:HOH:O	2.12	0.81
1:B:98:ASN:OD1	1:B:99:TYR:N	2.18	0.77
1:C:98[B]:ASN:OD1	1:C:99:TYR:N	2.22	0.73
1:C:35:TYR:CE1	1:C:368[A]:ILE:CG2	2.75	0.70
1:A:35:TYR:CE1	1:A:368:ILE:CG2	2.75	0.69
1:B:427:ASP:OD2	1:B:430:GLN:NE2	2.25	0.69
1:B:35:TYR:CE1	1:B:368[B]:ILE:CG2	2.75	0.69
1:B:260:ASN:ND2	5:B:2299:HOH:O	2.27	0.66
1:C:260:ASN:ND2	5:C:2327:HOH:O	2.30	0.65
1:D:54:ARG:NH1	5:D:2072:HOH:O	2.30	0.63
1:D:240:GLY:O	1:D:243:SER:OG	2.19	0.60
1:D:35:TYR:CE1	1:D:368[B]:ILE:CG2	2.86	0.58
1:A:145:ARG:NH1	5:A:2123:HOH:O	2.39	0.55
1:A:240:GLY:O	1:A:243:SER:OG	2.25	0.55
1:D:365:LYS:CD	1:D:365:LYS:N	2.70	0.55
1:D:368[A]:ILE:CG2	1:D:387:LEU:CD2	2.86	0.54
1:C:35:TYR:CE1	1:C:368[B]:ILE:CG2	2.91	0.54
1:B:35:TYR:CE1	1:B:368[A]:ILE:CG2	2.92	0.53
1:C:240:GLY:O	1:C:243:SER:OG	2.26	0.53
1:B:352:ARG:NE	5:B:2351:HOH:O	2.43	0.52
1:A:7:TYR:CE1	1:A:43:ARG:CD	2.94	0.51
1:C:368[B]:ILE:CG2	1:C:387:LEU:CD2	2.89	0.50
1:D:359:GLU:OE2	1:D:366:MET:N	2.46	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:215:PHE:CE1	1:D:294[B]:ILE:CD1	2.95	0.49
1:B:240:GLY:O	1:B:243:SER:OG	2.32	0.48
1:C:223:ASN:OD1	1:D:13:GLN:NE2	2.47	0.48
1:B:98:ASN:CB	5:B:2159:HOH:O	2.63	0.47
1:A:368:ILE:CG2	1:A:387:LEU:CD2	2.93	0.46
1:B:368[A]:ILE:CG2	1:B:387:LEU:CD2	2.94	0.46
1:A:93:ASP:OD1	1:A:98:ASN:ND2	2.50	0.45
1:B:440:MET:CE	1:B:443:LEU:CD1	2.96	0.44
1:C:215:PHE:CE1	1:C:294:ILE:CD1	3.00	0.43
1:D:38:ILE:N	1:D:39:PRO:CD	2.81	0.43
1:A:43:ARG:NH2	1:B:96:SER:OG	2.51	0.43
1:C:144:LYS:NZ	5:C:2229:HOH:O	2.51	0.43
1:C:20:LYS:NZ	4:C:700:TLA:O4	2.52	0.43
1:A:20:LYS:NZ	4:A:700:TLA:O4	2.52	0.42
1:D:427:ASP:OD2	1:D:430:GLN:N	2.52	0.42
1:B:123:SER:OG	1:B:163:ARG:NH1	2.52	0.42
1:A:13:GLN:NE2	1:B:223:ASN:OD1	2.52	0.42
1:B:215:PHE:CE1	1:B:294[B]:ILE:CD1	3.03	0.41
1:B:144:LYS:NZ	1:B:195:GLU:OE2	2.52	0.41
1:B:341:TYR:OH	5:B:2037:HOH:O	2.21	0.41
1:A:144:LYS:NZ	1:A:195:GLU:OE2	2.53	0.41
1:D:36:LYS:NZ	5:D:2051:HOH:O	2.53	0.41
1:B:179:ARG:NH2	5:B:2225:HOH:O	2.54	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
5:A:2153:HOH:O	5:C:2386:HOH:O[1_565]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/446 (89%)	390 (98%)	7 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	415/446 (93%)	408 (98%)	7 (2%)	0	100	100
1	C	416/446 (93%)	408 (98%)	8 (2%)	0	100	100
1	D	399/446 (90%)	395 (99%)	4 (1%)	0	100	100
All	All	1627/1784 (91%)	1601 (98%)	26 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	352/391 (90%)	350 (99%)	2 (1%)	92	89
1	B	368/391 (94%)	366 (100%)	2 (0%)	94	91
1	C	369/391 (94%)	367 (100%)	2 (0%)	94	91
1	D	354/391 (90%)	352 (99%)	2 (1%)	92	89
All	All	1443/1564 (92%)	1435 (99%)	8 (1%)	92	89

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	HIS
1	A	301	TYR
1	B	141	HIS
1	B	301	TYR
1	C	141	HIS
1	C	301	TYR
1	D	141	HIS
1	D	301	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 12 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TLA	A	700	2	9,9,9	1.11	0	12,12,12	0.91	0
4	TLA	B	700	2	9,9,9	1.09	0	12,12,12	1.04	0
4	TLA	C	700	2	9,9,9	1.21	0	12,12,12	0.86	0
4	TLA	D	700	2	9,9,9	1.16	0	12,12,12	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TLA	A	700	2	-	0/12/12/12	0/0/0/0
4	TLA	B	700	2	-	0/12/12/12	0/0/0/0
4	TLA	C	700	2	-	0/12/12/12	0/0/0/0
4	TLA	D	700	2	-	0/12/12/12	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/446 (90%)	0.01	16 (3%) 36 33	11, 20, 48, 80	0
1	B	419/446 (93%)	-0.11	11 (2%) 53 51	10, 17, 41, 75	0
1	C	418/446 (93%)	-0.13	12 (2%) 49 47	10, 17, 40, 68	0
1	D	404/446 (90%)	0.08	20 (4%) 28 25	12, 20, 53, 74	0
All	All	1645/1784 (92%)	-0.04	59 (3%) 41 37	10, 19, 46, 80	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	425	ILE	6.2
1	D	392	PHE	5.9
1	D	426	TYR	5.9
1	A	428	TYR	5.4
1	D	428	TYR	5.3
1	B	168	THR	4.9
1	D	432	TYR	4.8
1	D	425	ILE	4.5
1	A	426	TYR	4.3
1	C	168	THR	4.2
1	A	432	TYR	4.1
1	C	177	GLN	3.8
1	A	392	PHE	3.7
1	C	81	GLU	3.6
1	B	395	GLU	3.5
1	A	230	LEU	3.5
1	A	402	ASP	3.4
1	A	429	ASP	3.3
1	C	227	ASN	3.1
1	D	365	LYS	3.1
1	C	282[A]	CYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	315	SER	3.0
1	B	82	LYS	3.0
1	B	437	LEU	3.0
1	B	282[A]	CYS	3.0
1	D	402	ASP	3.0
1	D	282[A]	CYS	2.9
1	D	230	LEU	2.9
1	D	429	ASP	2.8
1	B	81	GLU	2.7
1	A	383	PRO	2.7
1	D	383	PRO	2.7
1	D	368[A]	ILE	2.7
1	A	393	PRO	2.6
1	B	402	ASP	2.5
1	A	364	LYS	2.5
1	D	82	LYS	2.5
1	B	183	VAL	2.5
1	B	177	GLN	2.4
1	D	167	SER	2.3
1	C	82	LYS	2.3
1	C	83	GLU	2.2
1	A	228	ASP	2.2
1	A	282[A]	CYS	2.2
1	D	337	GLY	2.2
1	A	384	ASP	2.2
1	C	280	VAL	2.2
1	D	293	ARG	2.1
1	C	228	ASP	2.1
1	D	228	ASP	2.1
1	B	436	ASN	2.1
1	A	81	GLU	2.1
1	A	363	GLY	2.1
1	D	434	LEU	2.1
1	C	439	GLY	2.1
1	D	393	PRO	2.1
1	D	433	GLN	2.0
1	B	185	LEU	2.0
1	C	225	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	TLA	B	700	10/10	0.12	0.61	16,28,45,46	0
4	TLA	D	700	10/10	0.11	-0.29	20,30,38,38	0
4	TLA	C	700	10/10	0.09	-0.34	17,27,38,41	0
4	TLA	A	700	10/10	0.09	-0.82	20,28,35,40	0
3	K	B	600	1/1	0.05	-1.25	17,17,17,17	0
3	K	C	601	1/1	0.06	-1.57	15,15,15,15	0
3	K	C	600	1/1	0.07	-1.67	11,11,11,11	0
3	K	D	600	1/1	0.04	-1.96	18,18,18,18	0
3	K	B	601	1/1	0.07	-2.00	10,10,10,10	0
3	K	A	601	1/1	0.07	-2.90	11,11,11,11	0
3	K	D	601	1/1	0.05	-2.96	12,12,12,12	0
3	K	A	600	1/1	0.03	-3.77	17,17,17,17	0
2	ZN	B	500	1/1	0.02	-4.99	18,18,18,18	0
2	ZN	D	500	1/1	0.01	-6.38	19,19,19,19	0
2	ZN	C	500	1/1	0.02	-6.85	18,18,18,18	0
2	ZN	A	500	1/1	0.01	-8.18	19,19,19,19	0

6.5 Other polymers ⓘ

There are no such residues in this entry.