



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 12:24 AM GMT

PDB ID : 1C0F
Title : CRYSTAL STRUCTURE OF DICTYOSTELIUM CAATP-ACTIN IN COMPLEX WITH GELSOLIN SEGMENT 1
Authors : Matsuura, Y.; Stewart, M.; Kawamoto, M.; Kamiya, N.; Saeki, K.; Yasunaga, T.; Wakabayashi, T.
Deposited on : 1999-07-16
Resolution : 2.40 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	S	127	
2	A	368	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4312 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GELSOLIN SEGMENT 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	S	127	Total	C	N	O	S	8	0	0
			1005	649	166	187	3			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
S	1	MET	-	EXPRESSION ARTIFACT	UNP P06396
S	2	GLY	-	EXPRESSION ARTIFACT	UNP P06396
S	3	SER	-	EXPRESSION ARTIFACT	UNP P06396
S	35	CYS	ASN	ENGINEERED	UNP P06396

- Molecule 2 is a protein called ACTIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	368	Total	C	N	O	S	27	0	0
			2872	1818	482	553	19			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	73	HIC	HIS	MODIFIED RESIDUE	UNP P07830

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	S	2	Total	Ca	0	0
			2	2		

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	249	Total	O	0	0
			249	249		
5	S	152	Total	O	0	0
			152	152		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: GELSOLIN SEGMENT 1

Chain S: 



- Molecule 2: ACTIN

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.00Å 69.55Å 184.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.40	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.40)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.183 , 0.231	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4312	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, HIC, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	S	0.73	1/1031 (0.1%)	1.34	8/1392 (0.6%)
2	A	1.56	10/2920 (0.3%)	1.71	37/3949 (0.9%)
All	All	1.39	11/3951 (0.3%)	1.62	45/5341 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	S	0	2
2	A	0	1
All	All	0	3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	4	ASP	CG-OD2	54.37	2.50	1.25
2	A	3	GLU	CB-CG	-33.48	0.88	1.52
2	A	50	LYS	CB-CG	31.24	2.36	1.52
2	A	326	LYS	CD-CE	24.29	2.12	1.51
1	S	1	MET	C-N	15.47	1.60	1.33
2	A	291	LYS	CG-CD	15.30	2.04	1.52
2	A	4	ASP	CG-OD1	10.23	1.48	1.25
2	A	2	GLY	N-CA	9.59	1.60	1.46
2	A	2	GLY	CA-C	8.47	1.65	1.51
2	A	1	ASP	C-N	7.53	1.46	1.33
2	A	1	ASP	CA-C	5.86	1.68	1.52

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	3	GLU	CA-CB-CG	-39.90	25.62	113.40
2	A	4	ASP	OD1-CG-OD2	-32.15	62.23	123.30
2	A	3	GLU	CB-CG-CD	-27.57	39.75	114.20
2	A	4	ASP	CB-CG-OD1	22.42	138.48	118.30
2	A	256	ARG	CD-NE-CZ	18.21	149.09	123.60
1	S	1	MET	C-N-CA	18.10	160.31	122.30
2	A	372	ARG	NE-CZ-NH2	-16.83	111.89	120.30
2	A	62	ARG	CD-NE-CZ	15.12	144.77	123.60
2	A	50	LYS	CA-CB-CG	-14.79	80.86	113.40
2	A	256	ARG	NE-CZ-NH2	-11.03	114.78	120.30
2	A	372	ARG	NE-CZ-NH1	9.42	125.01	120.30
2	A	326	LYS	CG-CD-CE	9.04	139.03	111.90
2	A	1	ASP	O-C-N	-8.70	108.42	123.20
2	A	286	ASP	CB-CG-OD2	8.03	125.53	118.30
2	A	62	ARG	NE-CZ-NH1	8.01	124.30	120.30
2	A	1	ASP	CB-CA-C	-7.77	94.86	110.40
2	A	50	LYS	CB-CG-CD	-7.23	92.79	111.60
2	A	266	PHE	CB-CG-CD2	7.09	125.76	120.80
2	A	375	PHE	CA-CB-CG	6.97	130.62	113.90
1	S	88	ASP	CB-CG-OD1	6.78	124.40	118.30
2	A	290	ARG	NE-CZ-NH2	-6.69	116.95	120.30
2	A	266	PHE	CB-CG-CD1	-6.64	116.16	120.80
1	S	89	TYR	CB-CG-CD1	-6.62	117.03	121.00
2	A	222	ASP	CB-CG-OD1	6.55	124.20	118.30
2	A	256	ARG	NE-CZ-NH1	6.45	123.53	120.30
2	A	375	PHE	CB-CG-CD1	-6.44	116.29	120.80
1	S	37	TYR	CB-CG-CD1	6.33	124.80	121.00
2	A	1	ASP	CA-C-N	6.28	128.76	116.20
2	A	99	GLU	OE1-CD-OE2	-6.23	115.82	123.30
2	A	169	TYR	CB-CG-CD2	-6.22	117.27	121.00
1	S	1	MET	CG-SD-CE	6.12	110.00	100.20
2	A	176	LEU	CA-CB-CG	5.95	128.99	115.30
2	A	116	ARG	NE-CZ-NH2	-5.64	117.48	120.30
1	S	1	MET	CB-CA-C	5.64	121.68	110.40
2	A	291	LYS	CB-CG-CD	-5.52	97.24	111.60
2	A	375	PHE	CB-CG-CD2	5.52	124.67	120.80
2	A	119	MET	CA-CB-CG	5.44	122.55	113.30
2	A	81	ASP	CB-CG-OD1	5.41	123.17	118.30
2	A	169	TYR	CB-CG-CD1	5.41	124.25	121.00
2	A	25	ASP	CB-CG-OD1	5.34	123.11	118.30
2	A	294	TYR	CB-CG-CD2	-5.24	117.86	121.00
1	S	111	TYR	CB-CG-CD2	-5.15	117.91	121.00
2	A	157	ASP	CB-CG-OD1	5.13	122.92	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	293	LEU	CB-CG-CD1	5.09	119.65	111.00
1	S	55	ARG	CB-CG-CD	5.02	124.66	111.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	4	ASP	Sidechain
1	S	1	MET	Mainchain,Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	S	1005	0	971	1	7
2	A	2872	0	2830	11	0
3	A	1	0	0	0	0
3	S	2	0	0	0	0
4	A	31	0	12	0	0
5	A	249	0	0	0	0
5	S	152	0	0	0	7
All	All	4312	0	3813	12	7

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (12) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:A:257:CYS:HB3	2:A:258:PRO:HD3	1.87	0.55
2:A:190:MET:HG3	2:A:209:VAL:HG11	1.92	0.52
2:A:220:ALA:HB1	2:A:226:GLU:HG3	1.95	0.49
2:A:10:ILE:HB	2:A:105:LEU:HD23	1.99	0.44
2:A:105:LEU:HD12	2:A:132:MET:HE2	1.99	0.44
2:A:242:LEU:HB3	2:A:243:PRO:HD2	2.00	0.44
2:A:219:VAL:HG22	2:A:258:PRO:HB2	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:A:272:ALA:HB1	2:A:276:GLU:HG2	2.01	0.43
1:S:5:VAL:HG12	1:S:7:HIS:H	1.83	0.43
2:A:332:PRO:HA	2:A:333:PRO:HD3	1.91	0.42
2:A:113:LYS:HG2	2:A:371:HIS:CE1	2.55	0.41
2:A:300:SER:HA	2:A:335:ARG:HB2	2.03	0.41

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:S:1:MET:CA	5:S:219:HOH:O[3_645]	1.31	0.89
1:S:1:MET:CE	5:S:168:HOH:O[3_645]	1.45	0.75
1:S:1:MET:N	5:S:219:HOH:O[3_645]	1.46	0.74
1:S:1:MET:SD	5:S:239:HOH:O[3_645]	1.46	0.74
1:S:1:MET:CG	5:S:255:HOH:O[3_645]	1.81	0.39
1:S:1:MET:CB	5:S:255:HOH:O[3_645]	2.03	0.17
1:S:1:MET:N	5:S:255:HOH:O[3_645]	2.16	0.04

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	125/127 (98%)	121 (97%)	4 (3%)	0	100	100
2	A	361/368 (98%)	353 (98%)	7 (2%)	1 (0%)	50	68
All	All	486/495 (98%)	474 (98%)	11 (2%)	1 (0%)	56	74

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	4	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	S	104/104 (100%)	102 (98%)	2 (2%)	69	87
2	A	309/309 (100%)	297 (96%)	12 (4%)	43	64
All	All	413/413 (100%)	399 (97%)	14 (3%)	49	70

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	S	36	LEU
1	S	55	ARG
2	A	1	ASP
2	A	3	GLU
2	A	51	ASP
2	A	62	ARG
2	A	176	LEU
2	A	185	LEU
2	A	228	GLN
2	A	233	SER
2	A	293	LEU
2	A	300	SER
2	A	354	GLN
2	A	375	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	280	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HIC	A	73	2	11,11,12	5.64	3 (27%)	12,14,16	1.72	3 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HIC	A	73	2	-	0/4/6/8	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	73	HIC	O-C	18.12	1.23	1.11
2	A	73	HIC	CA-C	3.18	1.54	1.48
2	A	73	HIC	CE1-NE2	-2.90	1.32	1.36

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	73	HIC	C-CA-N	-3.50	110.33	113.83
2	A	73	HIC	CE1-ND1-CG	2.48	109.62	104.93
2	A	73	HIC	ND1-CE1-NE2	-2.16	107.47	112.55

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	ATP	A	376	3	33,33,33	1.11	2 (6%)	52,52,52	1.75	6 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	376	3	-	0/22/38/38	0/1/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	376	ATP	C8-N7	-2.60	1.29	1.34
4	A	376	ATP	C5-N7	-2.12	1.32	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	376	ATP	C8-N9-C4	-6.91	101.62	106.90
4	A	376	ATP	C4-C5-N7	4.75	113.59	109.52
4	A	376	ATP	O4'-C1'-N9	4.59	112.71	108.44
4	A	376	ATP	N3-C2-N1	-4.05	125.33	128.71
4	A	376	ATP	C1'-N9-C4	2.99	131.81	126.64
4	A	376	ATP	O4'-C4'-C3'	2.08	109.38	105.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.