



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 03:22 AM GMT

PDB ID : 1C0R  
Title : COMPLEX OF VANCOMYCIN WITH D-LACTIC ACID  
Authors : Loll, P.J.; Kaplan, J.; Selinsky, B.; Axelsen, P.H.  
Deposited on : 1999-07-20  
Resolution : 1.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : NOT EXECUTED  
EDS : NOT EXECUTED  
Percentile statistics : 21963  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1028 (1.10-0.90)
Ramachandran outliers	78287	1056 (1.12-0.88)
Sidechain outliers	78261	1055 (1.12-0.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	7	
1	B	7	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 268 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VANCOMYCIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	7	Total	C	Cl	N	O	0	0	0
			80	53	2	8	17			
1	B	7	Total	C	Cl	N	O	0	1	0
			81	53	2	8	18			


- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			21	13	1	7		
2	B	2	Total	C	N	O	0	1
			32	19	1	12		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	2	Total	Cl	0	0
			2	2		

- Molecule 4 is LACTIC ACID (three-letter code: LAC) (formula: C<sub>3</sub>H<sub>6</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	25	Total	O	0	2
			25	25		
5	B	20	Total	O	0	5
			20	20		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: VANCOMYCIN

Chain A: 



- Molecule 1: VANCOMYCIN

Chain B: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	28.44Å 28.44Å 66.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.00	Depositor
% Data completeness (in resolution range)	97.0 (20.00-1.00)	Depositor
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	SHELXL-93	Depositor
R, $R_{free}$	0.114 , 0.142	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	268	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	10.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, GHP, CL, RER, LAC, OMZ, MLU, 3FG, OMY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.58	0/7	1.26	0/8
1	B	1.59	0/7	0.73	0/8
All	All	1.20	0/14	1.03	0/16

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	80	0	45	1	0
1	B	81	0	40	2	0
2	A	21	0	21	1	0
2	B	32	0	29	1	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
4	B	6	0	5	0	0
5	A	25	0	0	0	0
5	B	20	0	0	1	0
All	All	268	0	140	3	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (3) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:A:8:BGC:H6C1	5:B:2017[A]:HOH:O	1.96	0.64
1:B:6:OMY:CL	2:B:8[B]:BGC:H6C1	2.42	0.56
1:A:4:GHP:HA	1:B:5:GHP:O	2.14	0.47

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1/7 (14%)	0	1 (100%)	0	100	100
1	B	1/7 (14%)	1 (100%)	0	0	100	100
All	All	2/14 (14%)	1 (50%)	1 (50%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1/1 (100%)	1 (100%)	0	100	100
1	B	1/1 (100%)	1 (100%)	0	100	100
All	All	2/2 (100%)	2 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.



Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

Of 13 non-standard protein/DNA/RNA residues modelled in this entry, 2 are modelled with single atom - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	MLU	A	1	1	8,8,9	5.78	3 (37%)	6,9,11	0.50	0
1	OMZ	A	2	1	14,14,15	5.26	4 (28%)	17,19,21	1.43	4 (23%)
1	GHP	A	4	1,2	11,11,12	5.32	2 (18%)	12,14,16	10.00	1 (8%)
1	GHP	A	5	1	11,11,12	6.45	2 (18%)	12,14,16	13.84	2 (16%)
1	OMY	A	6	1	14,14,15	4.90	2 (14%)	17,19,21	1.69	3 (17%)
1	3FG	A	7	1	13,13,13	1.00	1 (7%)	18,18,18	1.21	1 (5%)
1	MLU	B	1	1	8,8,9	6.90	2 (25%)	6,9,11	0.59	0
1	OMZ	B	2	1	14,14,15	4.88	2 (14%)	17,19,21	1.16	2 (11%)
1	GHP	B	5	1	11,11,12	6.97	2 (18%)	12,14,16	18.43	1 (8%)
1	OMY	B	6	1	14,14,15	5.33	3 (21%)	17,19,21	1.47	2 (11%)
1	3FG	B	7	1	13,13,13	1.32	3 (23%)	18,18,18	1.12	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLU	A	1	1	-	0/6/8/10	0/0/0/0
1	OMZ	A	2	1	-	0/8/10/12	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	GHP	A	4	1,2	-	0/4/6/8	0/1/1/1
1	GHP	A	5	1	-	0/4/6/8	0/1/1/1
1	OMY	A	6	1	-	0/8/10/12	0/1/1/1
1	3FG	A	7	1	-	0/8/8/8	0/1/1/1
1	MLU	B	1	1	-	0/6/8/10	0/0/0/0
1	OMZ	B	2	1	-	0/8/10/12	0/1/1/1
1	GHP	B	5	1	-	0/4/6/8	0/1/1/1
1	OMY	B	6	1	-	0/8/10/12	0/1/1/1
1	3FG	B	7	1	-	0/8/8/8	0/1/1/1

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	5	GHP	O-C	21.48	1.26	1.11
1	A	5	GHP	O-C	19.60	1.24	1.11
1	B	6	OMY	O-C	19.48	1.24	1.11
1	B	1	MLU	O-C	19.12	1.24	1.11
1	A	2	OMZ	O-C	18.76	1.24	1.11
1	B	2	OMZ	O-C	17.65	1.23	1.11
1	A	6	OMY	O-C	17.53	1.23	1.11
1	A	4	GHP	O-C	17.16	1.23	1.11
1	A	1	MLU	O-C	15.82	1.22	1.11
1	A	5	GHP	CA-C	8.32	1.57	1.48
1	B	5	GHP	CA-C	8.03	1.57	1.48
1	B	2	OMZ	CA-C	3.80	1.55	1.48
1	A	4	GHP	CA-C	3.26	1.51	1.48
1	A	6	OMY	CA-C	3.21	1.54	1.48
1	A	2	OMZ	CA-C	3.18	1.54	1.48
1	B	1	MLU	CA-C	3.16	1.54	1.48
1	A	2	OMZ	CZ-CE1	-2.95	1.36	1.39
1	A	1	MLU	CA-C	2.92	1.53	1.48
1	B	6	OMY	CA-C	2.80	1.53	1.48
1	A	1	MLU	CN-N	2.63	1.55	1.46
1	A	2	OMZ	OH-CZ	2.47	1.41	1.36
1	B	7	3FG	CG1-CD1	2.46	1.43	1.39
1	B	6	OMY	OCZ-CZ	2.34	1.41	1.36
1	B	7	3FG	CA-C	2.28	1.59	1.52
1	A	7	3FG	CG1-CB	2.19	1.42	1.38
1	B	7	3FG	OXT-C	-2.05	1.22	1.30

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	5	GHP	C-CA-N	-63.74	106.05	113.27
1	A	5	GHP	C-CA-N	-47.72	107.86	113.27
1	A	4	GHP	C-CA-N	-34.55	109.35	113.27
1	A	6	OMY	CG-CB-CA	-4.21	105.52	111.75
1	A	6	OMY	CD1-CE1-CZ	-4.16	118.97	121.13
1	B	6	OMY	CG-CB-CA	-3.88	106.00	111.75
1	B	2	OMZ	OC-CB-CA	-3.14	101.49	109.21
1	A	2	OMZ	CD1-CE1-CZ	-3.05	119.54	121.13
1	A	2	OMZ	CE2-CZ-CE1	2.92	121.32	118.80
1	B	2	OMZ	CB-CA-N	2.76	113.40	109.14
1	A	5	GHP	C1-CA-N	2.71	120.41	112.92
1	A	2	OMZ	CB-CA-N	2.70	113.30	109.14
1	B	6	OMY	CE2-CZ-CE1	2.66	121.10	118.80
1	A	6	OMY	CE2-CZ-CE1	2.54	121.00	118.80
1	A	7	3FG	CD1-CG1-CB	-2.47	117.92	120.05
1	B	7	3FG	OXT-C-O	2.26	129.18	124.07
1	A	2	OMZ	CE2-CD2-CG	-2.12	118.96	121.20
1	B	7	3FG	CD2-CG2-CB	2.09	121.86	120.05

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.5 Carbohydrates ⓘ

5 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	BGC	A	8	1,2	10,11,12	1.54	2 (20%)	11,15,17	0.87	0
2	RER	A	9	2	9,10,11	0.97	1 (11%)	12,15,17	1.23	2 (16%)
2	BGC	B	8[A]	1,2	10,11,12	1.38	1 (10%)	11,15,17	1.94	4 (36%)
2	BGC	B	8[B]	1,2	10,11,12	1.92	3 (30%)	11,15,17	2.39	3 (27%)
2	RER	B	9	2	9,10,11	1.16	1 (11%)	12,15,17	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	8	1,2	-	0/2/19/22	0/1/1/1
2	RER	A	9	2	-	0/0/17/19	0/1/1/1
2	BGC	B	8[A]	1,2	-	0/2/19/22	0/1/1/1
2	BGC	B	8[B]	1,2	-	0/2/19/22	0/1/1/1
2	RER	B	9	2	-	0/0/17/19	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	8[B]	BGC	C4-C5	3.82	1.61	1.53
2	A	8	BGC	C4-C5	3.76	1.61	1.53
2	B	8[A]	BGC	C4-C5	3.30	1.60	1.53
2	B	8[B]	BGC	O5-C5	2.86	1.50	1.45
2	B	9	RER	C2-C3	2.85	1.57	1.53
2	B	8[B]	BGC	O6-C6	-2.83	1.30	1.42
2	A	9	RER	O4-C4	2.51	1.48	1.42
2	A	8	BGC	O6-C6	-2.32	1.32	1.42

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	8[B]	BGC	O5-C5-C4	-5.09	104.19	110.65
2	B	8[B]	BGC	O5-C5-C6	4.97	112.20	106.98
2	B	8[A]	BGC	O5-C5-C4	-4.01	105.56	110.65
2	B	8[A]	BGC	O5-C5-C6	2.97	110.10	106.98
2	A	9	RER	C5A-C5-C4	2.50	117.07	112.59
2	B	8[B]	BGC	O2-C2-C3	-2.50	104.79	110.18
2	A	9	RER	O4-C4-C3	-2.44	102.22	108.70
2	B	8[A]	BGC	C4-C3-C2	2.15	113.39	110.50
2	B	8[A]	BGC	O6-C6-C5	2.11	118.63	111.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	LAC	B	23	-	5,5,5	0.87	0	6,6,6	1.40	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LAC	B	23	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	23	LAC	O1-C1-C2	-2.38	116.64	122.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.