



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 03:52 AM GMT

PDB ID : 2C0T
Title : SRC FAMILY KINASE HCK WITH BOUND INHIBITOR A-641359
Authors : Borhani, D.W.; Burchat, A.; Calderwood, D.J.; Hirst, G.C.; Li, B.; Loew, A.
Deposited on : 2005-09-07
Resolution : 2.15 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1094 (2.18-2.14)
Clashscore	79885	1299 (2.18-2.14)
Ramachandran outliers	78287	1272 (2.18-2.14)
Sidechain outliers	78261	1272 (2.18-2.14)
RSRZ outliers	66119	1094 (2.18-2.14)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	454	
1	B	454	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7225 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TYROSINE-PROTEIN KINASE HCK.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	428	Total	C	N	O	P	S	0	1	0
			3467	2217	584	645	1	20			
1	B	430	Total	C	N	O	P	S	0	0	0
			3468	2217	583	647	1	20			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	502	GLU	GLN	ENGINEERED MUTATION	UNP P08631
A	503	GLU	GLN	ENGINEERED MUTATION	UNP P08631
A	504	ILE	GLN	ENGINEERED MUTATION	UNP P08631
B	502	GLU	GLN	ENGINEERED MUTATION	UNP P08631
B	503	GLU	GLN	ENGINEERED MUTATION	UNP P08631
B	504	ILE	GLN	ENGINEERED MUTATION	UNP P08631

- Molecule 2 is N-(4-{4-AMINO-1-[1-(TETRAHYDRO-2H-PYRAN-4-YL)PIPERIDIN-4-YL]-1H-PYRAZOLO[3,4-D]PYRIMIDIN-3-YL}-2-METHOXYPHENYL)-1-METHYL-1H-INDOLE-2-CARBOXAMIDE (three-letter code: L3G) (formula: C₃₂H₃₆N₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			43	32	8	3		
2	B	1	Total	C	N	O	0	0
			43	32	8	3		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Ca	0	0
			2	2		
3	A	2	Total	Ca	0	0
			2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	129	Total	O	0	0
			129	129		
4	B	71	Total	O	0	0
			71	71		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: TYROSINE-PROTEIN KINASE HCK

Chain A:



• Molecule 1: TYROSINE-PROTEIN KINASE HCK

Chain B:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	48.86Å 72.91Å 180.00Å 90.00° 95.87° 90.00°	Depositor
Resolution (Å)	20.00 – 2.15 40.44 – 2.15	Depositor EDS
% Data completeness (in resolution range)	70.8 (20.00-2.15) 70.5 (40.44-2.15)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.15 (at 2.14Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.194 , 0.253 0.198 , 0.256	Depositor DCC
R_{free} test set	2442 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	39.6	Xtriage
Anisotropy	0.688	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 60.5	EDS
Estimated twinning fraction	0.024 for h,-k,-h-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 48624 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7225	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: L3G, CA, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/3533	0.70	0/4767
1	B	0.57	0/3532	0.65	0/4764
All	All	0.62	0/7065	0.68	0/9531

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3467	0	3426	23	0
1	B	3468	0	3429	42	0
2	A	43	0	36	0	0
2	B	43	0	36	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	129	0	0	5	0
4	B	71	0	0	3	0
All	All	7225	0	6927	61	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (61) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:71:ILE:HD11	1:B:227:LYS:HE2	1.32	1.03
1:A:114:ARG:O	1:A:117:SER:HB2	1.83	0.79
1:A:214:GLN:NE2	4:A:2028:HOH:O	2.15	0.78
1:B:71:ILE:CD1	1:B:227:LYS:HE2	2.15	0.71
1:A:180:ARG:NH2	1:B:463:GLU:OE2	2.26	0.67
1:A:382:ALA:HB2	4:A:2083:HOH:O	1.98	0.63
1:B:352:GLU:HG3	1:B:415:ILE:HG12	1.80	0.63
1:A:180:ARG:HG2	4:A:2026:HOH:O	2.00	0.61
1:B:189:ILE:HG22	1:B:504:ILE:HG21	1.80	0.61
1:B:130:ARG:NH1	4:B:2063:HOH:O	2.34	0.60
1:A:195:PHE:HD2	1:A:200:GLU:HG2	1.65	0.60
1:A:463:GLU:OE2	1:B:180:ARG:NH2	2.36	0.59
1:B:195:PHE:HD2	1:B:200:GLU:HG2	1.69	0.56
1:B:281:PHE:HE2	1:B:310:ILE:HD11	1.71	0.56
1:B:417:SER:HB2	4:B:2044:HOH:O	2.06	0.55
1:A:190:SER:OG	1:A:192:ARG:HD2	2.07	0.55
1:A:463:GLU:OE2	1:B:180:ARG:NH1	2.41	0.54
1:B:425:LEU:O	1:B:428:GLU:HB2	2.08	0.54
1:B:359:ARG:HG2	1:B:382:ALA:CB	2.37	0.54
1:A:268:VAL:HG22	1:A:311:ILE:HG13	1.90	0.54
1:B:144:LEU:HD12	1:B:214:GLN:HG2	1.90	0.53
1:B:333:PRO:HB2	1:B:335:PRO:HD2	1.91	0.53
1:B:186:GLY:HA2	1:B:195:PHE:O	2.10	0.52
1:B:359:ARG:HG2	1:B:382:ALA:HB1	1.91	0.52
1:A:442:ASN:HB2	1:A:443:PRO:HD3	1.92	0.52
1:A:283:ALA:O	1:A:287:VAL:HG23	2.10	0.51
1:A:357:ILE:HD11	4:A:2089:HOH:O	2.10	0.51
1:A:281:PHE:HE2	1:A:310:ILE:HD11	1.76	0.51
1:B:146:SER:HA	1:B:218:VAL:O	2.10	0.51
1:B:281:PHE:CE2	1:B:310:ILE:HD11	2.46	0.50
1:B:378:ASP:OD1	2:B:1506:L3G:H9	2.11	0.50
1:B:195:PHE:CD2	1:B:200:GLU:HG2	2.48	0.49
1:B:293:HIS:HB3	1:B:296:LEU:HG	1.94	0.49
1:B:359:ARG:CG	1:B:382:ALA:HB1	2.43	0.49
1:A:253:GLY:HA3	1:A:270:THR:O	2.13	0.48
1:A:169:ARG:NH2	1:B:169:ARG:HH21	2.12	0.48
1:B:442:ASN:HB2	1:B:443:PRO:HD3	1.96	0.47
1:A:202:VAL:O	1:A:206:LYS:HG3	2.15	0.47
1:B:269:LYS:NZ	4:B:2023:HOH:O	2.10	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:209:ASN:O	1:B:209:ASN:CG	2.54	0.46
1:B:334:LEU:HD21	1:B:462:PRO:HD3	1.97	0.46
1:B:189:ILE:HG22	1:B:504:ILE:CG2	2.46	0.46
1:A:382:ALA:CB	4:A:2083:HOH:O	2.59	0.45
1:B:504:ILE:HA	1:B:505:PRO:HD3	1.76	0.45
1:B:380:GLY:C	1:B:381:LEU:N	2.70	0.45
1:B:198:LEU:O	1:B:202:VAL:HG13	2.17	0.45
1:B:457:ARG:HG3	1:B:466:TYR:CG	2.52	0.44
1:B:295:LYS:HA	1:B:295:LYS:HD3	1.78	0.44
1:B:323:PHE:O	1:B:329:GLY:HA3	2.18	0.44
1:B:268:VAL:HG22	1:B:311:ILE:HG13	1.99	0.44
1:B:180:ARG:HG3	1:B:188:TYR:CZ	2.53	0.44
1:B:83:GLN:C	1:B:84:MET:HG2	2.39	0.43
1:A:146:SER:HA	1:A:218:VAL:O	2.19	0.43
1:B:189:ILE:CG2	1:B:504:ILE:HG21	2.49	0.42
1:B:111:TYR:HE2	1:B:224:LYS:HE3	1.85	0.42
1:A:426:LEU:HD23	1:A:469:MET:HG2	2.00	0.42
1:A:192:ARG:HG2	1:A:193:SER:N	2.34	0.41
1:A:153:GLU:HB2	1:A:501:PTR:O2P	2.20	0.41
1:A:476:ARG:HA	1:A:477:PRO:HD3	1.93	0.41
1:B:224:LYS:HA	1:B:225:PRO:HD3	1.89	0.41
1:B:317:LYS:HG2	1:B:323:PHE:CD1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	422/454 (93%)	406 (96%)	16 (4%)	0	100	100
1	B	421/454 (93%)	401 (95%)	20 (5%)	0	100	100
All	All	843/908 (93%)	807 (96%)	36 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/393 (95%)	360 (96%)	14 (4%)	45	45
1	B	374/393 (95%)	358 (96%)	16 (4%)	40	38
All	All	748/786 (95%)	718 (96%)	30 (4%)	42	41

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	SER
1	A	117	SER
1	A	128	ILE
1	A	144	LEU
1	A	192	ARG
1	A	194	THR
1	A	199	GLN
1	A	223	SER
1	A	245	LYS
1	A	298	LYS
1	A	357	ILE
1	A	361	LEU
1	A	400	ILE
1	A	459	GLU
1	B	90	SER
1	B	119	GLU
1	B	144	LEU
1	B	164	ARG
1	B	169	ARG
1	B	178	LYS
1	B	192	ARG
1	B	194	THR
1	B	276	MET
1	B	298	LYS
1	B	336	LYS
1	B	359	ARG
1	B	361	LEU
1	B	406	GLU
1	B	417	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	459	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	251	GLN
1	A	261	ASN
1	B	251	GLN
1	B	292	GLN
1	B	442	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PTR	A	501	1,3	16,16,17	5.54	3 (18%)	20,22,24	0.93	2 (10%)
1	PTR	B	501	1,3	16,16,17	6.06	3 (18%)	20,22,24	1.12	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	501	1,3	-	0/9/11/13	0/1/1/1
1	PTR	B	501	1,3	-	0/9/11/13	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	501	PTR	O-C	23.30	1.27	1.11
1	A	501	PTR	O-C	20.99	1.25	1.11
1	A	501	PTR	OH-CZ	-6.03	1.25	1.40
1	B	501	PTR	OH-CZ	-5.66	1.26	1.40
1	A	501	PTR	CA-C	3.11	1.54	1.48
1	B	501	PTR	CA-C	2.76	1.53	1.48

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	501	PTR	C-CA-N	-3.94	109.89	113.83
1	A	501	PTR	CG-CB-CA	-2.11	109.44	114.42
1	A	501	PTR	P-OH-CZ	2.08	129.41	123.55

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	L3G	A	1506	-	49,49,49	1.38	3 (6%)	66,71,71	2.20	16 (24%)
2	L3G	B	1506	-	49,49,49	1.21	3 (6%)	66,71,71	2.31	16 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	L3G	A	1506	-	-	0/17/40/40	0/3/7/7
2	L3G	B	1506	-	-	0/17/40/40	0/3/7/7

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1506	L3G	C4-N2	-6.16	1.30	1.36
2	A	1506	L3G	C4-N2	-5.05	1.31	1.36
2	A	1506	L3G	N1-N2	5.02	1.43	1.37
2	A	1506	L3G	C24-N8	-3.01	1.35	1.39
2	B	1506	L3G	C24-N8	-2.52	1.35	1.39
2	B	1506	L3G	N1-N2	2.12	1.40	1.37

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1506	L3G	N3-C11-N4	-10.64	119.81	128.71
2	A	1506	L3G	N3-C11-N4	-8.82	121.33	128.71
2	A	1506	L3G	C23-N7-C22	6.22	120.66	109.16
2	B	1506	L3G	C21-C26-C24	5.75	111.29	106.27
2	A	1506	L3G	C21-C26-C24	5.72	111.26	106.27
2	B	1506	L3G	C23-N7-C22	5.50	119.31	109.16
2	A	1506	L3G	C20-C21-C26	-5.03	102.05	106.17
2	B	1506	L3G	C27-C22-N7	-4.94	103.52	111.49
2	A	1506	L3G	C20-N8-C24	4.53	110.80	106.34
2	B	1506	L3G	C20-C21-C26	-4.52	102.47	106.17
2	B	1506	L3G	C20-N8-C24	3.87	110.16	106.34
2	A	1506	L3G	C26-C24-N8	-3.55	105.44	109.17
2	B	1506	L3G	C10-N2-N1	-3.49	118.59	121.15
2	B	1506	L3G	O1-C8-C12	3.37	118.73	114.83
2	B	1506	L3G	C26-C24-N8	-3.32	105.68	109.17
2	A	1506	L3G	C10-N2-N1	-3.21	118.80	121.15
2	A	1506	L3G	O1-C8-C12	2.89	118.17	114.83
2	A	1506	L3G	C23-N7-C19	-2.82	105.84	112.63
2	A	1506	L3G	C17-C19-N7	2.69	117.87	112.39
2	A	1506	L3G	C15-O1-C8	2.61	121.45	117.59
2	A	1506	L3G	C28-C10-N2	-2.60	107.50	111.09
2	B	1506	L3G	C11-N4-C5	2.57	123.40	118.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1506	L3G	C27-C22-N7	-2.50	107.46	111.49
2	B	1506	L3G	C15-O1-C8	2.40	121.14	117.59
2	B	1506	L3G	C28-C23-N7	-2.32	107.75	111.49
2	B	1506	L3G	C2-C1-N1	-2.24	106.77	109.03
2	B	1506	L3G	C1-N1-N2	2.17	107.03	105.26
2	A	1506	L3G	C11-N4-C5	2.07	122.51	118.77
2	B	1506	L3G	C22-N7-C19	2.04	117.55	112.63
2	A	1506	L3G	C1-N1-N2	2.03	106.91	105.26
2	A	1506	L3G	C2-C1-N1	-2.02	107.00	109.03
2	B	1506	L3G	C8-C12-N6	2.01	120.41	116.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/454 (94%)	0.69	20 (4%) 30 34	46, 61, 79, 87	0
1	B	430/454 (94%)	0.72	34 (7%) 13 15	51, 64, 82, 94	0
All	All	858/908 (94%)	0.71	54 (6%) 19 22	46, 63, 80, 94	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	278	VAL	8.2
1	B	408	ILE	5.4
1	B	410	PHE	5.2
1	A	187	PHE	4.7
1	B	447	ARG	4.6
1	B	275	SER	4.4
1	B	400	ILE	4.3
1	B	279	GLU	4.2
1	B	271	MET	4.2
1	B	407	ALA	4.1
1	B	276	MET	4.1
1	B	405	PRO	3.6
1	B	452	GLY	3.6
1	B	443	PRO	3.5
1	B	449	LEU	3.5
1	A	194	THR	3.4
1	B	413	PHE	3.4
1	A	189	ILE	3.3
1	B	282	LEU	3.2
1	A	208	GLY	3.1
1	B	207	LYS	2.9
1	B	439	GLY	2.9
1	A	276	MET	2.9
1	B	404	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	442	ASN	2.7
1	B	446	ILE	2.6
1	B	381	LEU	2.6
1	B	409	ASN	2.6
1	A	342	ALA	2.5
1	B	445	VAL	2.5
1	A	192	ARG	2.5
1	A	197	THR	2.4
1	B	193	SER	2.4
1	B	281	PHE	2.4
1	A	296	LEU	2.4
1	A	190	SER	2.3
1	B	412	SER	2.3
1	B	406	GLU	2.3
1	B	357	ILE	2.2
1	A	341	SER	2.2
1	A	279	GLU	2.2
1	B	133	ALA	2.2
1	A	447	ARG	2.1
1	B	305	LYS	2.1
1	A	367	LEU	2.1
1	A	357	ILE	2.1
1	A	180	ARG	2.1
1	A	202	VAL	2.1
1	B	374	CYS	2.1
1	B	283	ALA	2.0
1	A	297	VAL	2.0
1	B	203	ASP	2.0
1	A	468	ILE	2.0
1	A	193	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	PTR	B	501	16/17	0.16	-0.03	55,58,60,62	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	PTR	A	501	16/17	0.14	-0.62	63,66,67,68	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	L3G	B	1506	43/43	0.14	-0.39	49,54,65,66	0
3	CA	B	1507	1/1	0.15	-0.93	53,53,53,53	0
2	L3G	A	1506	43/43	0.16	-1.12	40,46,59,63	0
3	CA	B	1508	1/1	0.05	-3.61	74,74,74,74	0
3	CA	A	1507	1/1	0.11	-3.85	52,52,52,52	0
3	CA	A	1508	1/1	0.03	-8.17	66,66,66,66	0

6.5 Other polymers

There are no such residues in this entry.