



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 10:18 AM GMT

PDB ID : 4C0G
Title : Structure of the NOT-box domain of human CNOT3
Authors : Boland, A.; Chen, Y.; Raisch, T.; Jonas, S.; Izaurralde, E.; Weichenrieder, O.
Deposited on : 2013-08-01
Resolution : 2.40 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance



The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2207 (2.40-2.40)
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)
RSRZ outliers	66119	2210 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	102	
1	B	102	
1	C	102	
1	D	102	
1	E	102	
1	F	102	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9619 atoms, of which 4559 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	90	Total	C	H	N	O	Se	0	0	0
			1562	535	756	126	143	2			
1	B	97	Total	C	H	N	O	Se	0	0	0
			1643	564	787	139	151	2			
1	C	96	Total	C	H	N	O	Se	0	0	0
			1591	550	753	135	151	2			
1	D	96	Total	C	H	N	O	Se	0	0	0
			1618	562	766	138	150	2			
1	E	91	Total	C	H	N	O	Se	0	0	0
			1572	539	761	128	142	2			
1	F	93	Total	C	H	N	O	Se	0	0	0
			1557	543	736	126	150	2			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	652	GLY	-	EXPRESSION TAG	UNP O75175
A	653	ALA	-	EXPRESSION TAG	UNP O75175
A	654	MSE	-	EXPRESSION TAG	UNP O75175
A	655	ALA	-	EXPRESSION TAG	UNP O75175
B	652	GLY	-	EXPRESSION TAG	UNP O75175
B	653	ALA	-	EXPRESSION TAG	UNP O75175
B	654	MSE	-	EXPRESSION TAG	UNP O75175
B	655	ALA	-	EXPRESSION TAG	UNP O75175
C	652	GLY	-	EXPRESSION TAG	UNP O75175
C	653	ALA	-	EXPRESSION TAG	UNP O75175
C	654	MSE	-	EXPRESSION TAG	UNP O75175
C	655	ALA	-	EXPRESSION TAG	UNP O75175
D	652	GLY	-	EXPRESSION TAG	UNP O75175
D	653	ALA	-	EXPRESSION TAG	UNP O75175
D	654	MSE	-	EXPRESSION TAG	UNP O75175
D	655	ALA	-	EXPRESSION TAG	UNP O75175
E	652	GLY	-	EXPRESSION TAG	UNP O75175

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	653	ALA	-	EXPRESSION TAG	UNP O75175
E	654	MSE	-	EXPRESSION TAG	UNP O75175
E	655	ALA	-	EXPRESSION TAG	UNP O75175
F	652	GLY	-	EXPRESSION TAG	UNP O75175
F	653	ALA	-	EXPRESSION TAG	UNP O75175
F	654	MSE	-	EXPRESSION TAG	UNP O75175
F	655	ALA	-	EXPRESSION TAG	UNP O75175

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	10	Total O 10 10	0	0
2	B	14	Total O 14 14	0	0
2	C	13	Total O 13 13	0	0
2	D	10	Total O 10 10	0	0
2	E	15	Total O 15 15	0	0
2	F	14	Total O 14 14	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 3

Chain A: 



- Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 3

Chain B: 



- Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 3

Chain C: 



- Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 3

Chain D: 




- Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 3

Chain E: 



- Molecule 1: CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 3

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.30Å 97.57Å 141.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.06 – 2.40 49.06 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.06-2.40) 99.6 (49.06-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.86 (at 2.39Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.221 , 0.257 0.211 , 0.246	Depositor DCC
R_{free} test set	1478 reflections (5.11%)	DCC
Wilson B-factor (Å ²)	54.6	Xtriage
Anisotropy	0.657	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	2 of 28975 reflections (0.007%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9619	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/832	0.40	0/1116
1	B	0.24	0/883	0.37	0/1184
1	C	0.24	0/863	0.37	0/1158
1	D	0.25	0/879	0.38	0/1178
1	E	0.26	0/837	0.39	0/1122
1	F	0.25	0/847	0.37	0/1138
All	All	0.25	0/5141	0.38	0/6896

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	806	756	0	10	0
1	B	856	787	13	10	0
1	C	838	753	19	8	0
1	D	852	766	21	4	0
1	E	811	761	0	5	0
1	F	821	736	0	9	0
2	A	10	0	0	0	0
2	B	14	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	13	0	0	0	0
2	D	10	0	0	0	0
2	E	15	0	0	0	0
2	F	14	0	0	0	0
All	All	5060	4559	53	43	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (43) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:659:THR:OG1	1:C:661:GLU:OE1	2.15	0.65
1:B:656:HIS:O	1:B:659:THR:OG1	2.17	0.61
1:B:749:ASP:O	1:B:750:ARG:HG2	2.03	0.59
1:B:700:THR:CG2	1:B:748:GLU:HG2	2.34	0.57
1:A:718:GLU:OE1	1:A:718:GLU:N	2.37	0.57
1:E:664:GLN:OE1	1:E:693:LYS:NZ	2.41	0.53
1:C:656:HIS:O	1:C:659:THR:HG23	2.09	0.53
1:F:699:HIS:CE1	1:F:746:TYR:HB3	2.44	0.52
1:A:668:THR:HG23	1:A:672:PHE:CE2	2.45	0.52
1:C:670:THR:OG1	1:F:669:GLU:OE1	2.26	0.51
1:F:678:LEU:O	1:F:681:THR:HG23	2.10	0.51
1:C:668:THR:HG23	1:C:672:PHE:CE2	2.47	0.50
1:E:702:TYR:HB3	1:E:704:MSE:HE3	1.93	0.50
1:E:668:THR:HG23	1:E:672:PHE:CE2	2.47	0.49
1:A:702:TYR:HB3	1:A:704:MSE:HE1	1.96	0.48
1:B:700:THR:HG22	1:B:748:GLU:HG2	1.96	0.47
1:A:660:VAL:HG21	1:A:661:GLU:N	2.29	0.47
1:A:682:LYS:HG3	1:A:686:LEU:HD12	1.97	0.47
1:D:689:LYS:HB3	1:D:693:LYS:NZ	2.30	0.46
1:B:749:ASP:C	1:B:750:ARG:HG2	2.37	0.45
1:F:668:THR:HG23	1:F:672:PHE:CE2	2.52	0.45
1:C:675:PHE:HA	1:C:684:GLN:HG3	1.99	0.45
1:A:737:LYS:HG3	1:A:740:PHE:HB2	1.98	0.45
1:A:738:GLU:N	1:A:738:GLU:OE1	2.50	0.45
1:B:668:THR:HG23	1:B:672:PHE:CE2	2.52	0.45
1:A:720:GLU:OE2	1:A:724:TYR:OH	2.35	0.44
1:B:726:TYR:CE1	1:B:737:LYS:HG3	2.53	0.44
1:F:694:GLN:O	1:F:695:SER:CB	2.65	0.44
1:D:656:HIS:CE1	1:D:658:ASP:CB	3.00	0.44
1:C:656:HIS:CB	1:C:658:ASP:OD1	2.67	0.43
1:F:699:HIS:CE1	1:F:702:TYR:HD1	2.37	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:749:ASP:O	1:B:750:ARG:CG	2.67	0.43
1:A:669:GLU:OE1	1:D:670:THR:OG1	2.36	0.43
1:E:674:ILE:HG23	1:E:684:GLN:HA	2.01	0.42
1:F:699:HIS:CE1	1:F:746:TYR:CB	3.03	0.41
1:A:694:GLN:O	1:A:695:SER:HB2	2.20	0.41
1:D:675:PHE:HA	1:D:684:GLN:HG3	2.03	0.41
1:C:662:PHE:C	1:C:662:PHE:CD1	2.93	0.41
1:F:692:LYS:NZ	1:F:748:GLU:OE1	2.54	0.41
1:B:743:GLU:HB2	1:B:746:TYR:CD2	2.55	0.41
1:B:678:LEU:O	1:B:681:THR:HG23	2.21	0.41
1:E:736:LYS:NZ	1:F:717:ASP:OD2	2.50	0.41
1:C:656:HIS:O	1:C:662:PHE:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	88/102 (86%)	84 (96%)	4 (4%)	0	100	100
1	B	95/102 (93%)	92 (97%)	3 (3%)	0	100	100
1	C	94/102 (92%)	93 (99%)	1 (1%)	0	100	100
1	D	94/102 (92%)	91 (97%)	3 (3%)	0	100	100
1	E	89/102 (87%)	87 (98%)	2 (2%)	0	100	100
1	F	91/102 (89%)	89 (98%)	2 (2%)	0	100	100
All	All	551/612 (90%)	536 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of

similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	81/88 (92%)	81 (100%)	0	100	100
1	B	85/88 (97%)	84 (99%)	1 (1%)	82	93
1	C	83/88 (94%)	82 (99%)	1 (1%)	82	93
1	D	85/88 (97%)	85 (100%)	0	100	100
1	E	81/88 (92%)	80 (99%)	1 (1%)	82	93
1	F	82/88 (93%)	82 (100%)	0	100	100
All	All	497/528 (94%)	494 (99%)	3 (1%)	92	98

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	742	PHE
1	C	742	PHE
1	E	742	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	707	GLN
1	B	694	GLN
1	B	707	GLN
1	D	656	HIS
1	D	707	GLN
1	E	707	GLN
1	F	699	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	90/102 (88%)	-0.02	2 (2%) 59 57	39, 63, 104, 124	0
1	B	97/102 (95%)	0.07	2 (2%) 60 58	40, 65, 115, 135	0
1	C	96/102 (94%)	0.23	4 (4%) 35 32	42, 70, 127, 163	0
1	D	96/102 (94%)	0.17	4 (4%) 35 32	38, 63, 109, 145	0
1	E	91/102 (89%)	0.01	1 (1%) 77 77	39, 58, 100, 145	0
1	F	93/102 (91%)	0.33	3 (3%) 45 43	41, 67, 135, 157	0
All	All	563/612 (91%)	0.13	16 (2%) 50 48	38, 65, 117, 163	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	658	ASP	3.9
1	C	660	VAL	3.6
1	C	659	THR	3.5
1	C	655	ALA	3.2
1	D	656	HIS	3.2
1	B	659	THR	2.8
1	D	686	LEU	2.4
1	D	725	ILE	2.3
1	F	680	GLY	2.3
1	F	750	ARG	2.3
1	A	668	THR	2.2
1	D	660	VAL	2.2
1	F	703	MSE	2.2
1	B	723	THR	2.0
1	A	714	THR	2.0
1	E	725	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.