



wwPDB X-ray Structure Validation Summary Report

Nov 21, 2014 – 02:52 AM EST

PDB ID : 4C0S
Title : Mammalian translation elongation factor eEF1A2
Authors : Crepin, T.; Shalak, V.F.; Yaremchuk, A.D.; Vlasenko, D.O.; McCarthy, A.A.;
Negrutskii, B.S.; Tukalo, M.A.; El'skaya, A.V.
Deposited on : 2013-08-07
Resolution : 2.70 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.16 November 2013
Xtriage (Phenix)	:	dev-1439
EDS	:	stable24195
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.1.3
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable24195

1 Overall quality at a glance

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1557 (2.70-2.70)
Clashscore	79885	1939 (2.70-2.70)
Ramachandran outliers	78287	1905 (2.70-2.70)
Sidechain outliers	78261	1905 (2.70-2.70)
RSRZ outliers	66119	1559 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	463	
1	B	463	

2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 6951 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ELONGATION FACTOR 1-ALPHA 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	451	Total	C	N	O	P	S	0	0	0
			3470	2196	600	656	2	16			
1	B	442	Total	C	N	O	P	S	0	0	0
			3414	2162	589	645	2	16			

- Molecule 2 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
2	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Mg 1	0	0
3	A	1	Total 1	Mg 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total 3	O 3	0	0
4	B	6	Total 6	O 6	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	135.37Å 135.37Å 304.62Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 2.70 24.81 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.0 (25.00-2.70) 99.0 (24.81-2.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.72Å)	Xtriage
Refinement program	REFMAC 5.6.0114	Depositor
R, R_{free}	0.202 , 0.256 0.203 , 0.255	Depositor DCC
R_{free} test set	2299 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	58.8	Xtriage
Anisotropy	0.219	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 45468 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6951	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, TPO, MG, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	1/3517 (0.0%)	0.65	0/4751
1	B	0.84	3/3462 (0.1%)	0.73	1/4680 (0.0%)
All	All	0.79	4/6979 (0.1%)	0.69	1/9431 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	214	TRP	CD2-CE2	7.00	1.49	1.41
1	B	196	TRP	CD2-CE2	6.23	1.48	1.41
1	A	196	TRP	CD2-CE2	5.08	1.47	1.41
1	B	78	TRP	CD2-CE2	5.02	1.47	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	240	ARG	NE-CZ-NH1	5.57	123.08	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	145	VAL	Peptide
1	B	238	PRO	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3470	0	3521	120	0
1	B	3414	0	3457	117	0
2	A	28	0	12	0	0
2	B	28	0	12	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	3	0	0	0	0
4	B	6	0	0	0	0
All	All	6951	0	7002	229	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

The worst 5 of 229 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:145:VAL:HA	1:A:146:LYS:HB2	1.32	1.11
1:B:240:ARG:HG3	1:B:240:ARG:HH11	1.23	1.02
1:B:147:GLN:NE2	1:B:236:LEU:H	1.58	1.00
1:B:102:MET:HE2	1:B:107:SER:HB2	1.49	0.94
1:A:281:ALA:HB2	1:A:334:PRO:O	1.69	0.92

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/463 (96%)	384 (86%)	48 (11%)	13 (3%)	7	16
1	B	438/463 (95%)	397 (91%)	31 (7%)	10 (2%)	10	24
All	All	883/926 (95%)	781 (88%)	79 (9%)	23 (3%)	8	20

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	317	VAL
1	A	348	ASN
1	A	448	ALA
1	A	452	THR
1	A	453	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	376/383 (98%)	325 (86%)	51 (14%)	5	13
1	B	371/383 (97%)	319 (86%)	52 (14%)	5	12
All	All	747/766 (98%)	644 (86%)	103 (14%)	5	13

5 of 103 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	392	LYS
1	B	36	LYS
1	B	371	LYS
1	A	401	ILE
1	B	6	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	132	GLN
1	B	136	HIS
1	B	314	ASN
1	B	26	HIS
1	B	343	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	A	163	1	9,9,10	5.25	4 (44%)	10,12,14	2.74	4 (40%)
1	TPO	A	239	1	10,10,11	6.21	1 (10%)	12,14,16	1.65	2 (16%)
1	SEP	B	163	1	9,9,10	6.07	2 (22%)	10,12,14	2.27	3 (30%)
1	TPO	B	239	1	10,10,11	5.52	1 (10%)	12,14,16	3.21	4 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	163	1	-	0/6/8/10	0/0/0/0
1	TPO	A	239	1	-	0/9/11/13	0/0/0/0
1	SEP	B	163	1	-	0/6/8/10	0/0/0/0
1	TPO	B	239	1	-	0/9/11/13	0/0/0/0

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	239	TPO	O-C	19.37	1.24	1.11
1	B	163	SEP	O-C	17.62	1.23	1.11
1	B	239	TPO	O-C	17.24	1.23	1.11
1	A	163	SEP	O-C	14.67	1.21	1.11
1	A	163	SEP	P-OG	4.39	1.75	1.60

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	239	TPO	CB-CA-N	8.97	127.14	110.01
1	B	163	SEP	P-OG-CB	6.31	134.77	118.63
1	A	163	SEP	P-OG-CB	5.96	133.88	118.63
1	B	239	TPO	OG1-CB-CA	5.04	117.33	105.79
1	A	163	SEP	OG-CB-CA	4.03	114.10	108.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GDP	A	500	3	30,30,30	1.50	5 (16%)	45,47,47	3.41	10 (22%)
2	GDP	B	500	3	30,30,30	1.67	6 (20%)	45,47,47	2.12	10 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GDP	A	500	3	-	0/16/32/32	0/3/3/3
2	GDP	B	500	3	-	0/16/32/32	0/3/3/3

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	GDP	C4-N9	-4.15	1.31	1.37
2	A	500	GDP	O4'-C1'	3.56	1.45	1.41
2	A	500	GDP	C4-N9	-3.46	1.32	1.37
2	B	500	GDP	C5-N7	-3.24	1.34	1.38
2	B	500	GDP	C2'-C1'	-3.21	1.48	1.53

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	GDP	C6-C5-N7	18.81	136.67	134.14
2	B	500	GDP	C6-C5-N7	7.40	135.14	134.14
2	A	500	GDP	C5-C4-N3	-6.34	118.77	126.07
2	A	500	GDP	C2-N3-C4	5.52	121.92	115.30
2	B	500	GDP	C6-N1-C2	5.22	123.15	120.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	451/463 (97%)	0.06	16 (3%) 42 47	30, 66, 114, 149	0
1	B	442/463 (95%)	-0.20	5 (1%) 77 82	29, 49, 86, 111	0
All	All	893/926 (96%)	-0.07	21 (2%) 56 62	29, 56, 107, 149	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	53	SER	4.0
1	A	430	ARG	3.4
1	A	258	GLY	3.1
1	B	258	GLY	3.0
1	A	51	LYS	2.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	SEP	A	163	10/11	0.18	1.09	41,49,76,84	0
1	TPO	A	239	11/12	0.24	0.58	117,125,150,152	0
1	TPO	B	239	11/12	0.21	0.28	75,88,129,139	0
1	SEP	B	163	10/11	0.16	-0.02	42,69,83,95	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GDP	B	500	28/28	0.12	-0.70	30,44,59,66	0
2	GDP	A	500	28/28	0.11	-0.95	31,41,49,49	0
3	MG	A	501	1/1	0.16	-	49,49,49,49	0
3	MG	B	501	1/1	0.20	-	53,53,53,53	0

6.5 Other polymers ⓘ

There are no such residues in this entry.