



Full wwPDB X-ray Structure Validation Report

Aug 22, 2014 – 12:47 AM EDT

PDB ID : 4C18
Title : The structure of the Tsi2 dimer with a disulfide bond
Authors : Robb, C.S.; Nano, F.E.; Boraston, A.B.
Deposited on : 2013-08-10
Resolution : 1.49 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable23489
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable23489

1 Overall quality at a glance

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| R_{free} | 66092 | 1513 (1.50-1.50) |
| Clashscore | 79885 | 1768 (1.50-1.50) |
| Ramachandran outliers | 78287 | 1720 (1.50-1.50) |
| Sidechain outliers | 78261 | 1718 (1.50-1.50) |
| RSRZ outliers | 66119 | 1514 (1.50-1.50) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 98 | |
| 1 | B | 98 | |

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1241 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TSI2.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|----|-----|---|---------|---------|-------|
| 1 | A | 76 | Total | C | N | O | S | 0 | 1 | 0 |
| | | | 574 | 362 | 97 | 112 | 3 | | | |
| 1 | B | 75 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 579 | 360 | 98 | 118 | 3 | | | |

There are 42 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| A | -20 | MET | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -19 | GLY | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -18 | SER | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -17 | SER | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -16 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -15 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -14 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -13 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -12 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -11 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -10 | SER | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -9 | SER | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -8 | GLY | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -7 | LEU | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -6 | VAL | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -5 | PRO | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -4 | ARG | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -3 | GLY | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -2 | SER | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | -1 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| A | 0 | MET | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -20 | MET | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -19 | GLY | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -18 | SER | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -17 | SER | - | EXPRESSION TAG | UNP Q9I0D9 |

Continued on next page...

Continued from previous page...

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| B | -16 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -15 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -14 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -13 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -12 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -11 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -10 | SER | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -9 | SER | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -8 | GLY | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -7 | LEU | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -6 | VAL | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -5 | PRO | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -4 | ARG | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -3 | GLY | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -2 | SER | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | -1 | HIS | - | EXPRESSION TAG | UNP Q9I0D9 |
| B | 0 | MET | - | EXPRESSION TAG | UNP Q9I0D9 |

- Molecule 2 is water.

| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|------------------|---------|---------|
| 2 | A | 48 | Total O 48 48 | 0 | 0 |
| 2 | B | 40 | Total O 40 40 | 0 | 0 |

3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TSI2

Chain A: 



- Molecule 1: TSI2

Chain B: 



4 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 1 | Depositor |
| Cell constants a, b, c, α , β , γ | 27.97Å 36.30Å 37.86Å 107.37° 95.23° 92.11° | Depositor |
| Resolution (Å) | 34.59 – 1.49 34.57 – 1.49 | Depositor EDS |
| % Data completeness (in resolution range) | 91.9 (34.59-1.49) 91.9 (34.57-1.49) | Depositor EDS |
| R_{merge} | 0.11 | Depositor |
| R_{sym} | (Not available) | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 3.23 (at 1.50Å) | Xtriage |
| Refinement program | REFMAC 5.7.0029 | Depositor |
| R, R_{free} | 0.223 , 0.259 0.230 , 0.269 | Depositor DCC |
| R_{free} test set | 1076 reflections (5.39%) | DCC |
| Wilson B-factor (Å ²) | 15.9 | Xtriage |
| Anisotropy | 0.710 | Xtriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.39 , 38.3 | EDS |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| L-test for twinning | $\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$ | Xtriage |
| Outliers | 0 of 21050 reflections | Xtriage |
| F_o, F_c correlation | 0.94 | EDS |
| Total number of atoms | 1241 | wwPDB-VP |
| Average B, all atoms (Å ²) | 24.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|-------------|-------------|---------------|
| | | RMSZ | $\# Z > 5$ | RMSZ | $\# Z > 5$ |
| 1 | A | 0.89 | 0/585 | 0.92 | 1/797 (0.1%) |
| 1 | B | 0.91 | 0/586 | 0.99 | 3/797 (0.4%) |
| All | All | 0.90 | 0/1171 | 0.96 | 4/1594 (0.3%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | B | 0 | 1 |

There are no bond length outliers.

All (4) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-----------|-------|-------------|----------|
| 1 | B | 70 | ARG | NE-CZ-NH1 | 6.53 | 123.56 | 120.30 |
| 1 | B | 20 | ARG | NE-CZ-NH2 | -5.86 | 117.37 | 120.30 |
| 1 | B | 50 | ASP | CB-CG-OD1 | 5.36 | 123.12 | 118.30 |
| 1 | A | 45 | ASP | CB-CG-OD2 | -5.35 | 113.48 | 118.30 |

There are no chirality outliers.

All (1) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|---------|
| 1 | B | -1 | HIS | Peptide |

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens

added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 574 | 0 | 559 | 3 | 0 |
| 1 | B | 579 | 0 | 562 | 1 | 0 |
| 2 | A | 48 | 0 | 0 | 2 | 0 |
| 2 | B | 40 | 0 | 0 | 0 | 0 |
| All | All | 1241 | 0 | 1121 | 3 | 0 |

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

All (3) close contacts within the same asymmetric unit are listed below.

| Atom-1 | Atom-2 | Distance(Å) | Clash(Å) |
|-----------------|----------------|-------------|----------|
| 1:A:18:ARG:NH1 | 2:A:2022:HOH:O | 1.95 | 0.99 |
| 1:A:18:ARG:NH2 | 2:A:2021:HOH:O | 2.26 | 0.68 |
| 1:A:18:ARG:HH21 | 1:B:17:ALA:CB | 2.29 | 0.46 |

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-----------|---------|----------|-------------|-----|
| 1 | A | 75/98 (76%) | 75 (100%) | 0 | 0 | 100 | 100 |
| 1 | B | 73/98 (74%) | 71 (97%) | 2 (3%) | 0 | 100 | 100 |
| All | All | 148/196 (76%) | 146 (99%) | 2 (1%) | 0 | 100 | 100 |

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|-------------|-----|
| 1 | A | 57/81 (70%) | 56 (98%) | 1 (2%) | 71 | 38 |
| 1 | B | 60/81 (74%) | 60 (100%) | 0 | 100 | 100 |
| All | All | 117/162 (72%) | 116 (99%) | 1 (1%) | 87 | 69 |

All (1) residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 18 | ARG |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 57 | GLN |
| 1 | A | 61 | GLN |

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|---------------|--------|-----------------------------|-----------------------|--------|
| 1 | A | 76/98 (77%) | 0.72 | 9 (11%) 5 5 | 15, 23, 46, 62 | 0 |
| 1 | B | 75/98 (76%) | 0.47 | 1 (1%) 74 81 | 15, 21, 31, 47 | 1 (1%) |
| All | All | 151/196 (77%) | 0.60 | 10 (6%) 18 19 | 15, 22, 42, 62 | 1 (0%) |

All (10) RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1 | A | 29 | ASP | 6.3 |
| 1 | A | 75 | PRO | 4.7 |
| 1 | B | -1 | HIS | 4.5 |
| 1 | A | 27 | GLN | 3.8 |
| 1 | A | 24 | ALA | 3.7 |
| 1 | A | 28 | ASN | 3.4 |
| 1 | A | 69 | ASP | 2.3 |
| 1 | A | 30 | ASP | 2.3 |
| 1 | A | 70 | ARG | 2.3 |
| 1 | A | 74 | GLU | 2.1 |

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.