



wwPDB X-ray Structure Validation Summary Report

Aug 15, 2014 – 09:30 AM EDT

PDB ID : 4C1N
Title : Corrinoid protein reactivation complex with activator
Authors : Hennig, S.E.; Goetzl, S.; Jeoung, J.H.; Bommer, M.; Lendzian, F.; Hildebrandt, P.; Dobbek, H.
Deposited on : 2013-08-13
Resolution : 2.53 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable23489
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable23489

1 Overall quality at a glance

The reported resolution of this entry is 2.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3240 (2.54-2.50)
Clashscore	79885	4080 (2.54-2.50)
Ramachandran outliers	78287	3990 (2.54-2.50)
Sidechain outliers	78261	3992 (2.54-2.50)
RSRZ outliers	66119	3241 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	442	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
1	C	442	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
1	E	442	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
1	G	442	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
2	B	309	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
2	D	309	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
2	F	309	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
2	H	309	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
3	I	509	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
3	J	509	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
3	X	509	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>
4	K	510	<div><div></div><div><div></div><div></div><div></div><div></div></div></div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 40155 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CARBON MONOXIDE DEHYDROGENASE CORRINOID/IRON-SULFURPROTEIN, GAMMA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	442	Total	C	N	O	S	0	0	0
			3389	2186	559	634	10			
1	C	442	Total	C	N	O	S	0	0	0
			3389	2186	559	634	10			
1	E	442	Total	C	N	O	S	0	0	0
			3389	2186	559	634	10			
1	G	441	Total	C	N	O	S	0	0	0
			3385	2184	558	633	10			

- Molecule 2 is a protein called CO DEHYDROGENASE/ACETYL-COASYNTHASE, IRON-SULFUR PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	309	Total	C	N	O	S	0	0	0
			2368	1518	393	441	16			
2	D	309	Total	C	N	O	S	0	0	0
			2368	1518	393	441	16			
2	F	309	Total	C	N	O	S	0	0	0
			2368	1518	393	441	16			
2	H	309	Total	C	N	O	S	0	0	0
			2368	1518	393	441	16			


- Molecule 3 is a protein called IRON-SULFUR CLUSTER BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	I	509	Total	C	N	O	S	1	0	0
			3882	2474	652	740	16			
3	J	509	Total	C	N	O	S	1	0	0
			3882	2474	652	740	16			
3	X	509	Total	C	N	O	S	0	0	0
			3882	2474	652	740	16			

- Molecule 4 is a protein called IRON-SULFUR CLUSTER BINDING PROTEIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	K	510	Total	C	N	O	S	0	0	0
			3891	2479	653	743	16			

- Molecule 5 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe_4S_4).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	Fe	S	0	0
			8	4	4		
5	C	1	Total	Fe	S	0	0
			8	4	4		
5	E	1	Total	Fe	S	0	0
			8	4	4		
5	G	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 6 is COBALAMIN (three-letter code: B12) (formula: $\text{C}_{62}\text{H}_{89}\text{CoN}_{13}\text{O}_{14}\text{P}$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	C	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		
6	I	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		
6	K	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		
6	X	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	119	Total	O	0	0
			119	119		
7	B	51	Total	O	0	0
			51	51		
7	C	116	Total	O	0	0
			116	116		
7	D	49	Total	O	0	0
			49	49		
7	E	120	Total	O	0	0
			120	120		
7	F	69	Total	O	0	0
			69	69		
7	G	80	Total	O	0	0
			80	80		
7	H	43	Total	O	0	0
			43	43		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	I	157	Total 157	O 157	0	0
7	J	130	Total 130	O 130	0	0
7	K	121	Total 121	O 121	0	0
7	X	143	Total 143	O 143	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CARBON MONOXIDE DEHYDROGENASE CORRINOID/IRON-SULFURPROTEIN, GAMMA SUBUNIT

Chain A:




- Molecule 1: CARBON MONOXIDE DEHYDROGENASE CORRINOID/IRON-SULFURPROTEIN, GAMMA SUBUNIT

Chain C:



- Molecule 1: CARBON MONOXIDE DEHYDROGENASE CORRINOID/IRON-SULFURPROTEIN, GAMMA SUBUNIT


Chain E:






- Molecule 1: CARBON MONOXIDE DEHYDROGENASE CORRINOID/IRON-SULFURPROTEIN, GAMMA SUBUNIT

Chain G:





- Molecule 2: CO DEHYDROGENASE/ACETYL-COASYNTHASE, IRON-SULFUR PROTEIN

Chain B:



- Molecule 2: CO DEHYDROGENASE/ACETYL-COASYNTHASE, IRON-SULFUR PROTEIN

Chain D:




- Molecule 2: CO DEHYDROGENASE/ACETYL-COASYNTHASE, IRON-SULFUR PROTEIN

Chain F:




- Molecule 2: CO DEHYDROGENASE/ACETYL-COASYNTHASE, IRON-SULFUR PROTEIN

Chain H:




- Molecule 3: IRON-SULFUR CLUSTER BINDING PROTEIN

Chain I:




- Molecule 3: IRON-SULFUR CLUSTER BINDING PROTEIN

Chain J:



- Molecule 3: IRON-SULFUR CLUSTER BINDING PROTEIN


Chain X:



- Molecule 4: IRON-SULFUR CLUSTER BINDING PROTEIN

Chain K:





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	67.13Å 128.18Å 163.37Å 77.61° 82.25° 88.76°	Depositor
Resolution (Å)	30.63 – 2.53 34.29 – 2.53	Depositor EDS
% Data completeness (in resolution range)	96.0 (30.63-2.53) 96.0 (34.29-2.53)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.54Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.191 , 0.243 0.191 , 0.241	Depositor DCC
R_{free} test set	8446 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 23.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	0 of 168978 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	40155	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, B12

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	1/3459 (0.0%)	0.44	1/4703 (0.0%)
1	C	0.35	0/3459	0.44	0/4703
1	E	0.50	1/3459 (0.0%)	0.48	2/4703 (0.0%)
1	G	0.33	0/3455	0.44	0/4698
2	B	0.39	0/2419	0.43	0/3289
2	D	0.39	0/2419	0.42	0/3289
2	F	0.28	0/2419	0.42	0/3289
2	H	0.22	0/2419	0.40	0/3289
3	I	0.28	0/3944	0.42	1/5342 (0.0%)
3	J	0.41	0/3944	0.45	1/5342 (0.0%)
3	X	0.38	0/3944	0.44	0/5342
4	K	0.41	2/3953 (0.1%)	0.45	2/5354 (0.0%)
All	All	0.37	4/39293 (0.0%)	0.44	7/53343 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	K	310	PRO	N-CD	5.25	1.55	1.47
1	A	202	PRO	N-CD	5.23	1.55	1.47
1	E	429	PRO	N-CD	5.16	1.55	1.47
4	K	628	PRO	N-CD	5.01	1.54	1.47

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	J	156	GLU	N-CA-C	-8.68	87.57	111.00
1	E	382	THR	C-N-CD	5.68	140.33	128.40
1	E	428	GLY	C-N-CD	5.63	140.23	128.40
3	I	123	ASP	C-N-CD	5.56	140.07	128.40
4	K	627	PHE	C-N-CD	5.50	139.94	128.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3389	0	3467	57	0
1	C	3389	0	3467	58	0
1	E	3389	0	3467	54	0
1	G	3385	0	3464	79	0
2	B	2368	0	2402	39	0
2	D	2368	0	2402	34	0
2	F	2368	0	2402	44	0
2	H	2368	0	2402	37	0
3	I	3882	0	3973	56	0
3	J	3882	0	3973	57	0
3	X	3882	0	3973	57	0
4	K	3891	0	3979	49	0
5	A	8	0	0	1	0
5	C	8	0	0	0	0
5	E	8	0	0	0	0
5	G	8	0	0	1	0
6	C	91	0	88	9	0
6	I	91	0	88	11	0
6	K	91	0	88	5	0
6	X	91	0	88	5	0
7	A	119	0	0	10	0
7	B	51	0	0	6	0
7	C	116	0	0	11	0
7	D	49	0	0	5	0
7	E	120	0	0	17	0
7	F	69	0	0	5	0
7	G	80	0	0	18	1
7	H	43	0	0	4	0
7	I	157	0	0	13	0
7	J	130	0	0	19	0
7	K	121	0	0	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	X	143	0	0	15	1
All	All	40155	0	39723	605	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

The worst 5 of 605 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:G:46:SER:O	7:G:2006:HOH:O	1.59	1.17
1:G:284:ALA:O	7:G:2051:HOH:O	1.70	1.06
1:E:367:THR:O	1:E:380:LYS:NZ	1.90	1.04
1:A:86:LYS:HD2	1:A:87:ARG:HG2	1.42	1.01
7:J:2066:HOH:O	3:X:169:ASP:HB3	1.60	0.99

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
7:G:2071:HOH:O	7:X:2105:HOH:O[1_645]	1.99	0.21

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	440/442 (100%)	422 (96%)	18 (4%)	0	100	100
1	C	440/442 (100%)	424 (96%)	16 (4%)	0	100	100
1	E	440/442 (100%)	425 (97%)	15 (3%)	0	100	100
1	G	439/442 (99%)	411 (94%)	28 (6%)	0	100	100
2	B	307/309 (99%)	294 (96%)	13 (4%)	0	100	100
2	D	307/309 (99%)	298 (97%)	9 (3%)	0	100	100
2	F	307/309 (99%)	297 (97%)	10 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	307/309 (99%)	296 (96%)	11 (4%)	0	100	100
3	I	507/509 (100%)	492 (97%)	15 (3%)	0	100	100
3	J	507/509 (100%)	493 (97%)	14 (3%)	0	100	100
3	X	507/509 (100%)	490 (97%)	17 (3%)	0	100	100
4	K	508/510 (100%)	485 (96%)	23 (4%)	0	100	100
All	All	5016/5041 (100%)	4827 (96%)	189 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	361/361 (100%)	359 (99%)	2 (1%)	92	98
1	C	361/361 (100%)	357 (99%)	4 (1%)	84	96
1	E	361/361 (100%)	355 (98%)	6 (2%)	73	92
1	G	361/361 (100%)	360 (100%)	1 (0%)	96	99
2	B	256/256 (100%)	256 (100%)	0	100	100
2	D	256/256 (100%)	253 (99%)	3 (1%)	82	95
2	F	256/256 (100%)	256 (100%)	0	100	100
2	H	256/256 (100%)	256 (100%)	0	100	100
3	I	422/422 (100%)	420 (100%)	2 (0%)	94	99
3	J	422/422 (100%)	419 (99%)	3 (1%)	91	98
3	X	422/422 (100%)	418 (99%)	4 (1%)	87	97
4	K	423/423 (100%)	418 (99%)	5 (1%)	82	95
All	All	4157/4157 (100%)	4127 (99%)	30 (1%)	91	98

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	418	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	I	127	LYS
3	X	169	ASP
1	G	50	LYS
3	I	429	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	418	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SF4	A	1444	1	12,12,12	5.15	12 (100%)	0,24,24	0.00	-
6	B12	C	1444	3	101,101,101	0.75	3 (2%)	152,166,166	1.21	15 (9%)
5	SF4	C	1445	1	12,12,12	4.53	12 (100%)	0,24,24	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SF4	E	1444	1	12,12,12	5.07	12 (100%)	0,24,24	0.00	-
5	SF4	G	1444	1	12,12,12	5.64	12 (100%)	0,24,24	0.00	-
6	B12	I	1631	3	101,101,101	0.72	3 (2%)	152,166,166	1.22	14 (9%)
6	B12	K	1631	4	101,101,101	0.73	3 (2%)	152,166,166	1.20	11 (7%)
6	B12	X	1631	3	101,101,101	0.72	3 (2%)	152,166,166	1.22	12 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SF4	A	1444	1	-	0/0/48/48	0/6/5/5
6	B12	C	1444	3	-	0/56/223/223	0/3/11/11
5	SF4	C	1445	1	-	0/0/48/48	0/6/5/5
5	SF4	E	1444	1	-	0/0/48/48	0/6/5/5
5	SF4	G	1444	1	-	0/0/48/48	0/6/5/5
6	B12	I	1631	3	-	0/56/223/223	0/3/11/11
6	B12	K	1631	4	-	0/56/223/223	0/3/11/11
6	B12	X	1631	3	-	0/56/223/223	0/3/11/11

The worst 5 of 60 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	1444	SF4	S1-FE3	-12.51	2.24	2.33
5	A	1444	SF4	S1-FE3	-9.58	2.26	2.33
5	E	1444	SF4	S1-FE3	-8.49	2.27	2.33
5	C	1445	SF4	S1-FE3	-6.67	2.28	2.33
5	G	1444	SF4	S3-FE2	-6.40	2.28	2.33

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	X	1631	B12	C35-C5-C6	5.82	127.23	118.75
6	K	1631	B12	C35-C5-C6	5.65	126.99	118.75
6	X	1631	B12	C35-C5-C4	-5.61	110.56	118.75
6	C	1444	B12	C35-C5-C6	5.58	126.88	118.75
6	I	1631	B12	C35-C5-C6	5.46	126.72	118.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/442 (100%)	-0.28	1 (0%) 93 95	21, 40, 54, 66	2 (0%)
1	C	442/442 (100%)	-0.18	2 (0%) 88 90	23, 39, 65, 86	1 (0%)
1	E	442/442 (100%)	-0.33	1 (0%) 93 95	23, 37, 51, 58	1 (0%)
1	G	441/442 (99%)	-0.10	5 (1%) 77 79	29, 47, 66, 82	0
2	B	309/309 (100%)	-0.30	1 (0%) 91 94	23, 36, 50, 67	0
2	D	309/309 (100%)	-0.32	0 100 100	24, 38, 53, 74	0
2	F	309/309 (100%)	-0.28	0 100 100	24, 36, 51, 63	0
2	H	309/309 (100%)	-0.18	0 100 100	29, 43, 60, 68	0
3	I	509/509 (100%)	-0.34	1 (0%) 93 95	24, 33, 45, 61	5 (0%)
3	J	509/509 (100%)	-0.31	0 100 100	21, 34, 47, 65	3 (0%)
3	X	509/509 (100%)	-0.32	0 100 100	23, 34, 47, 60	3 (0%)
4	K	510/510 (100%)	-0.26	0 100 100	25, 37, 53, 72	3 (0%)
All	All	5040/5041 (99%)	-0.27	11 (0%) 93 95	21, 37, 56, 86	18 (0%)

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	318	ILE	4.6
1	G	49	ALA	3.8
1	G	100	ASP	2.9
1	A	235	GLU	2.4
1	C	53	LEU	2.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	B12	X	1631	91/91	0.13	0.47	20,26,35,47	0
6	B12	K	1631	91/91	0.14	0.44	24,32,38,52	0
6	B12	C	1444	91/91	0.14	0.38	21,29,38,46	0
5	SF4	E	1444	8/8	0.14	0.31	29,38,41,43	0
5	SF4	A	1444	8/8	0.14	0.04	30,33,38,39	0
6	B12	I	1631	91/91	0.12	-0.04	17,24,31,41	0
5	SF4	G	1444	8/8	0.12	-0.70	36,42,46,47	0
5	SF4	C	1445	8/8	0.10	-1.27	43,47,58,61	0

6.5 Other polymers

There are no such residues in this entry.